

FORM PTO-100
(REV. 10-2000)

U.S. DEPARTMENT OF COMMERCE PATENT AND TRADEMARK OFFICE

ATTORNEY'S DOCKET NUMBER

P2534-3

TRANSMITTAL LETTER TO THE UNITED STATES
DESIGNATED/ELECTED OFFICE (DO/EO/US)
CONCERNING A FILING UNDER 35 U.S.C. 371

U.S. APPLICATION NO. (If known, see 37 CFR 1.5)

097869566

INTERNATIONAL APPLICATION NO.
PCT/US99/30720INTERNATIONAL FILING DATE
22 December 1999 (22.11.99)PRIORITY DATE CLAIMED
23 December 1998 (23.12.98)

TITLE OF INVENTION IL-1 RELATED POLYPEPTIDES

APPLICANT(S) FOR DO/EO/US GODDARD Audrey and PAN James

Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information.

1. ☒ This is a **FIRST** submission of items concerning a filing under 35 U.S.C. 371.
2. ☐ This is a **SECOND** or **SUBSEQUENT** submission of items concerning a filing under 35 U.S.C. 371.
3. ☒ This is an express request to promptly begin national examination procedures (35 U.S.C. 371(f)).
4. ☐ The US has been elected by the expiration of 19 months from the priority date (PCT Article 31)
5. ☒ A copy of the International Application as filed (35 U.S.C. 371(c)(2))
 - a. ☐ is attached hereto (required only if not communicated by the International Bureau).
 - b. ☐ has been communicated by the International Bureau.
 - c. ☒ is not required, as the application was filed in the United States Receiving Office (RO/US).
6. ☐ An English language translation of the International Application as filed (35 U.S.C. 371(c)(2)).
7. ☒ Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371(c)(3))
 - a. ☐ are attached hereto (required only if not communicated by the International Bureau).
 - b. ☐ have been communicated by the International Bureau.
 - c. ☐ have not been made; however, the time limit for making such amendments has NOT expired.
 - d. ☒ have not been made and will not be made.
8. ☐ An English language translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371(c)(3)).
9. ☒ An oath or declaration of the inventor(s) (35 U.S.C. 371(c)(4)).
10. ☐ An English language translation of the annexes to the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371(c)(5)).

Items 11 to 16 below concern document(s) or information included:

11. ☐ An Information Disclosure Statement under 37 CFR 1.97 and 1.98.
12. ☒ An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included.
13. ☐ A FIRST preliminary amendment.
☐ A SECOND or SUBSEQUENT preliminary amendment.
14. ☐ A substitute specification.
15. ☐ A change of power of attorney and/or address letter.
16. ☒ Other items or information:

Enclosed is the Petition for Revival Of An International Application For Patent Designating The U.S. Abandoned Unintentionally Under 37 CFR 1.137(b); fee of \$1240 for Petition for Revival (authorized to charge to Deposit Account No. 07-0630);

Enclosed is courtesy copy of International Preliminary Examination Report;

097-869566

INTERNATIONAL APPLICATION NO

PCT/US99/30720

ATTORNEY'S DOCKET NUMBER
P2534-317. ☒ The following fees are submitted:**BASIC NATIONAL FEE (37 CFR 1.492 (a) (1) - (5)):**

Neither international preliminary examination fee (37 CFR 1.482)
nor international search fee (37 CFR 1.445(a)(2)) paid to USPTO
and International Search Report not prepared by the EPO or JPO \$1000.00

International preliminary examination fee (37 CFR 1.482) not paid to
USPTO but International Search Report prepared by the EPO or JPO \$860.00

International preliminary examination fee (37 CFR 1.482) not paid to USPTO but
international search fee (37 CFR 1.445(a)(2)) paid to USPTO \$710.00

International preliminary examination fee paid to USPTO (37 CFR 1.482)
but all claims did not satisfy provisions of PCT Article 33(1)-(4) \$690.00

International preliminary examination fee paid to USPTO (37 CFR 1.482)
and all claims satisfied provisions of PCT Article 33(1)-(4) \$100.00

ENTER APPROPRIATE BASIC FEE AMOUNT =**CALCULATIONS** PTO USE ONLY

Surcharge of \$130.00 for furnishing the oath or declaration later than ☐ 20 ☒ 30
months from the earliest claimed priority date (37 CFR 1.492(e)).

\$ 130.00

CLAIMS	NUMBER FILED	NUMBER EXTRA	RATE	
Total claims	30 - 20 =	10	X \$18.00	\$ 180.00
Independent claims	4 - 3 =	1	X \$80.00	\$ 80.00
MULTIPLE DEPENDENT CLAIM(S) (if applicable)			+ \$270.00	\$

TOTAL OF ABOVE CALCULATIONS = \$ 1,250.00

☐ Applicant claims small entity status. See 37 CFR 1.27. The fees indicated above
are reduced by 1/2.

SUBTOTAL = \$ 1,250.00

Processing fee of \$130.00 for furnishing the English translation later than ☐ 20 ☐ 30
months from the earliest claimed priority date (37 CFR 1.492(f)).

\$

TOTAL NATIONAL FEE = \$ 1,250.00

Fee for recording the enclosed assignment (37 CFR 1.21(h)). The assignment must be
accompanied by an appropriate cover sheet (37 CFR 3.28, 3.31). \$40.00 per property +

\$ 40.00

TOTAL FEES ENCLOSED = \$ 1,290.00

Amount to be refunded:	\$
charged:	\$

a. ☐ A check in the amount of \$ _____ to cover the above fees is enclosed.

b. ☒ Please charge my Deposit Account No. 07-0630 in the amount of \$ 1,290.00 to cover the above fees.
A duplicate copy of this sheet is enclosed.

c. ☒ The Commissioner is hereby authorized to charge any additional fees which may be required, or credit any
overpayment to Deposit Account No. 07-0630. A duplicate copy of this sheet is enclosed.

NOTE: Where an appropriate time limit under 37 CFR 1.494 or 1.495 has not been met, a petition to revive (37 CFR 1.137(a) or (b)) must be filed and granted to restore the application to pending status.

SEND ALL CORRESPONDENCE TO:

Richard B. Love
GENENTECH, INC.
1 DNA Way
South San Francisco, CA 94080-4990

SIGNATURE


Richard B. Love

NAME

34,659

REGISTRATION NUMBER

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re Application of Goddard <i>et al.</i> Serial No.: 09/869,566 Filing Date: June 29, 2001 For: <i>IL-1 Related Polypeptides</i>	Group Art Unit: Not yet assigned Examiner: Not yet assigned Confirmation No.: 4737 EXPRESS MAIL NUMBER: EL 809449045 US DATE OF DEPOSIT: FEBRUARY 19, 2002  Glory L. Tabuena
--	---


CERTIFICATE RE: SEQUENCE LISTING
RESPONSE UNDER 37 CFR § 1.821(f) and (g)

Box MISSING PARTS
U.S. Patent and Trademark Office
P. O. Box 2327
Arlington, Virginia 22202

Sir:

I hereby state that the Sequence Listing submitted herewith is submitted in paper copy and a computer-readable diskette and that the information recorded in computer readable form is identical to the written sequence listing. I further state that this submission includes no new matter.

Respectfully submitted,
GENENTECH, INC.

By: 
Craig G. Svoboda
Reg. No. 39,044
Telephone No. (650) 225-1489

Date: February 19, 2002



09157

PATENT TRADEMARK OFFICE

IL-1 RELATED POLYPEPTIDESFIELD OF THE INVENTION

The present invention relates generally to the identification and isolation of novel DNAs having homology to interleukin-1 (IL-1) or interleukin-1 receptor antagonist (IL-1Ra) polypeptides, and to the recombinant production of novel polypeptides, designated herein as interleukin-1-like polypeptides ("IL-1lp").

BACKGROUND OF THE INVENTION

Interleukin-1 refers to two proteins (IL-1 α and IL-1 β) which play a key role early in the inflammatory response (for a review, see Dinarello, Blood, 87: 2095-2147 (1996) and references therein). both proteins are made as intracellular precursor proteins which are cleaved upon secretion to yield mature carboxy-terminal 17 kDa fragments which are biologically active. In the case of IL-1 β , this cleavage involves an intracellular cysteine protease, known as ICE, which is required to release the active fragment from the inactive precursor. The precursor of IL-1 α is active.

These two proteins act by binding to cell surface receptors found on almost all cell types and triggering a range of responses either alone or in concert with other secreted factors. These range from effects on proliferation (e.g. fibroblasts, T cells) apoptosis (e.g. A375 melanoma cells), cytokine induction (e.g. of TNF, IL-1, IL-8), receptor activation (e.g. E-selectin), eicosanoid production (e.g. PGE2) and the secretion of degradative enzymes (e.g. collagenase). To achieve these effects, IL-1 activates transcription factors such as NF-KB and AP-1. Several of the activities of IL-1 action on target cells are believed to be mediated through activation of kinase cascades that have also been associated with cellular stresses, such as the stress activated MAP kinase JNK/SAPK and p38.

A third member of the IL-1 family was subsequently discovered which acts as a natural antagonist of IL-1 α and IL-1 β by binding to the IL-1 receptor but not transducing an intracellular signal or a biological response. The protein is called IL-1Ra (for IL-1 receptor antagonist) or IRAP (for IL-1 receptor antagonist protein). At least three alternatively spliced forms of IL-1Ra exist: one encodes a secreted protein, also known as secretory IL-1Ra ("sIL-1Ra") (described in Eisenberg et al., Nature, 343: 341-346 (1990)), and the other two encode intracellular proteins. IL-1 α , IL-1 β and IL-1Ra exhibit approximately 25-30% sequence identity with each other and share a similar three dimensional structure consisting of twelve β -strands folded into a β -barrel, with an internal thrice repeated structural motif.

There are three known IL-1 receptor subunits. The active receptor complex consists of the type I receptor and IL-1 accessory protein (IL-1RAcP). The type I receptor is responsible for binding of the IL-1 α , IL-1 β and IL-1Ra ligands, and is able to do so in the absence of the IL-1RAcP. However, signal transduction requires the interaction of IL-1 α or IL-1 β with the IL-1RAcP. IL-1Ra does not interact with the IL-1RAcP and hence cannot induce signal transduction. A third receptor subunit, the type II receptor, binds IL-1 α and IL-1 β but cannot transduce signal due its lack of an intracellular domain. Instead, the type II receptor either acts as a decoy in its membrane bound form or as an IL-1 antagonist in a processed, secreted form, and hence inhibits IL-1 activity. The type II receptor weakly binds to IL-1Ra.

Many studies using IL-1Ra, soluble IL-1R derived from the extracellular domain of the type I IL-1 receptor, antibodies to IL-1 α or IL-1 β , and transgenic knockout mice for these genes have shown that IL-1 plays a role in a number of pathophysiologies (for a review, see Dinarello, Blood, 87: 2095-2147 (1996)). For example, IL-1Ra has been shown to be effective in animal models of septic shock, rheumatoid arthritis, graft-versus-host disease (GVHD), stroke, cardiac ischemia, psoriasis, inflammatory bowel disease, and asthma. In addition, IL-1Ra has demonstrated efficacy in clinical trials for rheumatoid arthritis and GVHD, and is also in clinical trials for inflammatory bowel disease, asthma and psoriasis.

More recently, interleukin-18 (IL-18) was placed in the IL-1 family (for a review, see Dinarello et al, J. Leukocyte Biol., 63: 658-664 (1998)). IL-18 shares the β -pleated, barrel-like form of IL-1 α and IL-1 β . In addition, IL-18 is the natural ligand for the IL-1 receptor family member formerly known as IL-1R-related protein (IL-1Rrp) (now known as the IL-18 receptor (IL-18R)). IL-18 has been shown to initiate the inflammatory cytokine cascade in a mixed population of peripheral blood mononuclear cells (PBMCs) by triggering the constitutive IL-18 receptors on lymphocytes and NK cells, inducing TNF production in the activated cells. TNF, in turn, stimulates IL-1 and IL-8 production in CD14+ cells. Because of its ability to induce TNF, IL-1, and both C-C and C-X-C chemokines, and because IL-18 induces Fas ligand as well as nuclear translocation of nuclear factor κ B (NF- κ B), IL-18 ranks with other pro-inflammatory cytokines as a likely contributor to systemic and local inflammation.

SUMMARY OF THE INVENTION

A family of cDNA clones (DNA85066, DNA96786, DNA94618, DNA102043, DNA114876, DNA102044, DNA92929, DNA96787, and DNA92505) has been identified, having homology to interleukin-1, that encode novel polypeptides. The novel polypeptides and variants thereof are collectively designated in the present application as "interleukin-1-like polypeptides" or "IL-1lp", as further defined herein. Accordingly, one aspect of the invention is an isolated IL-1lp polypeptide.

In another embodiment, the invention provides an isolated nucleic acid molecule encoding an IL-1lp polypeptide.

In another embodiment, the invention provides a method for producing an IL-1lp comprising culturing a host cell comprising a heterologous nucleic acid sequence encoding an IL-1lp polypeptide, under conditions wherein the IL-1lp polypeptide is expressed, and recovering the IL-1lp polypeptide from the host cell.

In another embodiment, the invention provides an anti-IL-1lp antibody.

In another embodiment, the invention provides chimeric molecules comprising an IL-1lp polypeptide fused to a heterologous polypeptide or amino acid sequence. An example of such a chimeric molecule comprises an IL-1lp polypeptide fused to an epitope tag sequence or a Fc region of an immunoglobulin.

In another embodiment, the invention provides an antibody which specifically binds to an IL-1lp polypeptide. Optionally, the antibody is a monoclonal antibody.

M 27.07.00

In yet another embodiment, the invention concerns agonists and antagonists of a native IL-1p polypeptide. In a particular embodiment, the agonist or antagonist is an anti-IL-1p antibody.

In a further embodiment, the invention concerns a method of identifying agonists or antagonists of a native IL-1p polypeptide, by contacting the native IL-1p polypeptide with a candidate molecule and monitoring a biological activity mediated by said polypeptide.

In a still further embodiment, the invention concerns a composition comprising an IL-1p polypeptide, or an agonist or antagonist as hereinabove defined, in combination with a pharmaceutically acceptable carrier.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 shows a nucleotide sequence (SEQ ID NO:1) and derived amino acid sequences (SEQ ID NOS:2-3) related to a native sequence hIL-1Ra1. The nucleotide sequence (SEQ ID NO:1) contains an intron believed to extend from nucleotide positions 181 to 432, with a splice donor site at nucleotide positions 181 to 186 and splice acceptor site at nucleotide positions 430 to 432. The amino acid sequences (SEQ ID NOS:2 and 3) are derived from the exonic sequences that are believed to make up the processed (intron-free) coding sequence.

Figure 2 shows the nucleotide sequence (SEQ ID NO:4) and derived amino acid sequence (SEQ ID NO:5) of a native sequence hIL-1Ra1 polypeptide fused at its N-terminus to a heterologous signal peptide (amino acid positions 1-15), flag peptide affinity handle (amino acid positions 16-23) and peptide linker (amino acid positions 24-36).

Figure 3 shows the nucleotide sequence (SEQ ID NO:6) and derived amino acid sequence (SEQ ID NO:7) of a native sequence hIL-1Ra1 polypeptide. The nucleotide sequence (SEQ ID NO:6) and derived amino acid sequence (SEQ ID NO:7) are believed to represent the processed (intron-free) form and intact hIL-1Ra1 polypeptide, respectively, of the nucleotide sequence (SEQ ID NO:1) and amino acid sequences (SEQ ID NOS:2-3) of Figure 1. The start and stop codons in the coding sequence are located at nucleotide positions 103-105 and 682-684, respectively. The putative signal sequence extends from amino acid positions 1 to 14. A putative cAMP- and cGMP-dependent protein kinase phosphorylation site is located at amino acid positions 33-36. Putative N-myristoylation sites are located at amino acid positions 50-55 and 87-92.

Figure 4 shows a nucleotide sequence (SEQ ID NO:8) of which nucleotides 145-629 correspond to EST AI014548.

Figure 5 shows the nucleotide sequence (SEQ ID NO:9) and derived amino acid sequence (SEQ ID NO:10) of a native sequence hIL-1Ra2 polypeptide. The start and stop codons in the coding sequence are located at nucleotide positions 96-98 and 498-500, respectively. The putative signal sequence extends from amino acid positions 1-26.

Figure 6 shows the nucleotide sequence (SEQ ID NO:11) of EST I433156.

Figure 7 shows the nucleotide sequence (SEQ ID NO:12) and derived amino acid sequence (SEQ ID NO:13) of a native sequence hIL-1Ra3 polypeptide. The start and stop codons in the coding sequence are located at nucleotide positions 1-3 and 466-468, respectively. The putative signal sequence extends from amino acid positions 1-33. Putative N-myristoylation sites are located at amino acid positions 29-34, 30-35, 60-65, 63-68, 73-78, 91-96 and 106-111. An interleukin-1-like sequence is located at amino acid positions 111-131.

Figure 8 shows the nucleotide sequence (SEQ ID NO:14) of EST 5120028.

Figure 9 shows the nucleotide sequence (SEQ ID NO:15) and derived amino acid sequence (SEQ ID NO:16) of a native sequence mIL-1Ra3 polypeptide. The start and stop codons in the coding sequence are located at nucleotide positions 145-147 and 610-612, respectively.

5 The putative signal sequence extends from amino acid positions 1-33. Putative N-myristoylation sites are located at amino acid positions 29-34, 60-65, 63-68, 91-96 and 106-111. An interleukin-1-like sequence is located at amino acid positions 111-131.

Figure 10 shows the nucleotide sequence (SEQ ID NO:17) of EST W08205.

10 Figure 11 is an autoradiograph of Northern blots depicting expression of hIL-1Ra3 mRNA in placental tissue and expression of mIL-1Ra3 mRNA in day-17 mouse embryo tissue.

Figure 12 is an amino acid sequence alignment of native sequence hIL-1Ra1L (SEQ ID NO:19), hIL-1Ra1V (SEQ ID NO:25), hIL-1Ra1S (SEQ ID NO:21), hIL-1Ra2 (SEQ ID NO:10), hIL-1Ra3 (SEQ ID NO:13) and mIL-1Ra3 (SEQ ID NO:16) polypeptides with secretory hIL-1Ra (also referred to as "sIL-1Ra" and "hIL-1Ra") (SEQ ID NO:26), hIL-1Ra β (SEQ ID NO:27) and TANGO-
15 77 (SEQ ID NO:28).

Figure 13A is a Western blot depicting the interleukin-18 receptor (IL-18R) binding activity of hIL-1Ra1. In the top panel (depicting a protein band at approximately 22 kD), a conditioned medium containing FLAGhIL-1Ra1 and FLAGIL-18R-ECD-Fc (shown in the left lane) and a conditioned medium containing FLAGhIL-1Ra1 and FLAGIL-18R-ECD-Fc (shown in the right lane) were each immunoprecipitated with protein G-sepharose, and the resulting precipitates were resolved by gel electrophoresis and Western blotting with anti-FLAG monoclonal antibody. In the middle and bottom panels (depicting protein bands at approximately 22 kD and 85 kD), a second aliquot from the FLAGhIL-1Ra1 and FLAGIL-18R-ECD-Fc conditioned medium used in the top panel (shown in the left lane) and a second aliquot from the FLAGhIL-1Ra1 and FLAGIL-18R-ECD-Fc conditioned medium used in the top panel (shown in the right lane) were each immunoprecipitated with anti-FLAG monoclonal antibody, and the resulting precipitates were resolved by gel electrophoresis and Western blotting with anti-FLAG monoclonal antibody.

Figure 13B is a Western blot depicting the IL-1R binding activity of hIL-1Ra3. In the top panel (depicting a protein band at approximately 20 kD), a conditioned medium containing hIL-1Ra3-FLAG and FLAGDR6-Fc (shown in the left lane), a conditioned medium containing hIL-1Ra3-FLAG and FLAGIL-18R-ECD-Fc (shown in the middle lane), and conditioned medium containing hIL-1Ra3-FLAG and FLAGIL-18R-ECD-Fc (shown in the right lane) were each immunoprecipitated with protein G sepharose, and the resulting precipitates were resolved by
35 gel electrophoresis and Western blotting with anti-FLAG monoclonal antibody. In the middle and bottom panels (depicting protein bands at approximately 20 kD and 85 kD), a second aliquot from the hIL-1Ra3-FLAG and FLAGDR6-Fc conditioned medium used in the top panel (shown in the left lane), a second aliquot from the hIL-1Ra3-FLAG and FLAGIL-18R-ECD-Fc conditioned medium used in the top panel (shown in the middle lane) and a second aliquot from the hIL-1Ra3-FLAG and FLAGIL-18R-ECD-Fc conditioned medium used in the top panel (shown in the right lane) were each immunoprecipitated with anti-FLAG monoclonal antibody, and the

resulting precipitates were resolved by gel electrophoresis and Western blotting with anti-FLAG monoclonal antibody.

Figure 14 is a Western blot depicting the interleukin-1 receptor (IL-1R) binding activity of mL-1Ra3. In the top panel (depicting a protein band at approximately 21 kD) and the bottom panel (depicting protein bands at approximately 85 kD) the FLAGIL-1R-ECD-Fc in conditioned medium (shown in the left lane) and the FLAGIL-18R-ECD-Fc in conditioned medium (shown in the right lane) were immobilized with protein G-agarose, the resulting solid phase was contacted with conditioned medium containing FLAGmIL-1Ra3, and the resulting bound complexes were resolved by gel electrophoresis and Western blotting with anti-FLAG monoclonal antibody.

Figure 15 shows the nucleotide sequence (SEQ ID NO:18) and derived amino acid sequence (SEQ ID NO:19) of a native sequence hIL-1Ra1L polypeptide. The start and stop codons in the coding sequence are located at nucleotide positions 4-6 and 625-627, respectively. The putative signal sequence extends from amino acid positions 1 to 34. A putative cAMP- and cGMP-dependent protein kinase phosphorylation site is located at amino acid positions 47-50. Putative N-myristoylation sites are located at amino acid positions 64-69 and 101-106.

Figure 16 shows the nucleotide sequence (SEQ ID NO:20) and derived amino acid sequence (SEQ ID NO:21) of a native sequence hIL-1Ra1S polypeptide. The start and stop codons in the coding sequence are located at nucleotide positions 4-6 and 505-507, respectively. A putative signal sequence extends from amino acid positions 1 to 46. A putative N-myristoylation site is located at amino acid positions 61-66.

Figure 17 shows the single stranded nucleotide sequence (SEQ ID NO:23) of EST AI343258 (lower strand) along with its complementary nucleotide sequence (SEQ ID NO:22) (upper strand).

Figure 18 is an amino acid sequence alignment of native sequence hIL-1Ra1 (SEQ ID NO:3), hIL-1Ra1L (SEQ ID NO:19), hIL-1Ra1V (SEQ ID NO:25) and hIL-1Ra1S (SEQ ID NO:21) polypeptides.

Figure 19 shows the nucleotide sequence (SEQ ID NO:24) and derived amino acid sequence (SEQ ID NO:25) of a native sequence hIL-1Ra1V polypeptide. The start and stop codons in the coding sequence are located at nucleotide positions 73-75 and 727-729, respectively. An alternate start codon is located at nucleotide positions 106-108. A putative signal sequence extends from amino acid positions 1 to 48.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

I. Definitions:

The terms "interleukin-1-like polypeptide", "interleukin-1-like protein", "IL-1lp", "IL-1lp polypeptide", and "IL-1lp protein" encompass any native sequence IL-1lp, and further encompass IL-1lp variants (which are further defined herein). The IL-1lp may be isolated from a variety of sources, such as from human tissue types or from another source, or prepared by recombinant and/or synthetic methods.

A "native sequence IL-1lp" comprises a polypeptide having the same amino acid sequence as a native sequence hIL-1Ra1, hIL-1Ra1L, hIL-1Ra1V, hIL-1Ra1S, hIL-1Ra2, hIL-1Ra3, or mL-1Ra3, (which are further defined herein). Such native sequence IL-1lp can be isolated from

nature or can be produced by recombinant and/or synthetic means. The term "native sequence IL-1lp" specifically encompasses naturally-occurring truncated or secreted forms (e.g., a processed, mature sequence) and naturally-occurring allelic variants of the IL-1lp.

The terms "naturally-occurring amino acid sequence" and "native amino acid sequence" mean any amino acid sequence found in a polypeptide existing in nature, i.e. present in a naturally-occurring polypeptide.

The terms "non-naturally-occurring amino acid sequence" and "non-native amino acid sequence" mean any amino acid sequence not found in a polypeptide existing in nature, i.e. not present in a naturally-occurring polypeptide.

"IL-1lp variant" is defined as any polypeptide that comprises a variant of hIL-1Ra1, hIL-1Ra1L, hIL-1Ra1V, hIL-1Ra1S, hIL-1Ra2, hIL-1Ra3, or mIL-1Ra3 (which are further defined herein).

Human interleukin-1 receptor antagonist analog 1 ("hIL-1Ra1"), hIL-1Ra1 polypeptide, and hIL-1Ra1 protein are defined as any native sequence hIL-1Ra1 or variant hIL-1Ra1.

A "native sequence hIL-1Ra1" means a polypeptide comprising a naturally-occurring amino acid sequence selected from the group consisting of: (1) the amino acid sequence of amino acid residues from at or about 37 to at or about 63 of Figure 2 (SEQ ID NO:5); (2) the amino acid sequence of amino acid residues from at or about 37 to at or about 203 of Figure 2 (SEQ ID NO:5); (3) the amino acid sequence of amino acid residues from at or about 15 to about 53 of Figure 3 (SEQ ID NO:7); (4) the amino acid sequence of amino acid residues from at or about 15 to at or about 193 of Figure 3 (SEQ ID NO:7); and (5) the amino acid sequence of any naturally-occurring truncated or secreted form or any naturally-occurring allelic variant of a polypeptide comprising the amino acid sequence of (1) or (2) or (3) or (4). In one embodiment of the invention, the native sequence hIL-1Ra1 comprises amino acids from at or about 37 to at or about 203 of Figure 2 (SEQ ID NO:5) or amino acids from at or about 15 to at or about 193 of Figure 3 (SEQ ID NO:7).

"hIL-1Ra1 variant" is defined as any hIL-1Ra1 N-terminal variant or hIL-1Ra1 full sequence variant (which are further defined herein).

"hIL-1Ra1 N-terminal variant" means any hIL-1Ra1 other than a native sequence hIL-1Ra1, which variant is an active hIL-1Ra1, as defined below, having at least about 80% amino acid sequence identity with an amino acid sequence selected from the group consisting of: (1) the amino acid sequence of amino acid residues from at or about 37 to at or about 63 of Figure 2 (SEQ ID NO:5); and (2) the amino acid sequence of amino acid residues from at or about 15 to at or about 53 of Figure 3 (SEQ ID NO:7). Such hIL-1Ra1 N-terminal variants include, for instance, hIL-1Ra1 polypeptides wherein one or more amino acid residues are added, or deleted, internally or at the N- or C-terminus, in the sequence of amino acid residues from at or about 37 to at or about 63 of Figure 2 (SEQ ID NO:5) or in the sequence of amino acid residues from at or about 15 to at or about 53 of Figure 3 (SEQ ID NO:7). Ordinarily, an hIL-1Ra1 N-terminal variant will have at least about 80% amino acid sequence identity, or at least about 85% amino acid sequence identity, or at least about 90% amino acid sequence identity, or at least about 95% amino acid sequence identity with an amino acid sequence selected from the group

consisting of: (1) the amino acid sequence of amino acid residues from at or about 37 to at or about 63 of Figure 2 (SEQ ID NO:5); and (2) the amino acid sequence of amino acid residues from at or about 15 to at or about 53 of Figure 3 (SEQ ID NO:7).

"hIL-1Ra1 full sequence variant" means any hIL-1Ra1 other than a native sequence hIL-1Ra1, which variant retains at least one biologic activity of a native sequence hIL-1Ra1, such as the ability to bind IL-18R, and which variant has at least about 80% amino acid sequence identity, or at least about 85% amino acid sequence identity, or at least about 90% amino acid sequence identity, or at least about 95% amino acid sequence identity with an amino acid sequence selected from the group consisting of: (1) the amino acid sequence of amino acid residues from at or about 37 to at or about 203 of Figure 2 (SEQ ID NO:5); and (2) the amino acid sequence of amino acid residues from at or about 15 to at or about 193 of Figure 3 (SEQ ID NO:7). Such hIL-1Ra1 full sequence variants include, for instance, hIL-1Ra1 polypeptides wherein one or more amino acid residues are added, or deleted, internally or at the N- or C-terminus, in the sequence of amino acid residues from at or about 37 to at or about 203 of Figure 2 (SEQ ID NO:5) or in the sequence of amino acid residues from at or about 15 to at or about 193 of Figure 3 (SEQ ID NO:7).

Human interleukin-1 receptor antagonist analog 1 long ("hIL-1Ra1L"), hIL-1Ra1L polypeptide, and hIL-1Ra1L protein are defined as any native sequence hIL-1Ra1L or hIL-1Ra1L variant (which are further defined herein).

A "native sequence hIL-1Ra1L" means a polypeptide comprising a naturally-occurring amino acid sequence selected from the group consisting of: (1) the amino acid sequence of amino acid residues from at or about 26 to at or about 44 of Figure 15 (SEQ ID NO:19); (2) the amino acid sequence of amino acid residues from at or about 26 to at or about 207 of Figure 15 (SEQ ID NO:19); and (3) the amino acid sequence of any naturally-occurring truncated or secreted form or any naturally-occurring allelic variant of a polypeptide comprising the amino acid sequence of (1) or (2). In one embodiment of the invention, the native sequence hIL-1Ra1L comprises amino acids from at or about 26 to at or about 207 of Figure 15 (SEQ ID NO:19).

"hIL-1Ra1L variant" is defined as any hIL-1Ra1L N-terminal variant or hIL-1Ra1L full sequence variant or hIL-1Ra1L fusion variant (which are further defined herein).

"hIL-1Ra1L N-terminal variant" means any hIL-1Ra1L other than a native sequence hIL-1Ra1L, which variant is an active hIL-1Ra1L, as defined below, having at least about 80% amino acid sequence identity with the amino acid sequence of amino acid residues from at or about 26 to at or about 44 of Figure 15 (SEQ ID NO:19). Such hIL-1Ra1L N-terminal variants include, for instance, hIL-1Ra1L polypeptides wherein one or more amino acid residues are added, or deleted, internally or at the N- or C-terminus, in the sequence of amino acid residues from at or about 26 to at or about 44 of Figure 15 (SEQ ID NO:19). Ordinarily, an hIL-1Ra1L N-terminal variant will have at least about 80% amino acid sequence identity, or at least about 85% amino acid sequence identity, or at least about 90% amino acid sequence identity, or at least about 95% amino acid sequence identity with the amino acid sequence of amino acid residues from at or about 26 to at or about 44 of Figure 15 (SEQ ID NO:19).

"hIL-1Ra1L full sequence variant" means any hIL-1Ra1L other than a native sequence hIL-1Ra1L, which variant retains at least one biologic activity of a native sequence hIL-1Ra1L, such as the ability to bind IL-18R, and which variant has at least about 80% amino acid sequence identity, or at least about 85% amino acid sequence identity, or at least about 90% amino acid sequence identity, or at least about 95% amino acid sequence identity with the amino acid sequence of amino acid residues from at or about 26 to at or about 207 of Figure 15 (SEQ ID NO:19). Such hIL-1Ra1L full sequence variants include, for instance, hIL-1Ra1L polypeptides wherein one or more amino acid residues are added, or deleted, internally or at the N- or C-terminus, in the sequence of amino acid residues from at or about 26 to at or about 207 of Figure 15 (SEQ ID NO:19).

"hIL-1Ra1L fusion variant" means a chimeric hIL-1Ra1L consisting of a native sequence hIL-1Ra1L fused at its N- or C-terminus to a heterologous amino acid or amino acid sequence. In one embodiment, the hIL-1Ra1L fusion variant polypeptide consists of a native sequence of hIL-1Ra1L fused at its N-terminus or C-terminus to a heterologous amino acid or amino acid sequence, wherein the heterologous amino acid or amino acid sequence is heterologous to the native sequence, i.e. the resulting chimeric sequence is non-naturally occurring. In another embodiment, the hIL-1Ra1L fusion variant consists of the amino acid sequence of amino acids from at or about 26 to at or about 207, inclusive of Figure 15 (SEQ ID NO:19), or the amino acid sequence of amino acid residues from at or about 1 to at or about 207, inclusive of Figure 15 (SEQ ID NO:19), fused at its N-terminus or C-terminus to a heterologous amino acid or amino acid sequence to form a non-naturally occurring fusion protein. Such hIL-1Ra1L fusion variants include, for instance, hIL-1Ra1L polypeptides wherein a heterologous secretion leader sequence is fused to the N-terminus of the sequence of amino acid residues from at or about 26 to at or about 207 of Figure 15 (SEQ ID NO:19), or amino acid residues from at or about 1 to at or about 207 of Figure 15 (SEQ ID NO:19).

Human interleukin-1 receptor antagonist analog 1 long allelic variant ("hIL-1Ra1V"), hIL-1Ra1V polypeptide, and hIL-1Ra1V protein are defined as any native sequence hIL-1Ra1V or hIL-1Ra1V variant (which are further defined herein).

A "native sequence hIL-1Ra1V" means a polypeptide comprising a naturally-occurring amino acid sequence selected from the group consisting of: (1) the amino acid sequence of amino acid residues from at or about 46 to at or about 55 of Figure 19 (SEQ ID NO:25); (2) the amino acid sequence of amino acid residues from at or about 46 to at or about 218 of Figure 19 (SEQ ID NO:25); (3) the amino acid sequence of amino acid residues from at or about 37 to at or about 218 of Figure 19 (SEQ ID NO:25); (4) the amino acid sequence of amino acid residues from at or about 12 to at or about 218 of Figure 19 (SEQ ID NO:25); and (5) the amino acid sequence of any naturally-occurring truncated or secreted form or any naturally-occurring allelic variant of a polypeptide comprising the amino acid sequence of (1) or (2) or (3) or (4). In one embodiment of the invention, the native sequence hIL-1Ra1V comprises amino acids from at or about 46 to at or about 218 of Figure 19 (SEQ ID NO:25), or amino acids from at or about 37 to at or about 218 of Figure 19 (SEQ ID NO:25), or amino acids from at or about 12 to at or about 218 of Figure 19 (SEQ ID NO:25), or amino acids from at or about 12 to at or about 218 of Figure 19 (SEQ ID NO:25), or amino acids from at or about 12 to at or about 218 of Figure 19 (SEQ ID NO:25).

(SEQ ID NO:25), or amino acids from at or about 1 to at or about 218 of Figure 19 (SEQ ID NO:25).

"hIL-1Ra1V variant" is defined as any hIL-1Ra1V N-terminal variant or hIL-1Ra1V full sequence variant or hIL-1Ra1V fusion variant (which are further defined herein).

"hIL-1Ra1V N-terminal variant" is defined as any hIL-1Ra1V other than a native sequence hIL-1Ra1V, which variant is an active hIL-1Ra1V, as defined below, having at least about 80% amino acid sequence identity with the amino acid sequence of amino acid residues from at or about 46 to at or about 89 of Figure 19 (SEQ ID NO:25). Such hIL-1Ra1V N-terminal variants include, for instance, hIL-1Ra1V polypeptides wherein one or more amino acid residues are added, internally or at the N- or C-terminus, in the sequence of amino acid residues from at or about 46 to at or about 89 of Figure 19 (SEQ ID NO:25). Ordinarily, an hIL-1Ra1V N-terminal variant will have at least about 80% amino acid sequence identity, or at least about 85% amino acid sequence identity, or at least about 90% amino acid sequence identity, or at least about 95% amino acid sequence identity with the sequence of amino acid residues from at or about 46 to at or about 89 of Figure 19 (SEQ ID NO:25).

"hIL-1Ra1V full sequence variant" means any hIL-1Ra1V other than a native sequence hIL-1Ra1V, which variant retains at least one biologic activity of a native sequence hIL-1Ra1V, such as the ability to bind IL-18R, and which variant has at least about 80% amino acid sequence identity, or at least about 85% amino acid sequence identity, or at least about 90% amino acid sequence identity, or at least about 95% amino acid sequence identity with the sequence of amino acid residues from at or about 46 to at or about 218 of Figure 19 (SEQ ID NO:25).

"hIL-1Ra1V fusion variant" means a chimeric hIL-1Ra1V consisting of a native sequence hIL-1Ra1V fused at its N- or C-terminus to a heterologous amino acid or amino acid sequence.

In one embodiment, the hIL-1Ra1V fusion variant polypeptide consists of a native sequence of hIL-1Ra1V fused at its N-terminus or C-terminus to a heterologous amino acid or amino acid sequence, wherein the heterologous amino acid or amino acid sequence is heterologous to the native sequence, i.e. the resulting chimeric sequence is non-naturally occurring. In another embodiment, the hIL-1Ra1V fusion variant consists of the amino acid sequence of amino acids from at or about 46 to at or about 218 of Figure 19 (SEQ ID NO:25), or the amino acid sequence of amino acids from at or about 37 to at or about 218 of Figure 19 (SEQ ID NO:25), or the amino acid sequence of amino acids from at or about 12 to at or about 218 of Figure 19 (SEQ ID NO:25), or the amino acid sequence of amino acids from at or about 1 to at or about 218 of Figure 19 (SEQ ID NO:25), fused at its N-terminus or C-terminus to a heterologous amino acid sequence to form a non-naturally occurring fusion protein. Such hIL-1Ra1V fusion variants include, for instance, hIL-1Ra1V polypeptides wherein a heterologous secretion leader sequence is fused to the N-terminus of the sequence of amino acid residues from at or about 46 to at or about 218 of Figure 19 (SEQ ID NO:25), or amino acid residues from at or about 37 to at or about 218 of Figure 19 (SEQ ID NO:25), or amino acid residues from at or about 12 to at or about 218 of Figure 19 (SEQ ID NO:25), or amino acid residues from at or about 1 to at or about 218 of Figure 19 (SEQ ID NO:25).

Human interleukin-1 receptor antagonist analog 1 short ("hIL-1Ra1S"), hIL-1Ra1S polypeptide, and hIL-1Ra1S protein are defined as any native sequence hIL-1Ra1S or hIL-1Ra1S variant (which are further defined herein).

A "native sequence hIL-1Ra1S" means a polypeptide comprising a naturally-occurring amino acid sequence selected from the group consisting of: (1) the amino acid sequence of amino acid residues from at or about 1 to at or about 38 of Figure 16 (SEQ ID NO:21); (2) the amino acid sequence of amino acid residues from at or about 26 to at or about 167 of Figure 16 (SEQ ID NO:21); (3) the amino acid sequence of amino acid residues from at or about 39 to at or about 167 of Figure 16 (SEQ ID NO:21); (4) the amino acid sequence of amino acid residues from at or about 47 to at or about 167 of Figure 16 (SEQ ID NO:21); and (5) the amino acid sequence of any naturally-occurring truncated or secreted form or any naturally-occurring allelic variant of a polypeptide comprising the amino acid sequence of (1) or (2) or (3) or (4). In one embodiment of the invention, the native sequence hIL-1Ra1S comprises amino acids from at or about 26 to at or about 167 of Figure 16 (SEQ ID NO:21), or amino acids from at or about 1 to at or about 167 of Figure 16 (SEQ ID NO:21). In another embodiment, the native sequence hIL-1Ra1S consists of amino acids from at or about 47 to at or about 167 of Figure 16 (SEQ ID NO:21) or amino acids from at or about 39 to at or about 167 of Figure 16 (SEQ ID NO:21).

"hIL-1Ra1S fusion variant" and "hIL-1Ra1S variant" mean a chimeric hIL-1Ra1S consisting of a native sequence hIL-1Ra1S fused at its N-terminus or C-terminus to a heterologous amino acid or amino acid sequence. In one embodiment, the hIL-1Ra1S fusion variant polypeptide consists of a native sequence of hIL-1Ra1S fused at its N-terminus or C-terminus to a heterologous amino acid or amino acid sequence, wherein the heterologous amino acid or amino acid sequence is heterologous to the native sequence, i.e. the resulting chimeric sequence is non-naturally occurring. In another embodiment, the hIL-1Ra1S fusion variant consists of the amino acid sequence of amino acids from at or about 47 to at or about 167 of Figure 16 (SEQ ID NO:21), or the amino acid sequence of amino acids from at or about 39 to at or about 167 of Figure 16 (SEQ ID NO:21), fused at its N-terminus or C-terminus to a heterologous amino acid or amino acid sequence to form a non-naturally occurring fusion protein. Such hIL-1Ra1S fusion variants include, for instance, hIL-1Ra1S polypeptides wherein a heterologous secretion leader sequence is fused to the N-terminus of the sequence of amino acid residues from at or about 47 to at or about 167 of Figure 16 (SEQ ID NO:21), or amino acid residues from at or about 39 to at or about 167 of Figure 16 (SEQ ID NO:21).

Human interleukin-1 receptor antagonist analog 2 ("hIL-1Ra2"), hIL-1Ra2 polypeptide, and hIL-1Ra2 protein are defined as any native sequence hIL-1Ra2 or hIL-1Ra2 fusion variant (which are further defined herein).

A "native sequence hIL-1Ra2" means (1) a polypeptide comprising the amino acid sequence of amino acid residues from at or about 1 to at or about 134 of Figure 5 (SEQ ID NO:10) or (2) a polypeptide consisting of a naturally-occurring truncated or secreted form of the polypeptide of (1). In one embodiment of the invention, the native sequence hIL-1Ra2 consists of amino acids from at or about 27 to at or about 134 of Figure 5 (SEQ ID NO:10), or amino acids from at or about 1 to at or about 134 of Figure 5 (SEQ ID NO:10).

“hIL-1Ra2 fusion variant” and “hIL-1Ra2 variant” mean a chimeric hIL-1Ra2 consisting of a native sequence hIL-1Ra2 fused at its N-terminus or C-terminus to a heterologous amino acid or amino acid sequence. In one embodiment, the hIL-1Ra2 fusion variant polypeptide consists of a native sequence of hIL-1Ra2 fused at its N-terminus or C-terminus to a heterologous amino acid or amino acid sequence, wherein the heterologous amino acid or amino acid sequence is heterologous to the native sequence, i.e. the resulting chimeric sequence is non-naturally occurring. In another embodiment, the hIL-1Ra2 variant consists of the amino acid sequence of amino acids from at or about 27 to at or about 134 of Figure 5 (SEQ ID NO:10), or amino acids from at or about 1 to at or about 134 of Figure 5 (SEQ ID NO:10), fused at its N-terminus or C-terminus to a heterologous amino acid or amino acid sequence to form a non-naturally occurring fusion protein. Such hIL-1Ra2 fusion variants include, for instance, hIL-1Ra2 polypeptides wherein a heterologous secretion leader sequence is fused to the N-terminus of the sequence of amino acids from at or about 27 to at or about 134 of Figure 5 (SEQ ID NO:10), or amino acids from at or about 1 to at or about 134 of Figure 5 (SEQ ID NO:10).

Human interleukin-1 receptor antagonist analog 3 (“hIL-1Ra3”), hIL-1Ra3 polypeptide, and hIL-1Ra3 protein are defined as any native sequence hIL-1Ra3 or variant hIL-1Ra3 (which are further defined herein).

A “native sequence hIL-1Ra3” means a polypeptide comprising an amino acid sequence selected from the group consisting of: (1) the amino acid sequence of amino acid residues from at or about 95 to at or about 134 of Figure 7 (SEQ ID NO:13); (2) the amino acid sequence of amino acid residues from at or about 34 to at or about 155 of Figure 7 (SEQ ID NO:13); and (3) the amino acid sequence of any naturally-occurring truncated or secreted form or any naturally-occurring allelic variant of a polypeptide comprising the amino acid sequence of (1) or (2). In one embodiment of the invention, the native sequence hIL-1Ra3 comprises amino acids from at or about 34 to at or about 155 of Figure 7 (SEQ ID NO:13), or amino acids from at or about 2 to at or about 155 of Figure 7 (SEQ ID NO:13).

“hIL-1Ra3 variant” is defined as any hIL-1Ra3 C-terminal variant or hIL-1Ra3 full sequence variant (which are further defined herein).

“hIL-1Ra3 C-terminal variant” means any hIL-1Ra3 other than a native sequence hIL-1Ra3, which variant is an active hIL-1Ra3, as defined below, having at least about 80% amino acid sequence identity with the amino acid sequence of amino acid residues from at or about 95 to at or about 134 of Figure 7 (SEQ ID NO:13) or the amino acid sequence of amino acid residues from at or about 80 to at or about 155 of Figure 7 (SEQ ID NO:13). Such hIL-1Ra3 C-terminal variants include, for instance, hIL-1Ra3 polypeptides wherein one or more amino acid residues are added, or deleted, internally or at the N- or C-terminus, in the sequence of amino acid residues from at or about 95 to at or about 134 of Figure 7 (SEQ ID NO:13) or in the sequence of amino acid residues from at or about 80 to at or about 155 of Figure 7 (SEQ ID NO:13). Ordinarily, an hIL-1Ra3 C-terminal variant will have at least about 80% amino acid sequence identity, or at least about 85% amino acid sequence identity, or at least about 90% amino acid sequence identity, or at least about 95% amino acid sequence identity with the amino acid sequence of amino acid residues from at or about 95 to at or about 134 of Figure 7 (SEQ ID

NO:13) or the amino acid sequence of amino acid residues from at or about 80 to at or about 155 of Figure 7 (SEQ ID NO:13).

"hIL-1Ra3 full sequence variant" means any hIL-1Ra3 other than a native sequence hIL-1Ra3, which variant retains at least one biologic activity of a native sequence hIL-1Ra3, such as the ability to bind IL-1R, and which variant has at least about 80% amino acid sequence identity, or at least about 85% amino acid sequence identity, or at least about 90% amino acid sequence identity, or at least about 95% amino acid sequence identity with the amino acid sequence of amino acid residues from at or about 34 to at or about 155 of Figure 7 (SEQ ID NO:13) or the amino acid sequence of amino acid residues from at or about 2 to at or about 155 of Figure 7 (SEQ ID NO:13). Such hIL-1Ra3 full sequence variants include, for instance, hIL-1Ra3 polypeptides wherein one or more amino acid residues are added, or deleted, internally or at the N- or C-terminus, in the sequence of amino acid residues from at or about 34 to at or about 155 of Figure 7 (SEQ ID NO:13) or the amino acid sequence of amino acid residues from at or about 2 to at or about 155 of Figure 7 (SEQ ID NO:13).

Murine interleukin-1 receptor antagonist analog 3 ("mIL-1Ra3"), mIL-1Ra3 polypeptide, and mIL-1Ra3 protein are defined as any native sequence mIL-1Ra3 or variant mIL-1Ra3.

A "native sequence mIL-1Ra3" means a polypeptide comprising an amino acid sequence selected from the group consisting of: (1) the amino acid sequence of amino acid residues from at or about 95 to at or about 134 of Figure 9 (SEQ ID NO:16); (2) the amino acid sequence of amino acid residues from at or about 34 to at or about 155 of Figure 9 (SEQ ID NO:16); and (3) the amino acid sequence of any naturally-occurring truncated or secreted form or naturally-occurring allelic variant of a polypeptide comprising the amino acid sequence of (1) or (2). In one embodiment of the invention, the native sequence mIL-1Ra3 comprises amino acids from at or about 34 to at or about 155 of Figure 9 (SEQ ID NO:16).

"mIL-1Ra3 variant" is defined as any mIL-1Ra3 C-terminal variant or mIL-1Ra3 full sequence variant (which are further defined herein).

"mIL-1Ra3 C-terminal variant" means any mIL-1Ra3 other than a native sequence mIL-1Ra3, which variant is an active mIL-1Ra3, as defined below, having at least about 80% amino acid sequence identity with the amino acid sequence of amino acids from at or about 95 to at or about 134 of Figure 9 (SEQ ID NO:16). Such mIL-1Ra3 C-terminal variants include, for instance, mIL-1Ra3 polypeptides wherein one or more amino acid residues are added, or deleted, internally or at the N- or C-terminus, in the sequence of amino acids from at or about 95 to at or about 134 of Figure 9 (SEQ ID NO:16). Ordinarily, an mIL-1Ra3 C-terminal variant will have at least about 80% amino acid sequence identity, or at least about 85% amino acid sequence identity, or at least about 90% amino acid sequence identity, and or at least about 95% amino acid sequence identity with the amino acid sequence of amino acids 95 to 134 of Figure 9 (SEQ ID NO:16).

"mIL-1Ra3 full sequence variant" means any mIL-1Ra3 other than a native sequence mIL-1Ra3, which variant retains at least one biologic activity of a native sequence mIL-1Ra3, such as the ability to bind IL-1R, and which variant has at least about 85% amino acid sequence identity, or at least about 90% amino acid sequence identity, or at least about 95% sequence

identity with the amino acid sequence of amino acid residues from at or about 34 to at or about 155 of Figure 9 (SEQ ID NO:16) or the amino acid sequence of amino acid residues from at or about 2 to at or about 155 of Figure 9 (SEQ ID NO:16). Such mL-1Ra3 full sequence variants include, for instance, mL-1Ra3 polypeptides wherein one or more amino acid residues are added, or deleted, internally or at the N- or C-terminus, in the sequence of amino acid residues from at or about 34 to at or about 155 of Figure 9 (SEQ ID NO:16) or in the sequence of amino acid residues from at or about 2 to at or about 155 of Figure 9 (SEQ ID NO:16).

"Human interleukin-1-like polypeptide", "hIL-1lp", "hIL-1lp polypeptide", "hIL-1lp protein", "human interleukin-1 receptor antagonist analog", "hIL-1Raa", "hIL-1Raa polypeptide", and "hIL-1Raa protein" are defined as any hIL-1Ra1, hIL-1Ra2 or hIL-1Ra3 polypeptide.

"Native sequence hIL-1lp" and "native sequence hIL-1Raa" are defined as any polypeptide that comprises a native sequence hIL-1Ra1, hIL-1Ra2, or hIL-1Ra3.

"hIL-1lp variant" is defined as any polypeptide that comprises a variant of hIL-1Ra1, hIL-1Ra2, or hIL-1Ra3.

"Interleukin-1 receptor", "interleukin-1 receptor polypeptide", "interleukin-1 receptor protein", "IL-1 receptor", "IL-1R", "IL-1R polypeptide", and "IL-1R protein", are defined as the family of cell surface proteins that bind to interleukin-1 (IL-1) and/or function in IL-1-induced signal transduction in a given species, such as human or mouse. IL-1R includes the human T cell-expressed IL-1 receptor disclosed in Sims, et al., *Proc. Natl. Acad. Sci. (USA)*, **86**: 8946-8950 (1989).

"Interleukin-18 receptor", "interleukin-18 receptor polypeptide", "interleukin-18 receptor protein", "IL-18 receptor", "IL-18R", "IL-18R polypeptide", and "IL-18R protein", are defined as the family of cell surface proteins that bind to interleukin-18 (IL-18) and/or function in IL-18-induced signal transduction in a given species, such as human or mouse. IL-18R includes the IL-1 receptor related protein (IL-1Rrp) described in Torigoe et al., *J. Biol. Chem.*, **272**: 25737-25742 (1997) and the IL-18 receptor accessory protein-like molecule (IL-18RacPL) described in Born et al., *J. Biol. Chem.*, **273**: 29445-29450 (1998).

"Interleukin-1-like family" and "IL-1-like family" are used to indicate the family of polypeptides related to the ligands of IL-1R or IL-18R. The IL-1-like family includes IL-1 receptor agonists and antagonists and related polypeptides such as IL-1 α (described in Bazan et al., *Nature*, **379**: 591 (1996), IL-1 β (Bazan et al.), IL-18 (interferon- γ inducing factor)(IGIF)(Bazan et al.), IL-1 receptor antagonist polypeptides such as secretory IL-1Ra (sIL-1Ra)(described in Eisenberg et al., *Nature*, **343**: 341-346 (1990)) and intracellular IL-1Ra (icIL-1Ra) (described in Haskill et al., *Proc. Natl. Acad. Sci. (USA)*, **88**: 3681-3685 (1991)), and the IL-1lp polypeptides of the invention.

"Percent (%) amino acid sequence identity" with respect to the IL-1lp sequences identified herein is defined as the percentage of amino acid residues in a candidate sequence that are identical with the amino acid residues in an IL-1lp sequence, after aligning the sequences and introducing gaps, if necessary, to achieve the maximum percent sequence identity, and not considering any conservative substitutions as part of the sequence identity. Alignment for purposes of determining percent amino acid sequence identity can be achieved in various ways

that are within the skill in the art, for instance, using publicly available computer software such as BLAST, BLAST-2, ALIGN, ALIGN-2 or Megalign (DNASTAR) software. Those skilled in the art can determine appropriate parameters for measuring alignment, including any algorithms needed to achieve maximal alignment over the full-length of the sequences being compared. For purposes herein, however, % amino acid sequence identity values are obtained as described below by using the sequence comparison computer program ALIGN-2, wherein the complete source code for the ALIGN-2 program is provided in Tables 3A-3Q. The ALIGN-2 sequence comparison computer program was authored by Genentech, Inc. and the source code shown in Tables 3A-3Q has been filed with user documentation in the U.S. Copyright Office, Washington D.C., 20559, where it is registered under U.S. Copyright Registration No. TXU510087. The ALIGN-2 program is publicly available through Genentech, Inc., South San Francisco, California or may be compiled from the source code provided in Tables 3A-3Q. The ALIGN-2 program should be compiled for use on a UNIX operating system, preferably digital UNIX V4.0D. All sequence comparison parameters are set by the ALIGN-2 program and do not vary.

For purposes herein, the % amino acid sequence identity of a given amino acid sequence A to, with, or against a given amino acid sequence B (which also can be phrased as a given amino acid sequence A that has or comprises a certain % amino acid sequence identity to, with, or against a given amino acid sequence B) is calculated as follows:

$$100 \text{ times the fraction } X/Y$$

where X is the number of amino acid residues scored as identical matches by the sequence alignment program ALIGN-2 in that program's alignment of A and B, and where Y is the total number of amino acid residues in B. It will be appreciated that where the length of amino acid sequence A is not equal to the length of amino acid sequence B, the % amino acid sequence identity of A to B will not equal the % amino acid sequence identity of B to A. As examples of % amino acid sequence identity calculations, Tables 2A-2B demonstrate how to calculate the % amino acid sequence identity of the amino acid sequence designated "Comparison Protein" to the amino acid sequence designated "PRO".

Unless specifically stated otherwise, all % amino acid sequence identity values used herein are obtained as described above using the ALIGN-2 sequence comparison computer program. However, % amino acid sequence identity may also be determined using the sequence comparison program NCBI-BLAST2 (Altschul et al., *Nucleic Acids Res.* 25:3389-3402 (1997)). The NCBI-BLAST2 sequence comparison program may be downloaded from <http://www.ncbi.nlm.nih.gov>. NCBI-BLAST2 uses several search parameters, wherein all of those search parameters are set to default values including, for example, unmask = yes, strand = all, expected occurrences = 10, minimum low complexity length = 15/5, multi-pass e-value = 0.01, constant for multi-pass = 25, dropoff for final gapped alignment = 25 and scoring matrix = BLOSUM62.

In situations where NCBI-BLAST2 is employed for amino acid sequence comparisons, the % amino acid sequence identity of a given amino acid sequence A to, with, or against a given

amino acid sequence B (which also can be phrased as a given amino acid sequence A that has or comprises a certain % amino acid sequence identity to, with, or against a given amino acid sequence B) is calculated as follows:

5 100 times the fraction X/Y

where X is the number of amino acid residues scored as identical matches by the sequence alignment program NCBI-BLAST2 in that program's alignment of A and B, and where Y is the total number of amino acid residues in B. It will be appreciated that where the length of amino acid sequence A is not equal to the length of amino acid sequence B, the % amino acid sequence identity of A to B will not equal the % amino acid sequence identity of B to A.

10 "Percent (%) nucleic acid sequence identity" with respect to the IL-11p polypeptide-encoding nucleic acid sequences identified herein is defined as the percentage of nucleotides in a candidate sequence that are identical with the nucleotides in an IL-11p polypeptide-encoding nucleic acid sequence, after aligning the sequences and introducing gaps, if necessary, to achieve the maximum percent sequence identity. Alignment for purposes of determining percent nucleic acid sequence identity can be achieved in various ways that are within the skill in the art, for instance, using publicly available computer software such as BLAST, BLAST-2, ALIGN, ALIGN-2 or Megalign (DNASTAR) software. Those skilled in the art can determine appropriate parameters for measuring alignment, including any algorithms needed to achieve maximal alignment over the full-length of the sequences being compared. For purposes herein, however, % nucleic acid sequence identity values are obtained as described below by using the sequence comparison computer program ALIGN-2, wherein the complete source code for the ALIGN-2 program is provided in Tables 3A-3Q. The ALIGN-2 sequence comparison computer program was authored by Genentech, Inc. and the source code shown in Tables 3A-3Q has been filed with user documentation in the U.S. Copyright Office, Washington D.C., 20559, where it is registered under U.S. Copyright Registration No. TXU510087. The ALIGN-2 program is publicly available through Genentech, Inc., South San Francisco, California or may be compiled from the source code provided in Tables 3A-3Q. The ALIGN-2 program should be compiled for use on a UNIX operating system, preferably digital UNIX V4.0D. All sequence comparison parameters are set by the ALIGN-2 program and do not vary.

25 For purposes herein, the % nucleic acid sequence identity of a given nucleic acid sequence C to, with, or against a given nucleic acid sequence D (which also can be phrased as a given nucleic acid sequence C that has or comprises a certain % nucleic acid sequence identity to, with, or against a given nucleic acid sequence D) is calculated as follows:

35 100 times the fraction W/Z

where W is the number of nucleotides scored as identical matches by the sequence alignment program ALIGN-2 in that program's alignment of C and D, and where Z is the total number of nucleotides in D. It will be appreciated that where the length of nucleic acid sequence C is not

equal to the length of nucleic acid sequence D, the % nucleic acid sequence identity of C to D will not equal the % nucleic acid sequence identity of D to C. As examples of % nucleic acid sequence identity calculations, Tables 2C-2D demonstrate how to calculate the % nucleic acid sequence identity of the nucleic acid sequence designated "Comparison DNA" to the nucleic acid sequence designated "PRO-DNA".

Unless specifically stated otherwise, all % nucleic acid sequence identity values used herein are obtained as described above using the ALIGN-2 sequence comparison computer program. However, % nucleic acid sequence identity may also be determined using the sequence comparison program NCBI-BLAST2 (Altschul et al., Nucleic Acids Res. 25:3389-3402 (1997)).

The NCBI-BLAST2 sequence comparison program may be downloaded from <http://www.ncbi.nlm.nih.gov>. NCBI-BLAST2 uses several search parameters, wherein all of those search parameters are set to default values including, for example, unmask = yes, strand = all, expected occurrences = 10, minimum low complexity length = 15/5, multi-pass e-value = 0.01, constant for multi-pass = 25, dropoff for final gapped alignment = 25 and scoring matrix = BLOSUM62.

In situations where NCBI-BLAST2 is employed for sequence comparisons, the % nucleic acid sequence identity of a given nucleic acid sequence C to, with, or against a given nucleic acid sequence D (which also can be phrased as a given nucleic acid sequence C that has or comprises a certain % nucleic acid sequence identity to, with, or against a given nucleic acid sequence D) is calculated as follows:

$$100 \text{ times the fraction } W/Z$$

where W is the number of nucleotides scored as identical matches by the sequence alignment program NCBI-BLAST2 in that program's alignment of C and D, and where Z is the total number of nucleotides in D. It will be appreciated that where the length of nucleic acid sequence C is not equal to the length of nucleic acid sequence D, the % nucleic acid sequence identity of C to D will not equal the % nucleic acid sequence identity of D to C.

The term "positives", in the context of the amino acid sequence identity comparisons performed as described above, includes amino acid residues in the sequences compared that are not only identical, but also those that have similar properties. Amino acid residues that score a positive value to an amino acid residue of interest are those that are either identical to the amino acid residue of interest or are a preferred substitution (as defined in Table 1 below) of the amino acid residue of interest.

For purposes herein, the % value of positives of a given amino acid sequence A to, with, or against a given amino acid sequence B (which also can be phrased as a given amino acid sequence A that has or comprises a certain % positives to, with, or against a given amino acid sequence B) is calculated as follows:

$$100 \text{ times the fraction } X/Y$$

where X is the number of amino acid residues scoring a positive value as defined above by the sequence alignment program ALIGN-2 in that program's alignment of A and B, and where Y is the total number of amino acid residues in B. It will be appreciated that where the length of amino acid sequence A is not equal to the length of amino acid sequence B, the % positives of A to B will not equal the % positives of B to A.

"Isolated," when used to describe the various polypeptides disclosed herein, means polypeptide that has been identified and separated and/or recovered from a component of its natural environment. Contaminant components of its natural environment are materials that would typically interfere with diagnostic or therapeutic uses for the polypeptide, and may include enzymes, hormones, and other proteinaceous or non-proteinaceous solutes. In preferred embodiments, the polypeptide will be purified (1) to a degree sufficient to obtain at least 15 residues of N-terminal or internal amino acid sequence by use of a spinning cup sequenator, or (2) to homogeneity by SDS-PAGE under non-reducing or reducing conditions using Coomassie blue or, preferably, silver stain. Isolated polypeptide includes polypeptide *in situ* within recombinant cells, since at least one component of the IL-11p natural environment will not be present. Ordinarily, however, isolated polypeptide will be prepared by at least one purification step.

An "isolated" nucleic acid molecule encoding a IL-11p polypeptide is a nucleic acid molecule that is identified and separated from at least one contaminant nucleic acid molecule with which it is ordinarily associated in the natural source of the IL-11p-encoding nucleic acid.

An isolated IL-11p-encoding nucleic acid molecule is other than in the form or setting in which it is found in nature. Isolated nucleic acid molecules therefore are distinguished from the IL-11p-encoding nucleic acid molecule as it exists in natural cells. However, an isolated nucleic acid molecule encoding a IL-11p polypeptide includes IL-11p-encoding nucleic acid molecules contained in cells that ordinarily express IL-11p where, for example, the nucleic acid molecule is in a chromosomal location different from that of natural cells.

The term "control sequences" refers to DNA sequences necessary for the expression of an operably linked coding sequence in a particular host organism. The control sequences that are suitable for prokaryotes, for example, include a promoter, optionally an operator sequence, and a ribosome binding site. Eukaryotic cells are known to utilize promoters, polyadenylation signals, and enhancers.

Nucleic acid is "operably linked" when it is placed into a functional relationship with another nucleic acid sequence. For example, DNA for a presequence or secretory leader is operably linked to DNA for a polypeptide if it is expressed as a preprotein that participates in the secretion of the polypeptide; a promoter or enhancer is operably linked to a coding sequence if it affects the transcription of the sequence; or a ribosome binding site is operably linked to a coding sequence if it is positioned so as to facilitate translation. Generally, "operably linked" means that the DNA sequences being linked are contiguous, and, in the case of a secretory leader, contiguous and in reading phase. However, enhancers do not have to be contiguous. Linking is accomplished by ligation at convenient restriction sites. If such sites do not exist,

the synthetic oligonucleotide adaptors or linkers are used in accordance with conventional practice.

The term "antibody" is used in the broadest sense and specifically covers single anti-IL-11p monoclonal antibodies (including agonist, antagonist, and neutralizing antibodies) and anti-IL-11p antibody compositions with polyepitopic specificity. The term "monoclonal antibody" as used herein refers to an antibody obtained from a population of substantially homogeneous antibodies, i.e., the individual antibodies comprising the population are identical except for possible naturally-occurring mutations that may be present in minor amounts.

"Stringency" of hybridization reactions is readily determinable by one of ordinary skill in the art, and generally is an empirical calculation dependent upon probe length, washing temperature, and salt concentration. In general, longer probes require higher temperatures for proper annealing, while shorter probes need lower temperatures. Hybridization generally depends on the ability of denatured DNA to reanneal when complementary strands are present in an environment below their melting temperature. The higher the degree of desired homology between the probe and hybridizable sequence, the higher the relative temperature which can be used. As a result, it follows that higher relative temperatures would tend to make the reaction conditions more stringent, while lower temperatures less so. For additional details and explanation of stringency of hybridization reactions, see Ausubel *et al.*, Current Protocols in Molecular Biology, Wiley Interscience Publishers, (1995).

"Stringent conditions" or "high stringency conditions", as defined herein, may be identified by those that: (1) employ low ionic strength and high temperature for washing, for example 0.015 M sodium chloride/0.0015 M sodium citrate/0.1% sodium dodecyl sulfate at 50°C; (2) employ during hybridization a denaturing agent, such as formamide, for example, 50% (v/v) formamide with 0.1% bovine serum albumin/0.1% Ficoll/0.1% polyvinylpyrrolidone/50mM sodium phosphate buffer at pH 6.5 with 750 mM sodium chloride, 75 mM sodium citrate at 42°C; or (3) employ 50% formamide, 5 x SSC (0.75 M NaCl, 0.075 M sodium citrate), 50 mM sodium phosphate (pH 6.8), 0.1% sodium pyrophosphate, 5 x Denhardt's solution, sonicated salmon sperm DNA (50 µg/ml), 0.1% SDS, and 10% dextran sulfate at 42°C, with washes at 42°C in 0.2 x SSC (sodium chloride/sodium citrate) and 50% formamide at 55°C, followed by a high-stringency wash consisting of 0.1 x SSC containing EDTA at 55°C.

"Moderately stringent conditions" may be identified as described by Sambrook *et al.*, Molecular Cloning: A Laboratory Manual, New York: Cold Spring Harbor Press, 1989, and include the use of washing solution and hybridization conditions (e.g., temperature, ionic strength and %SDS) less stringent than those described above. An example of moderately stringent conditions is overnight incubation at 37°C in a solution comprising: 20% formamide, 5 x SSC (150 mM NaCl, 15 mM trisodium citrate), 50 mM sodium phosphate (pH 7.6), 5 x Denhardt's solution, 10% dextran sulfate, and 20 mg/mL denatured sheared salmon sperm DNA, followed by washing the filters in 1 x SSC at about 37-50°C. The skilled artisan will recognize how to adjust the temperature, ionic strength, etc. as necessary to accommodate factors such as probe length and the like.

The term "epitope tagged" when used herein refers to a chimeric polypeptide comprising an IL-11p polypeptide fused to a "tag polypeptide". The tag polypeptide has enough residues to provide an epitope against which an antibody can be made, yet is short enough such that it does not interfere with activity of the polypeptide to which it is fused. The tag polypeptide preferably also is fairly unique so that the antibody does not substantially cross-react with other epitopes. Suitable tag polypeptides generally have at least six amino acid residues and usually between about 8 and 50 amino acid residues (preferably, between about 10 and 20 amino acid residues).

As used herein, the term "immunoadhesin" designates antibody-like molecules which combine the binding specificity of a heterologous protein (an "adhesin") with the effector functions of immunoglobulin constant domains. Structurally, the immunoadhesins comprise a fusion of an amino acid sequence with the desired binding specificity which is other than the antigen recognition and binding site of an antibody (i.e., is "heterologous"), and an immunoglobulin constant domain sequence. The adhesin part of an immunoadhesin molecule typically is a contiguous amino acid sequence comprising at least the binding site of a receptor or a ligand. The immunoglobulin constant domain sequence in the immunoadhesin may be obtained from any immunoglobulin, such as IgG-1, IgG-2, IgG-3, or IgG-4 subtypes, IgA (including IgA-1 and IgA-2), IgE, IgD or IgM.

"Active" or "activity" for the purposes herein refers to form(s) of IL-11p which retain one or more of the biologic activities of native or naturally-occurring IL-11p, or which exhibit immunological cross-reactivity with a native or naturally-occurring IL-11p.

As used herein, a "biologic activity" or "biological activity" of an IL-11p means any effector function exhibited by the IL-11p in the physiology or pathophysiology of a mammal, excluding any immunogenic or antigenic functions of the IL-11p. Immunogenic and antigenic functions of an IL-11p refer to the ability of the IL-11p to generate a humoral or cell-mediated immune response specific to the IL-11p, and the ability of the IL-11p to specifically recognize and interact with anti-IL-11p antibodies, B cells or T cells, respectively, in a mammal.

As used herein, "immunological cross-reactivity" with an IL-11p means that the candidate polypeptide is capable of competitively inhibiting the binding of the IL-11p to polyclonal or monoclonal antibodies raised against the IL-11p.

In one embodiment, IL-11p activity includes the ability to agonize or antagonize one or more biological activities of any IL-1-like family member, e.g. an IL-11p activity that antagonizes an IL-1-mediated or IL-18-mediated inflammatory response. In another embodiment, IL-11p activity includes the ability to bind to the IL-18 receptor and/or IL-1 receptor.

The term "antagonist" is used in the broadest sense, and includes any molecule that partially or fully blocks, inhibits, or neutralizes a biological activity of a native IL-11p polypeptide disclosed herein. In a similar manner, the term "agonist" is used in the broadest sense and includes any molecule that mimics a biological activity of a native IL-11p polypeptide disclosed herein. Suitable agonist or antagonist molecules specifically include agonist or antagonist antibodies or antibody fragments, fragments or amino acid sequence variants of native IL-11p polypeptides, peptides, small organic molecules, etc.

"Treatment" refers to both therapeutic treatment and prophylactic or preventative measures, wherein the object is to prevent or slow down (lessen) the targeted pathologic condition or disorder. Those in need of treatment include those already with the disorder as well as those prone to have the disorder or those in whom the disorder is to be prevented.

"Chronic" administration refers to administration of the agent(s) in a continuous mode as opposed to an acute mode, so as to maintain the initial therapeutic effect (activity) for an extended period of time. "Intermittent" administration is treatment that is not consecutively done without interruption, but rather is cyclic in nature.

"Mammal" for purposes of treatment refers to any animal classified as a mammal, including humans, domestic and farm animals, and zoo, sports, or pet animals, such as dogs, cats, cattle, horses, sheep, pigs, etc. Preferably, the mammal is human.

Administration "in combination with" one or more further therapeutic agents includes simultaneous (concurrent) and consecutive administration in any order.

The terms "inflammatory disorders" and "inflammatory diseases" are used interchangeably herein and refer to pathological states resulting in inflammation. Examples of such disorders include inflammatory skin diseases such as psoriasis and atopic dermatitis; systemic scleroderma and sclerosis; responses associated with inflammatory bowel disease (such as Crohn's disease and ulcerative colitis); ischemic reperfusion disorders including surgical tissue reperfusion injury, myocardial ischemic conditions such as myocardial infarction, cardiac arrest, reperfusion after cardiac surgery and constriction after percutaneous transluminal coronary angioplasty, stroke, and abdominal aortic aneurysms; cerebral edema secondary to stroke; cranial trauma; hypovolemic shock; asphyxia; adult respiratory distress syndrome; acute lung injury; Behcet's Disease; dermatomyositis; polymyositis; multiple sclerosis; dermatitis; meningitis; encephalitis; uveitis; osteoarthritis; autoimmune diseases such as rheumatoid arthritis, Sjorgen's syndrome, vasculitis, and insulin-dependent diabetes mellitus (IDDM); diseases involving leukocyte diapedesis; central nervous system (CNS) inflammatory disorder; meningitis; multiple organ injury syndrome secondary to septicemia or trauma; inflammatory diseases of the liver, including alcoholic hepatitis and hepatic fibrosis; pathologic host responses to infection, including pathologic inflammation in granulomatous diseases, hepatitis, and bacterial pneumonia; antigen-antibody complex mediated diseases including glomerulonephritis; sepsis; sarcoidosis; immunopathologic responses to tissue/organ transplantation, including graft-versus host disease (GVHD); inflammations of the lung, including pleurisy, alveolitis, vasculitis, pneumonia, chronic bronchitis, bronchiectasis, diffuse panbronchiolitis, hypersensitivity pneumonitis, idiopathic pulmonary fibrosis (IPF), and cystic fibrosis; inflammation in renal diseases, including acute or chronic nephritic conditions such as lupus nephritis; pancreatitis; etc. The preferred indications include rheumatoid arthritis, osteoarthritis, sepsis, acute lung injury, adult respiratory distress syndrome, idiopathic pulmonary fibrosis, ischemic reperfusion (including surgical tissue reperfusion injury, stroke, myocardial ischemia, and acute myocardial infarction), asthma, psoriasis, graft-versus-host disease (GVHD), and inflammatory bowel disease such as ulcerative colitis.

As used herein, the terms "asthma", "asthmatic disorder", "asthmatic disease", and "bronchial asthma" refer to a condition of the lungs in which there is widespread narrowing of lower airways. "Atopic asthma" and "allergic asthma" refer to asthma that is a manifestation of an IgE-mediated hypersensitivity reaction in the lower airways, including, e.g., moderate or severe chronic asthma, such as conditions requiring the frequent or constant use of inhaled or systemic steroids to control the asthma symptoms. A preferred indication is allergic asthma.

II. Detailed Description of the Invention

The present invention provides newly identified and isolated nucleotide sequences encoding polypeptides referred to in the present application as IL-11p. In particular, cDNAs encoding IL-11p polypeptides have been identified and isolated, as disclosed in further detail in the Examples below.

Using NCBI-BLAST2 sequence alignment computer programs, it has been found that a full-length native sequence hIL-1Ra1 (shown in Figure 3 and SEQ ID NO:7) has some amino acid sequence identity with human IL-1 receptor antagonist beta (hIL-1Ra β) and TANGO-77 protein, a full-length native sequence hIL-1Ra1L (shown in Figure 15 and SEQ ID NO:19) has some amino acid sequence identity with human IL-1 receptor antagonist beta (hIL-1Ra β) and TANGO-77 protein, a full-length native sequence hIL-1Ra1V (shown in Figure 19 and SEQ ID NO:25) has some amino acid sequence identity with human IL-1 receptor antagonist beta (hIL-1Ra β) and TANGO-77 protein, a full-length native sequence hIL-1Ra1S (shown in Figure 16 and SEQ ID NO:21) appears to be an allelic variant of TANGO-77 protein and has some amino acid sequence identity with human IL-1 receptor antagonist beta (hIL-1Ra β), a full-length native sequence hIL-1Ra2 (shown in Figure 5 and SEQ ID NO:10) has some amino acid sequence identity with hIL-1Ra β , a full-length native sequence hIL-1Ra3 (shown in Figure 7 and SEQ ID NO:13) has some amino acid sequence identity with human intracellular IL-1 receptor antagonist (hicIL-1Ra), and a full-length native sequence mL-1Ra3 (shown in Figure 9 and SEQ ID NO:16) has some amino acid sequence identity with mouse IL-1 receptor antagonist (mIL-1Ra) and has some amino acid sequence identity with hicIL-1Ra. hIL-1Ra β is described in EP 0855404 published July 29, 1998. TANGO-77 protein is described in WO 99/06426 published February 11, 1999. hicIL-1Ra is described in WO 95/10298 published April 20, 1995 and in Haskill et al., *Proc. Natl. Acad. Sci. (USA)*, **88**: 3681-3685 (1991). mIL-1Ra is described in Zahedi et al., *J. Immunol.*, **146**: 4228-4233 (1991), Matsushime et al., *Blood*, **78**: 616-623 (1991), Zahedi et al., *Cytokine*, **6**: 1-9 (1994), Eisenberg et al., *Proc. Natl. Acad. Sci. (USA)*, **88**: 5232-5236 (1991) and Shuck et al., *Eur. J. Immunol.*, **21**: 2775-2780 (1991). Accordingly, it is presently believed that the IL-11p polypeptides disclosed in the present application are newly identified members of the interleukin-1-like family and possess inflammatory or anti-inflammatory activities, or other cellular response activating or inhibiting activities, typical of the IL-1-like family.

In addition to the full-length native sequence IL-11p polypeptides described herein, it is contemplated that IL-11p variants can be prepared. Such embodiments of the invention include all IL-11p polypeptides that are IL-11p variants as defined herein, such as hIL-1Ra1 variants, hIL-1Ra1L variants, hIL-1Ra1S variants, hIL-1Ra2 variants, hIL-1Ra3 variants, and mL-1Ra3 variants.

IL-11p variants can be prepared by introducing appropriate nucleotide changes into the IL-11p DNA, and/or by synthesis of the desired IL-11p polypeptide. Those skilled in the art will appreciate that amino acid changes may alter post-translational processes of the IL-11p, such as changing the number or position of glycosylation sites or altering the membrane anchoring characteristics.

Variations in the native full-length sequence IL-11p or in various domains of the IL-11p described herein, can be made, for example, using any of the techniques and guidelines for conservative and non-conservative mutations set forth, for instance, in U.S. Patent No. 5,364,934. Variations may be a substitution, deletion or insertion of one or more codons encoding the IL-11p that results in a change in the amino acid sequence of the IL-11p as compared with the native sequence IL-11p. Optionally the variation is by substitution of at least one amino acid with any other amino acid in one or more of the domains of the IL-11p. Guidance in determining which amino acid residue may be inserted, substituted or deleted without adversely affecting the desired activity may be found by comparing the sequence of the IL-11p with that of homologous known protein molecules and minimizing the number of amino acid sequence changes made in regions of high homology. Amino acid substitutions can be the result of replacing one amino acid with another amino acid having similar structural and/or chemical properties, such as the replacement of a leucine with a serine, i.e., conservative amino acid replacements. Insertions or deletions may optionally be in the range of 1 to 5 amino acids. The variation allowed may be determined by systematically making insertions, deletions or substitutions of amino acids in the sequence and testing the resulting variants for activity in the *in vitro* assay described in the Examples below.

Table 1 below lists conservative amino acid substitutions (under the heading of "Preferred Substitutions") that are useful in generating variants of the native sequence IL-11p. If such substitutions result in alteration of biological activity, it is useful to introduce more substantial changes, such as the "Exemplary Substitutions" denoted in Table 1 or the substantial changes described below in reference to amino acid classes, at the active site in question.

Table 1

	Original Residue	Exemplary Substitutions	Preferred Substitutions
5	Ala (A)	val; leu; ile	val
	Arg (R)	lys; gln; asn	lys
	Asn (N)	gln; his; lys; arg	gln
	Asp (D)	glu	glu
10	Cys (C)	ser	ser
	Gln (Q)	asn	asn
	Glu (E)	asp	asp
	Gly (G)	pro; ala	ala
	His (H)	asn; gln; lys; arg	arg
15	Ile (I)	leu; val; met; ala; phe; norleucine	leu
	Leu (L)	norleucine; ile; val; met; ala; phe	ile
	Lys (K)	arg; gln; asn	arg
20	Met (M)	leu; phe; ile	leu
	Phe (F)	leu; val; ile; ala; tyr	leu
	Pro (P)	ala	ala
	Ser (S)	thr	thr
	Thr (T)	ser	ser
25	Trp (W)	tyr; phe	tyr
	Tyr (Y)	trp; phe; thr; ser	phe
	Val (V)	ile; leu; met; phe; ala; norleucine	leu

Substantial modifications in function or immunological identity of the IL-11p polypeptide are accomplished by selecting substitutions that differ significantly in their effect on maintaining (a) the structure of the polypeptide backbone in the area of the substitution, for example, as a sheet or helical conformation, (b) the charge or hydrophobicity of the molecule at the target site, or (c) the bulk of the side chain. Naturally occurring residues are divided into groups based on

common side-chain properties:

- (1) hydrophobic: norleucine, met, ala, val, leu, ile;
- (2) neutral hydrophilic: cys, ser, thr;
- (3) acidic: asp, glu;
- (4) basic: asn, gln, his, lys, arg;
- (5) residues that influence chain orientation: gly, pro; and
- (6) aromatic: trp, tyr, phe.

Non-conservative substitutions will entail exchanging a member of one of these classes for another class. Such substituted residues also may be introduced into the conservative substitution sites or, more preferably, into the remaining (non-conserved) sites.

The variations can be made using methods known in the art such as oligonucleotide-mediated (site-directed) mutagenesis, alanine scanning, and PCR mutagenesis. Site-directed mutagenesis (Carter *et al.*, *Nucl. Acids Res.*, 13:4331 (1986); Zoller *et al.*, *Nucl. Acids Res.*, 10:6487 (1987)), cassette mutagenesis (Wells *et al.*, *Gene*, 34:315 (1985)), restriction selection

mutagenesis [Wells *et al.*, *Philos. Trans. R. Soc. London SerA*, 317:415 (1986)] or other known techniques can be performed on the cloned DNA to produce the IL-11p variant DNA.

Scanning amino acid analysis can also be employed to identify one or more amino acids along a contiguous sequence. Among the preferred scanning amino acids are relatively small, neutral amino acids. Such amino acids include alanine, glycine, serine, and cysteine. Alanine is typically a preferred scanning amino acid among this group because it eliminates the side-chain beyond the beta-carbon and is less likely to alter the main-chain conformation of the variant [Cunningham and Wells, *Science*, 244: 1081-1085 (1989)]. Alanine is also typically preferred because it is the most common amino acid. Further, it is frequently found in both buried and exposed positions [Creighton, *The Proteins*, (W.H. Freeman & Co., N.Y.); Chothia, *J. Mol. Biol.*, 150:1 (1976)]. If alanine substitution does not yield adequate amounts of variant, an isoteric amino acid can be used.

Covalent modifications of IL-11p are included within the scope of this invention. One type of covalent modification includes reacting targeted amino acid residues of an IL-11p polypeptide with an organic derivatizing agent that is capable of reacting with selected side chains or the N- or C- terminal residues of the IL-11p. Derivatization with bifunctional agents is useful, for instance, for crosslinking IL-11p to a water-insoluble support matrix or surface for use in the method for purifying anti-IL-11p antibodies, and vice-versa. Commonly used crosslinking agents include, *e.g.*, 1,1-bis(diazoacetyl)-2-phenylethane, glutaraldehyde, N-hydroxysuccinimide esters, for example, esters with 4-azidosalicylic acid, homobifunctional imidoesters, including disuccinimidyl esters such as 3,3'-dithiobis(succinimidylpropionate), bifunctional maleimides such as bis-N-maleimido-1,8-octane and agents such as methyl-3-[(p-azidophenyl)dithio]propionimide.

Other modifications include deamidation of glutaminyl and asparaginyl residues to the corresponding glutamyl and aspartyl residues, respectively, hydroxylation of proline and lysine, phosphorylation of hydroxyl groups of seryl or threonyl residues, methylation of the α -amino groups of lysine, arginine, and histidine side chains [T.E. Creighton, *Proteins: Structure and Molecular Properties*, W.H. Freeman & Co., San Francisco, pp. 79-86 (1983)], acetylation of the N-terminal amine, and amidation of any C-terminal carboxyl group.

Another type of covalent modification of the IL-11p polypeptide included within the scope of this invention comprises altering the native glycosylation pattern of the polypeptide. "Altering the native glycosylation pattern" is intended for purposes herein to mean deleting one or more carbohydrate moieties found in native sequence IL-11p (either by removing the underlying glycosylation site or by deleting the glycosylation by chemical and/or enzymatic means), and/or adding one or more glycosylation sites that are not present in the native sequence IL-11p. In addition, the phrase includes qualitative changes in the glycosylation of the native proteins, involving a change in the nature and proportions of the various carbohydrate moieties present.

Addition of glycosylation sites to the IL-11p polypeptide may be accomplished by altering the amino acid sequence. The alteration may be made, for example, by the addition of, or substitution by, one or more serine or threonine residues to the native sequence IL-11p (for O-

linked glycosylation sites). The IL-11p amino acid sequence may optionally be altered through changes at the DNA level, particularly by mutating the DNA encoding the IL-11p polypeptide at preselected bases such that codons are generated that will translate into the desired amino acids.

Another means of increasing the number of carbohydrate moieties on the IL-11p polypeptide is by chemical or enzymatic coupling of glycosides to the polypeptide. Such methods are described in the art, e.g., in WO 87/05330 published 11 September 1987, and in Aplin and Wriston, CRC Crit. Rev. Biochem., pp. 259-306 (1981).

Removal of carbohydrate moieties present on the IL-11p polypeptide may be accomplished chemically or enzymatically or by mutational substitution of codons encoding for amino acid residues that serve as targets for glycosylation. Chemical deglycosylation techniques are known in the art and described, for instance, by Hakimuddin, *et al.*, Arch. Biochem. Biophys., 259:52 (1987) and by Edge *et al.*, Anal. Biochem., 118:131 (1981). Enzymatic cleavage of carbohydrate moieties on polypeptides can be achieved by the use of a variety of endo- and exo-glycosidases as described by Thotakura *et al.*, Meth. Enzymol., 138:350 (1987).

Another type of covalent modification of IL-11p comprises linking the IL-11p polypeptide to one of a variety of nonproteinaceous polymers, e.g., polyethylene glycol (PEG), polypropylene glycol, or polyoxyalkylenes, in the manner set forth in U.S. Patent Nos. 4,640,835; 4,496,689; 4,301,144; 4,670,417; 4,791,192 or 4,179,337.

The IL-11p of the present invention may also be modified in a way to form a chimeric molecule comprising IL-11p fused to another, heterologous polypeptide or amino acid sequence.

In one embodiment, such a chimeric molecule comprises a fusion of the IL-11p with a tag polypeptide which provides an epitope to which an anti-tag antibody can selectively bind. The epitope tag is generally placed at the amino- or carboxyl- terminus of the IL-11p. The presence of such epitope-tagged forms of the IL-11p can be detected using an antibody against the tag polypeptide. Also, provision of the epitope tag enables the IL-11p to be readily purified by affinity purification using an anti-tag antibody or another type of affinity matrix that binds to the epitope tag. Various tag polypeptides and their respective antibodies are well known in the art. Examples include poly-histidine (poly-his) or poly-histidine-glycine (poly-his-gly) tags; the flu HA tag polypeptide and its antibody 12CA5 [Field *et al.*, Mol. Cell. Biol., 8:2159-2165 (1988)]; the c-myc tag and the 8F9, 3C7, 6E10, G4, B7 and 9E10 antibodies thereto [Evan *et al.*, Molecular and Cellular Biology, 5:3610-3616 (1985)]; and the Herpes Simplex virus glycoprotein D (gD) tag and its antibody [Paborsky *et al.*, Protein Engineering, 3(6):547-553 (1990)]. Other tag polypeptides include the Flag-peptide [Hopp *et al.*, BioTechnology, 6:1204-1210 (1988)]; the KT3 epitope peptide [Martin *et al.*, Science, 255:192-194 (1992)]; an α -tubulin epitope peptide [Skinner *et al.*, J. Biol. Chem., 266:15163-15166 (1991)]; and the T7 gene 10 protein peptide tag [Lutz-Freyermuth *et al.*, Proc. Natl. Acad. Sci. USA, 87:6393-6397 (1990)].

In an alternative embodiment, the chimeric molecule may comprise a fusion of the IL-11p with an immunoglobulin or a particular region of an immunoglobulin. For a bivalent form of the chimeric molecule (also referred to as an immunoadhesin), such a fusion could be to the Fc

region of an IgG molecule. The Ig fusions preferably include the substitution of a soluble form of an IL-11p polypeptide in place of at least one variable region within an Ig molecule. In a particularly preferred embodiment, the immunoglobulin fusion includes the hinge, CH2 and CH3, or the hinge, CH1, CH2 and CH3 regions of an IgG1 molecule. For the production of immunoglobulin fusions see also US Patent No. 5,428,130 issued June 27, 1995.

In one aspect, the invention provides an isolated nucleic acid comprising DNA encoding an IL-11p polypeptide that retains at least one biologic activity of a native sequence IL-11p, such as the IL-1R binding activity of a native sequence hIL-1Ra3 or mL-1Ra3, or the IL-18R binding activity of a native sequence hIL-1Ra1, hIL-1Ra1L, or hIL-1Ra1V, which DNA has at least at or about 80% sequence identity, or at least at or about 85% sequence identity, or at least at or about 90% sequence identity, or at least at or about 95% sequence identity to (a) a DNA molecule selected from the group consisting of: (1) a DNA molecule encoding an IL-11p polypeptide comprising amino acid residues from at or about 37 to at or about 203, inclusive of Figure 2 (SEQ ID NO:5), (2) a DNA molecule encoding an IL-11p polypeptide comprising amino acid residues from at or about 15 to at or about 193, inclusive of Figure 3 (SEQ ID NO:7), (3) a DNA molecule encoding an IL-11p polypeptide comprising amino acid residues from at or about 34 to at or about 155, inclusive of Figure 7 (SEQ ID NO:13), (4) a DNA molecule encoding an IL-11p polypeptide comprising amino acid residues from at or about 34 to at or about 155, inclusive of Figure 9 (SEQ ID NO:16), (5) a DNA molecule encoding an IL-11p polypeptide comprising amino acid residues from at or about 26 to at or about 207, inclusive of Figure 15 (SEQ ID NO:19), and (6) a DNA molecule encoding an IL-11p polypeptide comprising amino acid residues from at or about 46 to at or about 218 of Figure 19 (SEQ ID NO:25), or (b) the complement of the DNA molecule of (a).

In another aspect, the invention provides an isolated nucleic acid comprising DNA encoding an IL-11p polypeptide that retains at least one biologic activity of a native sequence IL-11p, such as the IL-18R binding activity of a native sequence hIL-1Ra1, hIL-1Ra1L, or hIL-1Ra1V, which DNA has at least at or about 80% sequence identity, or at least at or about 85% sequence identity, or at least at or about 90% sequence identity, or at least at or about 95% sequence identity to (a) a DNA molecule selected from the group consisting of: (1) a DNA molecule encoding an IL-11p polypeptide comprising amino acid residues from at or about 1 to at or about 207, inclusive of Figure 15 (SEQ ID NO:19), and (2) a DNA molecule encoding an IL-11p polypeptide comprising amino acid residues from at or about 1 to at or about 218 of Figure 19 (SEQ ID NO:25), or (b) the complement of the DNA molecule of (a).

In another aspect, the invention provides an isolated nucleic acid comprising DNA encoding an IL-11p polypeptide that retains at least one biologic activity of a native sequence IL-11p, such as the IL-18R binding activity of a native sequence hIL-1Ra1, or the IL-1R binding activity of a native sequence hIL-1Ra3 or mL-1Ra3, which DNA has at least at or about 80% sequence identity, or at least at or about 85% sequence identity, or at least at or about 90% sequence identity, or at least at or about 95% sequence identity to (a) a DNA molecule selected from the group consisting of: (1) a DNA molecule encoding an IL-11p polypeptide comprising amino acid residues from at or about 95 to at or about 134, inclusive of Figure 7 (SEQ ID NO:13),

and (2) a DNA molecule encoding an IL-11p polypeptide comprising amino acid residues from at or about 95 to at or about 134, inclusive of Figure 9 (SEQ ID NO:16), or (b) the complement of the DNA molecule of (a).

In another aspect, the invention provides an isolated nucleic acid comprising DNA encoding an IL-11p polypeptide that retains at least one biologic activity of a native sequence IL-11p, such as the IL-18R binding activity of a native sequence hIL-1Ra1, or the IL-1R binding activity of a native sequence hIL-1Ra3 or mIL-1Ra3, which DNA has at least at or about 80% sequence identity, or at least at or about 85% sequence identity, or at least at or about 90% sequence identity, or at least at or about 95% sequence identity to (a) a DNA molecule encoding an IL-11p polypeptide comprising amino acid residues from at or about 80 to at or about 155, inclusive of Figure 7 (SEQ ID NO:13), or (b) the complement of the DNA molecule of (a).

In another aspect, the invention provides an isolated nucleic acid comprising DNA encoding an IL-11p polypeptide that retains at least one biologic activity of a native sequence IL-11p, such as the IL-18R binding activity of a native sequence hIL-1Ra1, or the IL-1R binding activity of a native sequence hIL-1Ra3 or mIL-1Ra3, which DNA has at least at or about 80% sequence identity, or at least at or about 85% sequence identity, or at least at or about 90% sequence identity, or at least at or about 95% sequence identity to (a) a DNA molecule selected from the group consisting of: (1) a DNA molecule encoding an IL-11p polypeptide comprising amino acid residues from at or about 2 to at or about 155, inclusive of Figure 7 (SEQ ID NO:13), and (2) a DNA molecule encoding an IL-11p polypeptide comprising amino acid residues from at or about 2 to at or about 155, inclusive of Figure 9 (SEQ ID NO:16), or (b) the complement of the DNA molecule of (a).

In another aspect, the invention provides an isolated nucleic acid comprising DNA having at least at or about 80% sequence identity, or at least at or about 85% sequence identity, or at least at or about 90% sequence identity, or at least at or about 95% sequence identity to (a) a DNA molecule selected from the group consisting of: (1) a DNA molecule encoding an IL-11p polypeptide comprising amino acid residues from at or about 95 to at or about 134, inclusive of Figure 7 (SEQ ID NO:13), and (2) a DNA molecule encoding an IL-11p polypeptide comprising amino acid residues from at or about 95 to at or about 134, inclusive of Figure 9 (SEQ ID NO:16), or (b) the complement of the DNA molecule of (a).

In another aspect, the invention provides an isolated nucleic acid comprising DNA having at least at or about 80% sequence identity, or at least at or about 85% sequence identity, or at least at or about 90% sequence identity, or at least at or about 95% sequence identity to (a) a DNA molecule encoding an IL-11p polypeptide comprising amino acid residues from at or about 80 to at or about 155, inclusive of Figure 7 (SEQ ID NO:13), or (b) the complement of the DNA molecule of (a).

In another aspect, the invention provides an isolated nucleic acid comprising DNA having at least at or about 80% sequence identity, or at least at or about 85% sequence identity, or at least at or about 90% sequence identity, or at least at or about 95% sequence identity to (a) a DNA molecule selected from the group consisting of: (1) a DNA molecule encoding an IL-11p polypeptide comprising amino acid residues from at or about 2 to at or about 155, inclusive of

Figure 7 (SEQ ID NO:13), and (2) a DNA molecule encoding an IL-11p polypeptide comprising amino acid residues from at or about 2 to at or about 155, inclusive of Figure 9 (SEQ ID NO:16), or (b) the complement of the DNA molecule of (a).

In another aspect, the invention concerns an isolated nucleic acid molecule encoding an IL-11p polypeptide, comprising DNA hybridizing to the complement of a nucleic acid sequence selected from the group consisting of: (1) the nucleic acid sequence consisting of nucleotide positions from at or about 238 to at or about 465 in the sense strand of Figure 7 (SEQ ID NO:12); (2) the nucleic acid sequence consisting of nucleotide positions from at or about 427 to at or about 609 in the sense strand of Figure 9 (SEQ ID NO:15); and (3) the nucleic acid sequence consisting of nucleotide positions from at or about 79 to at or about 135 in the sense strand of Figure 15 (SEQ ID NO:18). Preferably, hybridization occurs under stringent hybridization and wash conditions.

In another aspect, the invention concerns an isolated nucleic acid molecule, comprising DNA that is at least 90 nucleotides in length and that hybridizes to the complement of a nucleic acid sequence selected from the group consisting of: (1) the nucleic acid sequence consisting of nucleotide positions from at or about 238 to at or about 465 in the sense strand of Figure 7 (SEQ ID NO:12); (2) the nucleic acid sequence consisting of nucleotide positions from at or about 427 to at or about 609 in the sense strand of Figure 9 (SEQ ID NO:15); and (3) the nucleic acid sequence consisting of nucleotide positions from at or about 115 to at or about 135 in the sense strand of Figure 15 (SEQ ID NO:18). Preferably, hybridization occurs under stringent hybridization and wash conditions.

In another aspect, the invention concerns an isolated nucleic acid molecule comprising DNA encoding an IL-11p polypeptide that retains at least one biologic activity of a native sequence IL-11p, such as the IL-1R binding activity of a native sequence hIL-1Ra3 or mL-1Ra3, or the IL-18R binding activity of a native sequence hIL-1Ra1, hIL-1Ra1L, or hIL-1Ra1V, which DNA hybridizes to the complement of a nucleic acid sequence selected from the group consisting of: (1) the nucleic acid sequence consisting of nucleotide positions from at or about 118 to at or about 231 in the sense strand of Figure 2 (SEQ ID NO:4); (2) the nucleic acid sequence consisting of nucleotide positions from at or about 100 to at or about 465 in the sense strand of Figure 7 (SEQ ID NO:12); (3) the nucleic acid sequence consisting of nucleotide positions from at or about 244 to at or about 609 in the sense strand of Figure 9 (SEQ ID NO:15); and (4) the nucleic acid sequence consisting of nucleotide positions from at or about 208 to at or about 339 in the sense strand of Figure 19 (SEQ ID NO:24). Preferably, hybridization occurs under stringent hybridization and wash conditions.

In another aspect, the invention concerns an isolated nucleic acid molecule comprising DNA encoding an IL-11p polypeptide that retains at least one biologic activity of a native sequence IL-11p, such as the IL-1R binding activity of a native sequence hIL-1Ra3 or mL-1Ra3, which DNA hybridizes to the complement of a nucleic acid sequence selected from the group consisting of: (1) the nucleic acid sequence consisting of nucleotide positions from at or about 4 to at or about 465 in the sense strand of Figure 7 (SEQ ID NO:12); and (2) the nucleic acid sequence consisting of nucleotide positions from at or about 148 to at or about 609 in the sense strand

of Figure 9 (SEQ ID NO:15). Preferably, hybridization occurs under stringent hybridization and wash conditions.

In a further aspect, the invention concerns an isolated nucleic acid molecule comprising DNA encoding an IL-11p polypeptide that retains at least one biologic activity of a native sequence IL-11p, such as the IL-1R binding activity of a native sequence hIL-1Ra3 or mL-1Ra3, or the IL-18R binding activity of a native sequence hIL-1Ra1, hIL-1Ra1L, or hIL-1Ra1V, which DNA has at least at or about 80% sequence identity, or at least at or about 85% sequence identity, or at least at or about 90% sequence identity, or at least at or about 95% sequence identity to (a) a DNA encoding an IL-11p, such as a mature IL-11p polypeptide, encoded by the cDNA insert in the vector deposited as ATCC Deposit No. 203588 (DNA85066-2534), ATCC Deposit No. 203587 (DNA96786-2534), ATCC Deposit No. 203589 (DNA96787-2534), ATCC Deposit No. 203590 (DNA92505-2534), ATCC Deposit No. 203846 (DNA102043-2534), or ATCC Deposit No. 203973 (DNA114876-2534), or (b) the complement of the DNA molecule of (a). In a preferred embodiment, the nucleic acid comprises a DNA encoding an IL-11p polypeptide, such as a mature IL-11p polypeptide, encoded by the cDNA insert in the vector deposited as ATCC Deposit No. 203588 (DNA85066-2534), ATCC Deposit No. 203587 (DNA96786-2534), ATCC Deposit No. 203586 (DNA92929-2534), ATCC Deposit No. 203589 (DNA96787-2534), ATCC Deposit No. 203590 (DNA92505-2534), ATCC Deposit No. 203846 (DNA102043-2534), ATCC Deposit No. 203973 (DNA114876-2534), or ATCC Deposit No. 203855 (DNA102044-2534).

In a further aspect, the invention concerns an isolated nucleic acid molecule comprising DNA encoding an IL-11p polypeptide that retains at least one biologic activity of a native sequence IL-11p, such as the IL-1R binding activity of a native sequence hIL-1Ra3 or mL-1Ra3, or the IL-18R binding activity of a native sequence hIL-1Ra1, hIL-1Ra1L, or hIL-1Ra1V, which DNA has at least at or about 80% sequence identity, or at least at or about 85% sequence identity, or at least at or about 90% sequence identity, or at least at or about 95% sequence identity to (a) a DNA encoding the entire amino acid sequence encoded by the longest open reading frame in the cDNA insert of a vector selected from the group consisting of the vectors deposited as ATCC Deposit No. 203588 (DNA85066-2534), ATCC Deposit No. 203587 (DNA96786-2534), ATCC Deposit No. 203589 (DNA96787-2534), ATCC Deposit No. 203590 (DNA92505-2534), ATCC Deposit No. 203846 (DNA102043-2534), and ATCC Deposit No. 203973 (DNA114876-2534), or (b) the complement of the DNA molecule of (a). In a preferred embodiment, the nucleic acid comprises (a) DNA encoding the entire amino acid sequence encoded by the longest open reading frame in the cDNA insert of a vector selected from the group consisting of the vectors deposited as ATCC Deposit No. 203588 (DNA85066-2534), ATCC Deposit No. 203587 (DNA96786-2534), ATCC Deposit No. 203586 (DNA92929-2534), ATCC Deposit No. 203589 (DNA96787-2534), and ATCC Deposit No. 203590 (DNA92505-2534), or (b) the complement of the DNA of (a). In another preferred embodiment, the nucleic acid comprises (a) DNA encoding the entire amino acid sequence encoded by the longest open reading frame in the cDNA insert of a vector selected from the group consisting of the vectors deposited as ATCC Deposit No. 203846 (DNA102043-2534), ATCC Deposit No. 203855 (DNA102044-2534), and ATCC Deposit No. 203973 (DNA114876-2534), or (b) the complement of the DNA of (a).

In another aspect, the invention concerns an isolated nucleic acid molecule comprising DNA encoding an IL-11p polypeptide that retains at least one biologic activity of a native sequence IL-11p, such as the IL-1R binding activity of a native sequence IL-1Ra3 or mL-1Ra3, or the IL-18R binding activity of a native sequence hIL-1Ra1, hIL-1Ra1L, or hIL-1Ra1V, which DNA has at least at or about 80% sequence identity, or at least at or about 85% sequence identity, or at least at or about 90% sequence identity, or at least at or about 95% sequence identity to (a) DNA encoding an amino acid sequence selected from the group consisting of: (1) the entire amino acid sequence encoded by the longest open reading frame in the cDNA insert in the vector deposited as ATCC Deposit No. 203588, (2) the entire amino acid sequence, or the entire amino acid sequence excluding the 36 N-terminal amino acid residues of such sequence, encoded by the longest open reading frame in the cDNA insert in the vector deposited as ATCC Deposit No. 203587, (3) the entire amino acid sequence, or the entire amino acid sequence excluding the N-terminal amino acid residue of such sequence, encoded by the longest open reading frame in the cDNA insert in the vector deposited as ATCC Deposit No. 203589, (4) the entire amino acid sequence, or the entire amino acid sequence excluding the N-terminal amino acid residue of such sequence, encoded by the longest open reading frame in the cDNA insert in the vector deposited as ATCC Deposit No. 203590, (5) the entire amino acid sequence, or the entire amino acid sequence excluding the N-terminal amino acid residue of such sequence, or the entire amino acid sequence excluding the 34 N-terminal amino acid residues of such sequence, encoded by the longest open reading frame in the cDNA insert in the vector deposited as ATCC Deposit No. 203846, and (6) the entire amino acid sequence, or the entire amino acid sequence excluding the N-terminal amino acid residue of such sequence, or the entire amino acid sequence excluding the 45 N-terminal amino acid residues of such sequence, encoded by the longest open reading frame in the cDNA insert in the vector deposited as ATCC Deposit No. 203973, or (b) the complement of the DNA of (a).

In a preferred embodiment, the nucleic acid comprises (a) DNA encoding an amino acid sequence selected from the group consisting of: (1) the entire amino acid sequence encoded by the longest open reading frame in the cDNA insert in the vector deposited as ATCC Deposit No. 203588, (2) the entire amino acid sequence, or the entire amino acid sequence excluding the 36 N-terminal amino acid residues of such sequence, encoded by the longest open reading frame in the cDNA insert in the vector deposited as ATCC Deposit No. 203587, (3) the entire amino acid sequence, or the entire amino acid sequence excluding the N-terminal amino acid residue of such sequence, encoded by the longest open reading frame in the cDNA insert in the vector deposited as ATCC Deposit No. 203589, (4) the entire amino acid sequence, or the entire amino acid sequence excluding the N-terminal amino acid residue of such sequence, encoded by the longest open reading frame in the cDNA insert in the vector deposited as ATCC Deposit No. 203590, (5) the entire amino acid sequence, or the entire amino acid sequence excluding the N-terminal amino acid residue of such sequence, or the entire amino acid sequence excluding the 34 N-terminal amino acid residues of such sequence, encoded by the longest open reading frame in the cDNA insert in the vector deposited as ATCC Deposit No. 203846, and (6) the entire amino acid sequence, or the entire amino acid sequence excluding the N-terminal amino acid residue

of such sequence, or the entire amino acid sequence excluding the 45 N-terminal amino acid residues of such sequence, encoded by the longest open reading frame in the cDNA insert in the vector deposited as ATCC Deposit No. 203973, or (b) the complement of the DNA of (a).

In a still further aspect, the invention concerns an isolated nucleic acid molecule comprising (a) DNA encoding an IL-11p polypeptide that retains at least one biologic activity of a native sequence IL-11p, such as the IL-18R binding activity of a native sequence hIL-1Ra1, or the IL-1R binding activity of a native sequence hIL-1Ra3 or mIL-1Ra3, and which IL-11p polypeptide has at least at or about 80% sequence identity, or at least at or about 85% sequence identity, or at least at or about 90% sequence identity, or at least at or about 95% sequence identity to an amino acid sequence selected from the group consisting of: (1) amino acid residues from at or about 37 to at or about 203, inclusive of Figure 2 (SEQ ID NO:5), (2) amino acid residues from at or about 15 to at or about 193, inclusive of Figure 3 (SEQ ID NO:7), (3) amino acid residues from at or about 34 to at or about 155, inclusive of Figure 7 (SEQ ID NO:13), (4) amino acid residues from at or about 34 to at or about 155, inclusive of Figure 9 (SEQ ID NO:16), (5) amino acid residues from at or about 26 to at or about 207, inclusive of Figure 15 (SEQ ID NO:19), and (6) amino acid residues from at or about 46 to at or about 218, inclusive of Figure 19 (SEQ ID NO:25), or (b) the complement of the DNA of (a).

In a still further aspect, the invention concerns an isolated nucleic acid molecule comprising (a) DNA encoding an IL-11p polypeptide that retains at least one biologic activity of a native sequence IL-11p, such as the IL-18R binding activity of a native sequence hIL-1Ra1, or the IL-1R binding activity of a native sequence hIL-1Ra3 or mIL-1Ra3, and which IL-11p polypeptide has at least at or about 80% sequence identity, or at least at or about 85% sequence identity, or at least at or about 90% sequence identity, or at least at or about 95% sequence identity to an amino acid sequence selected from the group consisting of: (1) amino acid residues from at or about 95 to at or about 134, inclusive of Figure 7 (SEQ ID NO:13), and (2) amino acid residues from at or about 95 to at or about 134, inclusive of Figure 9 (SEQ ID NO:16), or (b) the complement of the DNA of (a).

In a still further aspect, the invention concerns an isolated nucleic acid molecule comprising (a) DNA encoding an IL-11p polypeptide that retains at least one biologic activity of a native sequence IL-11p, such as the IL-18R binding activity of a native sequence hIL-1Ra1, or the IL-1R binding activity of a native sequence hIL-1Ra3 or mIL-1Ra3, and which IL-11p polypeptide has at least at or about 80% sequence identity, or at least at or about 85% sequence identity, or at least at or about 90% sequence identity, or at least at or about 95% sequence identity to the amino acid sequence of amino acid residues from at or about 80 to at or about 155, inclusive of Figure 7 (SEQ ID NO:13), or (b) the complement of the DNA of (a).

In a still further aspect, the invention concerns an isolated nucleic acid molecule comprising (a) DNA encoding an IL-11p polypeptide that retains at least one biologic activity of a native sequence IL-11p, such as the IL-18R binding activity of a native sequence hIL-1Ra1, or the IL-1R binding activity of a native sequence hIL-1Ra3 or mIL-1Ra3, and which IL-11p polypeptide has at least at or about 80% sequence identity, or at least at or about 85% sequence identity, or at least at or about 90% sequence identity, or at least at or about 95% sequence

identity to an amino acid sequence selected from the group consisting of: (1) amino acid residues from at or about 2 to at or about 155, inclusive of Figure 7 (SEQ ID NO:13), and (2) amino acid residues from at or about 2 to at or about 155, inclusive of Figure 9 (SEQ ID NO:16), or (b) the complement of the DNA of (a).

5 In a still further aspect, the invention concerns an isolated nucleic acid molecule comprising (a) DNA encoding an IL-11p polypeptide having at least at or about 80% sequence identity, or at least at or about 85% sequence identity, or at least at or about 90% sequence identity, or at least at or about 95% sequence identity to an amino acid sequence selected from the group consisting of: (1) amino acid residues from at or about 95 to at or about 134, inclusive
10 of Figure 7 (SEQ ID NO:13), and (2) amino acid residues from at or about 95 to at or about 134, inclusive of Figure 9 (SEQ ID NO:16), or (b) the complement of the DNA of (a).

In a still further aspect, the invention concerns an isolated nucleic acid molecule comprising (a) DNA encoding an IL-11p polypeptide having at least at or about 80% sequence identity, or at least at or about 85% sequence identity, or at least at or about 90% sequence identity, or at least at or about 95% sequence identity to the amino acid sequence of amino acid residues from at or about 80 to at or about 155, inclusive of Figure 7 (SEQ ID NO:13), or (b) the complement of the DNA of (a).
15

In a still further aspect, the invention concerns an isolated nucleic acid molecule comprising (a) DNA encoding an IL-11p polypeptide having at least at or about 80% sequence identity, or at least at or about 85% sequence identity, or at least at or about 90% sequence identity, or at least at or about 95% sequence identity to an amino acid sequence selected from the group consisting of: (1) amino acid residues from at or about 2 to at or about 155, inclusive of Figure 7 (SEQ ID NO:13), and (2) amino acid residues from at or about 2 to at or about 155, inclusive of Figure 9 (SEQ ID NO:16), or (b) the complement of the DNA of (a).
20

25 In a still further aspect, the invention concerns an isolated nucleic acid molecule comprising (a) DNA encoding an IL-11p polypeptide that retains at least one biologic activity of a native sequence IL-11p, such as the IL-18R binding activity of a native sequence hIL-1Ra1, hIL-1Ra1L, or hIL-1Ra1V, and which IL-11p polypeptide has at least at or about 80% sequence identity, or at least at or about 85% sequence identity, or at least at or about 90% sequence identity, or at least at or about 95% sequence identity to an amino acid sequence selected from the group consisting of: (1) amino acid residues from at or about 1 to at or about 207, inclusive of Figure 15 (SEQ ID NO:19), and (2) amino acid residues from at or about 1 to at or about 218, inclusive of Figure 19 (SEQ ID NO:25), or (b) the complement of the DNA of (a).
30

In a further aspect, the invention concerns an isolated nucleic acid molecule comprising
35 DNA encoding an IL-11p that retains at least one biologic activity of a native sequence IL-11p, such as the IL-1R binding activity of a native sequence hIL-1Ra3 or mL-1Ra3, or the IL-18R binding activity of a native sequence hIL-1Ra1, hIL-1Ra1L, hIL-1Ra1S, or hIL-1Ra1V, which DNA is produced by hybridizing a test DNA molecule under stringent conditions with (a) a DNA molecule encoding an IL-11p polypeptide selected from the group consisting of: (1) an IL-11p polypeptide comprising the sequence of amino acid residues from at or about 37 to at or about
40 203, inclusive of Figure 2 (SEQ ID NO:5), (2) an IL-11p polypeptide comprising the sequence of

amino acid residues from at or about 15 to at or about 193, inclusive of Figure 3 (SEQ ID NO:7), (3) an IL-11p polypeptide comprising the sequence of amino acid residues from at or about 34 to at or about 155, inclusive of Figure 7 (SEQ ID NO:13), (4) an IL-11p polypeptide comprising the sequence of amino acid residues from at or about 34 to at or about 155, inclusive of Figure 9 (SEQ ID NO:16), (5) an IL-11p polypeptide comprising the sequence of amino acid residues from at or about 26 to at or about 207, inclusive of Figure 15 (SEQ ID NO:19), and (6) an IL-11p polypeptide comprising the sequence of amino acid residues from at or about 46 to at or about 218, inclusive of Figure 19 (SEQ ID NO:25), or (b) the complement of the DNA molecule of (a), and, if the test DNA molecule encodes an IL-11p that retains at least one biologic activity of a native sequence IL-11p, such as the IL-1R binding activity of a native sequence hIL-1Ra3 or mL-1Ra3, or the IL-18R binding activity of a native sequence hIL-1Ra1, hIL-1Ra1L, or hIL-1Ra1V, and if the test DNA molecule has at least at or about an 80% sequence identity, or at least at or about an 85% sequence identity, or at least at or about a 90% sequence identity, or at least at or about a 95% sequence identity to the DNA molecule of (a) or (b), isolating the test DNA molecule.

In another aspect, the invention provides an isolated nucleic acid molecule comprising (a) DNA encoding an a polypeptide, such as IL-11p polypeptide, selected from the group consisting of: (1) a polypeptide, such as an hIL-1Ra1 polypeptide, comprising amino acid residues from at or about 37 to at or about 63, inclusive of Figure 2 (SEQ ID NO:5); (2) a polypeptide, such as an hIL-1Ra1 polypeptide, comprising amino acid residues from at or about 15 to at or about 53, inclusive of Figure 3 (SEQ ID NO:7); (3) a polypeptide, such as an hIL-1Ra2 polypeptide, comprising amino acid residues from at or about 1 to at or about 134, inclusive of Figure 5 (SEQ ID NO:10); (4) a polypeptide comprising amino acid residues from at or about 10 to at or about 134, inclusive of Figure 5 (SEQ ID NO:10); (5) a polypeptide, such as an hIL-1Ra2 polypeptide, consisting of amino acid residues from at or about 27 to at or about 134, inclusive of Figure 5 (SEQ ID NO:10); (6) a polypeptide, such as an hIL-1Ra2 fusion variant polypeptide, consisting of a native amino acid sequence of hIL-1Ra2 consisting of amino acid residues from at or about 27 to at or about 134, inclusive of Figure 5 (SEQ ID NO:10) fused at its N-terminus or C-terminus to a heterologous amino acid or amino acid sequence; (7) a polypeptide, such as an hIL-1Ra3 polypeptide, comprising amino acid residues from at or about 95 to at or about 134, inclusive of Figure 7 (SEQ ID NO:13); and (8) a polypeptide, such as a mL-1Ra3 polypeptide, comprising amino acid residues from at or about 95 to at or about 134, inclusive of Figure 9 (SEQ ID NO:16); or (b) the complement of the DNA of (a).

In another aspect, the invention provides an isolated nucleic acid molecule comprising (a) DNA encoding a polypeptide, such as an IL-11p polypeptide, selected from the group consisting of: (1) a polypeptide, such as an hIL-1Ra1L polypeptide, comprising amino acid residues from at or about 26 to at or about 44, inclusive of Figure 15 (SEQ ID NO:19); (2) a polypeptide, such as an hIL-1Ra1L polypeptide, comprising amino acid residues from at or about 1 to at or about 44, inclusive of Figure 15 (SEQ ID NO:19); (3) a polypeptide, such as an hIL-1Ra1L polypeptide, comprising amino acid residues from at or about 26 to at or about 78, inclusive of Figure 15 (SEQ ID NO:19); (4) a polypeptide, such as an hIL-1Ra1L polypeptide,

comprising amino acid residues from at or about 1 to at or about 78, inclusive of Figure 15 (SEQ ID NO:19); (5) a polypeptide, such as an hIL-1Ra1S polypeptide, comprising amino acid residues from at or about 1 to at or about 38, inclusive of Figure 16 (SEQ ID NO:21); (6) a polypeptide, such as an hIL-1Ra1V polypeptide, comprising amino acid residues from at or about 37 to at or about 55, inclusive of Figure 19 (SEQ ID NO:25); (7) a polypeptide, such as an hIL-1Ra1V polypeptide, comprising amino acid residues from at or about 12 to at or about 55, inclusive of Figure 19 (SEQ ID NO:25); (8) a polypeptide, such as an hIL-1Ra1V polypeptide, comprising amino acid residues from at or about 1 to at or about 55, inclusive of Figure 19 (SEQ ID NO:25); (9) a polypeptide, such as an hIL-1Ra1V polypeptide, comprising amino acid residues from at or about 46 to at or about 55, inclusive of Figure 19 (SEQ ID NO:25); (10) a polypeptide, such as an hIL-1Ra1V polypeptide, comprising amino acid residues from at or about 46 to at or about 89, inclusive of Figure 19 (SEQ ID NO:25); (11) a polypeptide, such as an hIL-1Ra1V polypeptide, comprising amino acid residues from at or about 37 to at or about 89, inclusive of Figure 19 (SEQ ID NO:25); (12) a polypeptide, such as an hIL-1Ra1V polypeptide, comprising amino acid residues from at or about 12 to at or about 89, inclusive of Figure 19 (SEQ ID NO:25); and (13) a polypeptide, such as an hIL-1Ra1V polypeptide, comprising amino acid residues from at or about 1 to at or about 89, inclusive of Figure 19 (SEQ ID NO:25); or (b) the complement of the DNA of (a).

In another aspect, the invention provides an isolated nucleic acid molecule comprising (a) DNA encoding a polypeptide, such as an hIL-1Ip polypeptide, selected from the group consisting of: (1) a polypeptide, such as an hIL-1Ra1L polypeptide, consisting of a native amino acid sequence of hIL-1Ra1L consisting of amino acid residues from at or about 26 to at or about 207, inclusive of Figure 15 (SEQ ID NO:19) fused at its N-terminus or C-terminus to a heterologous amino acid or amino acid sequence; (2) a polypeptide, such as an hIL-1Ra1L polypeptide, consisting of a native amino acid sequence of hIL-1Ra1L consisting of amino acid residues from at or about 1 to at or about 207, inclusive of Figure 15 (SEQ ID NO:19) fused at its N-terminus or C-terminus to a heterologous amino acid or amino acid sequence; (3) a polypeptide, such as an hIL-1Ra1L polypeptide, consisting of amino acid residues from at or about 26 to at or about 207, inclusive of Figure 15 (SEQ ID NO:19); (4) a polypeptide, such as an hIL-1Ra1L polypeptide, consisting of amino acid residues from at or about 1 to at or about 207, inclusive of Figure 15 (SEQ ID NO:19); (5) a polypeptide, such as an hIL-1Ra1S fusion variant polypeptide, consisting of a native amino acid sequence of hIL-1Ra1S consisting of amino acid residues from at or about 26 to at or about 167, inclusive of Figure 16 (SEQ ID NO:21) fused at its N-terminus or C-terminus to a heterologous amino acid or amino acid sequence; (6) a polypeptide, such as an hIL-1Ra1S polypeptide, consisting of amino acid residues from at or about 26 to at or about 167, inclusive of Figure 16 (SEQ ID NO:21); (7) a polypeptide, such as an hIL-1Ra1S fusion variant polypeptide, consisting of a native amino acid sequence of hIL-1Ra1S consisting of amino acid residues from at or about 1 to at or about 167, inclusive of Figure 16 (SEQ ID NO:21) fused at its N-terminus or C-terminus to a heterologous amino acid or amino acid sequence; (8) a polypeptide, such as an hIL-1Ra1S polypeptide, consisting of amino acid residues from at or about 1 to at or about 167, inclusive of Figure 16 (SEQ ID NO:21);

(9) a polypeptide, such as an hIL-1Ra1S fusion variant polypeptide, consisting of a native amino acid sequence of hIL-1Ra1S consisting of amino acid residues from at or about 39 to at or about 167, inclusive of Figure 16 (SEQ ID NO:21) fused at its N-terminus or C-terminus to a heterologous amino acid or amino acid sequence; (10) a polypeptide, such as an hIL-1Ra1S fusion variant polypeptide, consisting of a native amino acid sequence of hIL-1Ra1S consisting of amino acid residues from at or about 47 to at or about 167, inclusive of Figure 16 (SEQ ID NO:21) fused at its N-terminus or C-terminus to a heterologous amino acid or amino acid sequence; (11) a polypeptide, such as an hIL-1Ra1S polypeptide, consisting of amino acid residues from at or about 39 to at or about 167, inclusive of Figure 16 (SEQ ID NO:21); (12) a polypeptide, such as an hIL-1Ra1S polypeptide, consisting of amino acid residues from at or about 47 to at or about 167, inclusive of Figure 16 (SEQ ID NO:21); (13) a polypeptide, such as an hIL-1Ra1V polypeptide, consisting of a native amino acid sequence of hIL-1Ra1V consisting of amino acid residues from at or about 1 to at or about 218, inclusive of Figure 19 (SEQ ID NO:25) fused at its N-terminus or C-terminus to a heterologous amino acid or amino acid sequence; (14) a polypeptide, such as an hIL-1Ra1V polypeptide, consisting of amino acid residues from at or about 1 to at or about 218, inclusive of Figure 19 (SEQ ID NO:25); (15) a polypeptide, such as an hIL-1Ra1V polypeptide, consisting of a native amino acid sequence of hIL-1Ra1V consisting of amino acid residues from at or about 12 to at or about 218, inclusive of Figure 19 (SEQ ID NO:25) fused at its N-terminus or C-terminus to a heterologous amino acid or amino acid sequence; (16) a polypeptide, such as an hIL-1Ra1V polypeptide, consisting of amino acid residues from at or about 12 to at or about 218, inclusive of Figure 19 (SEQ ID NO:25); (17) a polypeptide, such as an hIL-1Ra1V fusion variant polypeptide, consisting of a native amino acid sequence of hIL-1Ra1V consisting of amino acid residues from at or about 37 to at or about 218, inclusive of Figure 19 (SEQ ID NO:25) fused at its N-terminus or C-terminus to a heterologous amino acid or amino acid sequence; (18) a polypeptide, such as an hIL-1Ra1V fusion variant polypeptide, consisting of a native amino acid sequence of hIL-1Ra1V consisting of amino acid residues from at or about 46 to at or about 218, inclusive of Figure 19 (SEQ ID NO:25) fused at its N-terminus or C-terminus to a heterologous amino acid or amino acid sequence; (19) a polypeptide, such as an hIL-1Ra1V polypeptide, consisting of amino acid residues from at or about 37 to at or about 218, inclusive of Figure 19 (SEQ ID NO:25); and (20) a polypeptide, such as an hIL-1Ra1V polypeptide, consisting of amino acid residues from at or about 46 to at or about 218, inclusive of Figure 19 (SEQ ID NO:25); or (b) the complement of the DNA of (a).

In another aspect, the invention provides an isolated DNA molecule selected from the group consisting of: (1) a DNA molecule encoding a polypeptide, such as an hIL-1Ra1 polypeptide, comprising the amino acid sequence of amino acid residues from at or about 37 to at or about 203, inclusive of Figure 2 (SEQ ID NO:5); (2) a DNA molecule encoding a polypeptide, such as an hIL-1Ra1 polypeptide, comprising the amino acid sequence of amino acid residues from at or about 15 to at or about 193, inclusive of Figure 3 (SEQ ID NO:7); (3) a DNA molecule encoding a polypeptide, such as an hIL-1Ra2 polypeptide, comprising the amino acid sequence of amino acid residues from at or about 1 to at or about 134, inclusive of Figure 5 (SEQ ID

NO:10); (4) a DNA molecule encoding a polypeptide comprising the amino acid sequence of amino acid residues from at or about 10 to at or about 134, inclusive of Figure 5 (SEQ ID NO:10); (5) a DNA molecule encoding a polypeptide, such as an hIL-1Ra2 fusion variant polypeptide, consisting of a native amino acid sequence of hIL-1Ra2 consisting of amino acid residues from at or about 27 to at or about 134, inclusive of Figure 5 (SEQ ID NO:10) fused at its N-terminus or C-terminus to a heterologous amino acid or amino acid sequence; (6) a DNA molecule encoding a polypeptide, such as an hIL-1Ra2 polypeptide, consisting of the amino acid sequence of amino acid residues from at or about 27 to at or about 134, inclusive of Figure 5 (SEQ ID NO:10); (7) a DNA molecule encoding a polypeptide, such as an hIL-1Ra3 polypeptide, comprising the amino acid sequence of amino acid residues from at or about 95 to at or about 134, inclusive of Figure 7 (SEQ ID NO:13); (8) a DNA molecule encoding a polypeptide, such as a mIL-1Ra3 polypeptide, comprising the amino acid sequence of amino acid residues from about 95 to at or about 134, inclusive of Figure 9 (SEQ ID NO:16); and (9) the complement of any of the DNA molecules of (1)-(8).

In another aspect, the invention provides an isolated DNA molecule selected from the group consisting of: (1) a DNA molecule encoding a polypeptide, such as an hIL-1Ra1L polypeptide, comprising the amino acid sequence of amino acid residues from at or about 26 to at or about 207, inclusive of Figure 15 (SEQ ID NO:19); (2) a DNA molecule encoding a polypeptide, such as an hIL-1Ra1S polypeptide, comprising the amino acid sequence of amino acid residues from at or about 26 to at or about 167, inclusive of Figure 16 (SEQ ID NO:21); (3) a DNA molecule encoding a polypeptide, such as an hIL-1Ra1V polypeptide, comprising the amino acid sequence of amino acid residues from at or about 37 to at or about 218, inclusive of Figure 19 (SEQ ID NO:25); (4) a DNA molecule encoding a polypeptide, such as an hIL-1Ra1V polypeptide, comprising the amino acid sequence of amino acid residues from at or about 46 to at or about 218, inclusive of Figure 19 (SEQ ID NO:25); and (5) the complement of any of the DNA molecules of (1)-(4).

In another aspect, the invention provides an isolated DNA molecule selected from the group consisting of: (1) a DNA molecule encoding a polypeptide, such as an hIL-1Ra1L polypeptide, comprising the amino acid sequence of amino acid residues from at or about 1 to at or about 207, inclusive of Figure 15 (SEQ ID NO:19); (2) a DNA molecule encoding a polypeptide, such as an hIL-1Ra1S polypeptide, comprising the amino acid sequence of amino acid residues from at or about 1 to at or about 167, inclusive of Figure 16 (SEQ ID NO:21); (3) a DNA molecule encoding a polypeptide, such as an hIL-1Ra1V polypeptide, comprising the amino acid sequence of amino acid residues from at or about 12 to at or about 218, inclusive of Figure 19 (SEQ ID NO:25); (4) a DNA molecule encoding a polypeptide, such as an hIL-1Ra1V polypeptide, comprising the amino acid sequence of amino acid residues from at or about 1 to at or about 218, inclusive of Figure 19 (SEQ ID NO:25); and (5) the complement of any of the DNA molecules of (1)-(4).

In another aspect, the invention provides an isolated DNA molecule selected from the group consisting of: (1) a DNA molecule encoding a polypeptide, such as an hIL-1Ra3

polypeptide, comprising amino acid residues from at or about 80 to at or about 155, inclusive of Figure 7 (SEQ ID NO:13); and (2) the complement of the DNA molecule of (1).

In another aspect, the invention provides an isolated DNA molecule selected from the group consisting of: (1) a DNA molecule encoding a polypeptide, such as an hIL-1Ra1 polypeptide, comprising the amino acid sequence of amino acid residues from at or about 1 to at or about 203, inclusive of Figure 2 (SEQ ID NO:5); (2) a DNA molecule encoding a polypeptide, such as an hIL-1Ra1 polypeptide, comprising the amino acid sequence of amino acid residues from at or about 1 to at or about 193, inclusive of Figure 3 (SEQ ID NO:7); (3) a DNA molecule encoding a polypeptide, such as an hIL-1Ra3 polypeptide, comprising the amino acid sequence of amino acid residues from at or about 34 to at or about 155, inclusive of Figure 7 (SEQ ID NO:13); (4) a DNA molecule encoding a polypeptide, such as a mL-1Ra3 polypeptide, comprising the amino acid sequence of amino acid residues from at or about 34 to at or about 155, inclusive of Figure 9 (SEQ ID NO:16); and (5) the complement of any of the DNA molecules of (1)-(4).

In another aspect, the invention provides an isolated DNA molecule selected from the group consisting of: (1) a DNA molecule encoding a polypeptide, such as an hIL-1Ra3 polypeptide, comprising the amino acid sequence of amino acid residues from at or about 1 to at or about 155, inclusive of Figure 7 (SEQ ID NO:13); (2) a DNA molecule encoding a polypeptide, such as a mL-1Ra3 polypeptide, comprising the amino acid sequence of amino acid residues from at or about 1 to at or about 155, inclusive of Figure 9 (SEQ ID NO:16); and (3) the complement of any of the DNA molecules of (1)-(2).

In another aspect, the invention provides an isolated DNA molecule selected from the group consisting of: (1) a DNA molecule encoding a polypeptide, such as an hIL-1Ra3 polypeptide, comprising the amino acid sequence of amino acid residues from at or about 2 to at or about 155, inclusive of Figure 7 (SEQ ID NO:13); (2) a DNA molecule encoding a polypeptide, such as a mL-1Ra3 polypeptide, comprising the amino acid sequence of amino acid residues from at or about 2 to at or about 155, inclusive of Figure 9 (SEQ ID NO:16); and (3) the complement of any of the DNA molecules of (1)-(2).

In another aspect, the invention provides an isolated nucleic acid molecule comprising (a) a DNA molecule encoding a polypeptide selected from the group consisting of: (1) a polypeptide comprising an hIL-1Ra1 polypeptide, such as a mature hIL-1Ra1 polypeptide, encoded by the cDNA insert in the vector deposited as ATCC Deposit No. 203588; (2) a polypeptide comprising an hIL-1Ra1 polypeptide, such as a mature hIL-1Ra1 polypeptide, encoded by the cDNA insert in the vector deposited as ATCC Deposit No. 203587; (3) a polypeptide consisting of an hIL-1Ra2 polypeptide, such as a mature hIL-1Ra2 polypeptide, encoded by the cDNA insert in the vector deposited as ATCC Deposit No. 203586, which hIL-1Ra2 polypeptide is fused at its N-terminus or C-terminus to a heterologous amino acid or amino acid sequence; (4) a polypeptide consisting of an hIL-1Ra2 polypeptide, such as a mature hIL-1Ra2 polypeptide, encoded by the cDNA insert in the vector deposited as ATCC Deposit No. 203586; (5) a polypeptide comprising an hIL-1Ra3 polypeptide, such as a mature hIL-1Ra3 polypeptide, encoded by the cDNA insert in the vector deposited as ATCC Deposit No. 203589; and (6) a polypeptide comprising a mL-1Ra3 polypeptide, such as a mature mL-1Ra3

polypeptide, encoded by the cDNA insert in the vector deposited as ATCC Deposit No. 203590; or (b) the complement of the DNA molecule of (a).

In another aspect, the invention provides an isolated nucleic acid molecule comprising (a) a DNA molecule encoding a polypeptide selected from the group consisting of: (1) a polypeptide comprising an hIL-1Ra1L polypeptide, such as a mature hIL-1Ra1L polypeptide, encoded by the cDNA insert in the vector deposited as ATCC Deposit No.203846; (2) a polypeptide consisting of an hIL-1Ra1S polypeptide, such as a mature hIL-1Ra1S polypeptide, encoded by the cDNA insert in the vector deposited as ATCC Deposit No.203855, which hIL-1Ra1S polypeptide is fused at its N-terminus or C-terminus to a heterologous amino acid or amino acid sequence; (3) a polypeptide consisting of an hIL-1Ra1S polypeptide, such as a mature hIL-1Ra1S polypeptide, encoded by the cDNA insert in the vector deposited as ATCC Deposit No.203855; (4) a polypeptide comprising an hIL-1Ra1V polypeptide, such as a mature hIL-1Ra1V polypeptide, encoded by the cDNA insert in the vector deposited as ATCC Deposit No. 203973; or (b) the complement of the DNA molecule of (a).

In another aspect, the invention provides an isolated nucleic acid molecule comprising (a) a DNA molecule encoding a polypeptide comprising an hIL-1Ra1S polypeptide, such as a mature hIL-1Ra1S polypeptide, encoded by the cDNA insert in the vector deposited as ATCC Deposit No. 203855; or (b) the complement of the DNA molecule of (a).

In another aspect, the invention provides an isolated nucleic acid molecule comprising (a) a DNA encoding a polypeptide selected from the group consisting of: (1) a polypeptide comprising the entire amino acid sequence encoded by the longest open reading frame in the cDNA insert in the vector deposited as ATCC Deposit No. 203588; (2) a polypeptide comprising the entire amino acid sequence, or the entire amino acid sequence excluding the 36 N-terminal amino acid residues of such sequence, encoded by the longest open reading frame in the cDNA insert in the vector deposited as ATCC Deposit No. 203587; (3) a polypeptide comprising the entire amino acid sequence, or the entire amino acid sequence excluding the N-terminal amino acid residue of such sequence, or the entire amino acid sequence excluding the 9 N-terminal amino acid residues of such sequence, encoded by the longest open reading frame in the cDNA insert in the vector deposited as ATCC Deposit No. 203586; (4) a polypeptide comprising the entire amino acid sequence, or the entire amino acid sequence excluding the N-terminal amino acid residue of such sequence, encoded by the longest open reading frame in the cDNA insert in the vector deposited as ATCC Deposit No. 203589; and (5) a polypeptide comprising the entire amino acid sequence, or the entire amino acid sequence excluding the N-terminal amino acid residue of such sequence, encoded by the longest open reading frame in the cDNA insert in the vector deposited as ATCC Deposit No. 203590; or (b) the complement of the DNA of (a).

In another aspect, the invention provides an isolated nucleic acid molecule comprising (a) a DNA encoding a polypeptide comprising the entire amino acid sequence encoded by the longest open reading frame in the cDNA insert of a vector selected from the group consisting of the vectors deposited as ATCC Deposit Nos. 203588, 203586, 203589, 203590, and 203973, or (b) the complement of the DNA of (a).

In another aspect, the invention provides an isolated nucleic acid molecule comprising (a) a DNA encoding a polypeptide selected from the group consisting of: (1) a polypeptide comprising the entire amino acid sequence, or the entire amino acid sequence excluding the N-terminal amino acid residue of such sequence, or the entire amino acid sequence excluding the 34 N-terminal amino acid residues of such sequence, encoded by the longest open reading frame in the cDNA insert in the vector deposited as ATCC Deposit No. 203846; (2) a polypeptide comprising the entire amino acid sequence, or the entire amino acid sequence excluding the N-terminal amino acid residue of such sequence, or the entire amino acid sequence excluding the 25 N-terminal amino acid residues of such sequence, encoded by the longest open reading frame in the cDNA insert in the vector deposited as ATCC Deposit No. 204855; and (3) a polypeptide comprising the entire amino acid sequence, or the entire amino acid sequence excluding the N-terminal amino acid residue of such sequence, or the entire amino acid sequence excluding the 11 N-terminal amino acid residues of such sequence, or the entire amino acid sequence excluding the 36 N-terminal amino acid residues of such sequence, or the entire amino acid sequence excluding the 45 N-terminal amino acid residues of such sequence, encoded by the longest open reading frame in the cDNA insert in the vector deposited as ATCC Deposit No. 203973; or (b) the complement of the DNA of (a).

In another aspect, the invention provides an isolated nucleic acid molecule comprising (a) a DNA encoding a non-naturally occurring, chimeric polypeptide formed by fusing the entire amino acid sequence excluding the 38 N-terminal amino acid residues of such sequence, or the entire amino acid sequence excluding the 46 N-terminal amino acid residues of such sequence, encoded by the longest open reading frame in the cDNA insert in the vector deposited as ATCC Deposit No. 203855, at its N-terminus or C-terminus to a heterologous amino acid or amino acid sequence; or (b) the complement of the DNA of (a).

In another aspect, the invention provides an isolated nucleic acid molecule comprising (a) a DNA encoding a polypeptide consisting of the entire amino acid sequence excluding the 38 N-terminal amino acid residues of such sequence, or the entire amino acid sequence excluding the 46 N-terminal amino acid residues of such sequence, encoded by the longest open reading frame in the cDNA insert in the vector deposited as ATCC Deposit No. 203855; or (b) the complement of the DNA of (a).

In another aspect, the invention provides an isolated nucleic acid molecule comprising (a) a DNA molecule encoding a polypeptide comprising the entire amino acid sequence encoded by the longest open reading frame in the cDNA insert of a vector selected from the group consisting of the vectors deposited as ATCC Deposit Nos. 203846, 203855 and 203973, or (b) the complement of the DNA of (a).

In another aspect, the invention provides an isolated DNA molecule selected from the group consisting of: (1) a DNA molecule which encodes a polypeptide, such as an hIL-1Ra1V fusion variant polypeptide, consisting of a native amino acid sequence of hIL-1Ra1V consisting of amino acid residues from at or about 37 to at or about 218, inclusive of Figure 19 (SEQ ID NO:25) fused at its N-terminus or C-terminus to a heterologous amino acid or amino acid sequence, and which DNA molecule comprises the nucleic acid sequence in the sense strand of

Figure 19 (SEQ ID NO:24) that encodes the native amino acid sequence; (2) a DNA molecule which encodes a polypeptide, such as an hIL-1Ra1V fusion variant polypeptide, consisting of a native amino acid sequence of hIL-1Ra1V consisting of amino acid residues from at or about 46 to at or about 218, inclusive of Figure 19 (SEQ ID NO:25) fused at its N-terminus or C-terminus to a heterologous amino acid or amino acid sequence, and which DNA molecule comprises the nucleic acid sequence in the sense strand of Figure 19 (SEQ ID NO:24) that encodes the native amino acid sequence; (3) a DNA molecule which encodes a polypeptide, such as an hIL-1Ra1V polypeptide, consisting of a native amino acid sequence of hIL-1Ra1V consisting of amino acid residues from at or about 37 to at or about 218, inclusive of Figure 19 (SEQ ID NO:25), and which DNA molecule comprises the nucleic acid sequence in the sense strand of Figure 19 (SEQ ID NO:24) that encodes the native amino acid sequence; (4) a DNA molecule which encodes a polypeptide, such as an hIL-1Ra1V polypeptide, consisting of a native amino acid sequence of hIL-1Ra1V consisting of amino acid residues from at or about 46 to at or about 218, inclusive of Figure 19 (SEQ ID NO:25), and which DNA molecule comprises the nucleic acid sequence in the sense strand of Figure 19 (SEQ ID NO:24) that encodes the native amino acid sequence; and (5) the complement of any of the DNA molecules of (1)-(4).

In another aspect, the invention provides an isolated DNA molecule selected from the group consisting of: (1) a DNA molecule which encodes a polypeptide, such as an hIL-1Ra1 polypeptide, and which DNA molecule comprises the nucleic acid sequence of nucleotide positions from at or about 118 to at or about 618, inclusive in the sense strand of Figure 2 (SEQ ID NO:4); (2) a DNA molecule which encodes a polypeptide, such as an hIL-1Ra1 polypeptide, and which DNA molecule comprises the nucleic acid sequence of nucleotide positions from at or about 145 to at or about 681, inclusive in the sense strand of Figure 3 (SEQ ID NO:6); (3) a DNA molecule which encodes a polypeptide, such as an hIL-1Ra2 polypeptide, and which DNA molecule comprises the nucleic acid sequence of nucleotide positions from at or about 96 to at or about 497, inclusive in the sense strand of Figure 5 (SEQ ID NO:9); (4) a DNA molecule which comprises the nucleic acid sequence of nucleotide positions from at or about 123 to at or about 497, inclusive in the sense strand of Figure 5 (SEQ ID NO:9); (5) a DNA molecule which encodes a polypeptide, such as an hIL-1Ra2 fusion variant polypeptide, consisting of a native amino acid sequence of hIL-1Ra2 consisting of amino acid residues from at or about 27 to at or about 134, inclusive of Figure 5 (SEQ ID NO:10) fused at its N-terminus or C-terminus to a heterologous amino acid or amino acid sequence, and which DNA molecule comprises the nucleic acid sequence in the sense strand of Figure 5 (SEQ ID NO:9) that encodes the native amino acid sequence; (6) a DNA molecule which encodes a polypeptide, such as an hIL-1Ra2 polypeptide, consisting of a native amino acid sequence of hIL-1Ra2 consisting of amino acid residues from at or about 27 to at or about 134, inclusive of Figure 5 (SEQ ID NO:10), and which DNA molecule comprises the nucleic acid sequence in the sense strand of Figure 5 (SEQ ID NO:9) that encodes the native amino acid sequence; (7) a DNA molecule which encodes a polypeptide, such as an hIL-1Ra3 polypeptide, and which DNA molecule comprises the nucleic acid sequence of nucleotide positions from at or about 283 to at or about 402, inclusive in the sense strand of Figure 7 (SEQ ID NO:12); (8) a DNA molecule which encodes a polypeptide, such as a mL-1Ra3

polypeptide, and which DNA molecule comprises the nucleic acid sequence of nucleotide positions from at or about 427 to at or about 546, inclusive in the sense strand of Figure 9 (SEQ ID NO:15); and (9) the complement of any of the DNA molecules of (1)-(8).

In another aspect, the invention provides an isolated DNA molecule selected from the group consisting of: (1) a DNA molecule which encodes a polypeptide, such as an hIL-1Ra1L polypeptide, and which DNA molecule comprises the nucleic acid sequence of nucleotide positions from at or about 79 to at or about 624, inclusive in the sense strand of Figure 15 (SEQ ID NO:18); (2) a DNA molecule which encodes a polypeptide, such as an hIL-1Ra1S polypeptide, and which DNA molecule comprises the nucleic acid sequence of nucleotide positions from at or about 79 to at or about 504, inclusive in the sense strand of Figure 16 (SEQ ID NO:20); (3) a DNA molecule which encodes a polypeptide, such as an hIL-1Ra1S fusion variant polypeptide, consisting of a native amino acid sequence of hIL-1Ra1S consisting of amino acid residues from at or about 39 to at or about 167, inclusive of Figure 16 (SEQ ID NO:21) fused at its N-terminus or C-terminus to a heterologous amino acid or amino acid sequence, and which DNA molecule comprises the nucleic acid sequence in the sense strand of Figure 16 (SEQ ID NO:20) that encodes the native amino acid sequence; (4) a DNA molecule which encodes a polypeptide, such as an hIL-1Ra1S fusion variant polypeptide, consisting of a native amino acid sequence of hIL-1Ra1S consisting of amino acid residues from at or about 47 to at or about 167, inclusive of Figure 16 (SEQ ID NO:21) fused at its N-terminus or C-terminus to a heterologous amino acid or amino acid sequence, and which DNA molecule comprises the nucleic acid sequence in the sense strand of Figure 16 (SEQ ID NO:20) that encodes the native amino acid sequence; (5) a DNA molecule which encodes a polypeptide, such as an hIL-1Ra1S polypeptide, consisting of a native amino acid sequence of hIL-1Ra1S consisting of amino acid residues from at or about 39 to at or about 167, inclusive of Figure 16 (SEQ ID NO:21), and which DNA molecule comprises the nucleic acid sequence in the sense strand of Figure 16 (SEQ ID NO:20) that encodes the native amino acid sequence; (6) a DNA molecule which encodes a polypeptide, such as an hIL-1Ra1S polypeptide, consisting of a native amino acid sequence of hIL-1Ra1S consisting of amino acid residues from at or about 47 to at or about 167, inclusive of Figure 16 (SEQ ID NO:21), and which DNA molecule comprises the nucleic acid sequence in the sense strand of Figure 16 (SEQ ID NO:20) that encodes the native amino acid sequence; (7) a DNA molecule which encodes a polypeptide, such as an hIL-1Ra1V polypeptide, and which DNA molecule comprises the nucleic acid sequence of nucleotide positions from at or about 181 to at or about 729, inclusive in the sense strand of Figure 19 (SEQ ID NO:24); (8) a DNA molecule which encodes a polypeptide, such as an hIL-1Ra1V polypeptide, and which DNA molecule comprises the nucleic acid sequence of nucleotide positions from at or about 208 to at or about 729, inclusive in the sense strand of Figure 19 (SEQ ID NO:24); and (9) the complement of any of the DNA molecules of (1)-(8).

In another aspect, the invention provides an isolated DNA molecule selected from the group consisting of: (1) a DNA molecule which encodes a polypeptide, such as an hIL-1Ra1L polypeptide, and which DNA molecule comprises the nucleic acid sequence of nucleotide positions from at or about 4 to at or about 624, inclusive in the sense strand of Figure 15 (SEQ ID NO:18); (2) a DNA molecule which encodes a polypeptide, such as an hIL-1Ra1S polypeptide,

and which DNA molecule comprises the nucleic acid sequence of nucleotide positions from at or about 4 to at or about 504, inclusive in the sense strand of Figure 16 (SEQ ID NO:20); (3) a DNA molecule which encodes a polypeptide, such as an hIL-1Ra1V polypeptide, and which DNA molecule comprises the nucleic acid sequence of nucleotide positions from at or about 106 to at or about 729, inclusive in the sense strand of Figure 19 (SEQ ID NO:24); (4) a DNA molecule which encodes a polypeptide, such as an hIL-1Ra1 V polypeptide, and which comprises the nucleic acid sequence of nucleotide positions from at or about 73 to at or about 729, inclusive in the sense strand of Figure 19 (SEQ ID NO:24); and (5) the complement of any of the DNA molecules of (1)-(4).

In another aspect, the invention provides an isolated DNA molecule selected from the group consisting of: (1) a DNA molecule comprising the nucleic acid sequence of nucleotide positions from at or about 103 to at or about 681, inclusive in the sense strand of Figure 3 (SEQ ID NO:6); (2) a DNA molecule comprising the nucleic acid sequence of nucleotide positions from at or about 100 to at or about 465, inclusive in the sense strand of Figure 7 (SEQ ID NO:12); (3) a DNA molecule comprising the nucleic acid sequence of nucleotide positions from at or about 244 to at or about 609, inclusive in the sense strand of Figure 9 (SEQ ID NO:15); and (4) the complement of any of the DNA molecules of (1)-(3).

In another aspect, the invention provides an isolated DNA molecule comprising (a) the complete DNA sequence in the sense strand of Figure 2 (SEQ ID NO:4), Figure 3 (SEQ ID NO:6), Figure 5 (SEQ ID NO:9), Figure 7 (SEQ ID NO:12), or Figure 9 (SEQ ID NO:15), or (b) the complement of (a).

In another aspect, the invention provides an isolated DNA molecule comprising (a) the complete DNA sequence in the sense strand of Figure 15 (SEQ ID NO:18), Figure 16 (SEQ ID NO:20), or Figure 19 (SEQ ID NO:24), or (b) the complement of (a).

In a specific aspect, the invention provides an isolated nucleic acid molecule comprising DNA encoding an IL-11p polypeptide, with or without the N-terminal signal sequence and/or the initiating methionine, or is complementary to such IL-11p encoding nucleic acid molecule. The signal peptide has been tentatively identified as extending from amino acid position 1 to at or about amino acid position 14 in the IL-11p sequence of Figure 3 (SEQ ID NO:7), from amino acid position 1 to at or about amino acid position 26 in the IL-11p sequence of Figure 5 (SEQ ID NO:10), from amino acid position 1 to at or about amino acid position 33 in the IL-11p sequence of Figure 7 (SEQ ID NO:13), and from amino acid position 1 to at or about amino acid position 33 in the IL-11p sequence of Figure 9 (SEQ ID NO:16).

The IL-11p sequence of amino acids from at or about 1 to at or about 207 of Figure 15 (SEQ ID NO:19) is believed to behave as a mature sequence (without a presequence that is removed in post-translational processing) in certain animal cells. In addition, it is believed that other animal cells recognize and remove in post-translational processing one or more signal peptide(s) contained in the sequence of amino acid positions 1 to about 34 of Figure 15 (SEQ ID NO:19).

The IL-11p sequence of amino acids from at or about 1 to at or about 167 of Figure 16 (SEQ ID NO:21) is believed to behave as a mature sequence (without a presequence that is

removed in post-translational processing) in certain animal cells. In addition, it is believed that other animal cells recognize and remove in post-translational processing one or more signal peptide(s) contained in the sequence of amino acid positions 1 to about 46 in the IL-1lp sequence of Figure 16 (SEQ ID NO:21).

5 The IL-1lp sequence of amino acids from at or about 1 to at or about 218 of Figure 19 (SEQ ID NO:25) is believed to behave as a mature sequence (without a presequence that is removable in post-translational processing) in certain animal cells. The IL-1lp sequence of amino acids from at or about 12 to at or about 218 of Figure 19 (SEQ ID NO:25) that results from
10 initiation of translation at the start codon occurring at nucleotide positions 106-108 is also believed to behave as mature sequence in certain animal cells. It is further believed that other animal cells recognize and remove in post-translational processing one or more signal peptide(s) contained in the sequence of amino acid positions 1 to 45 in the IL-1lp polypeptide of amino acid positions 1 to 218 of Figure 19 (SEQ ID NO:25) or contained in the sequence of amino acid positions 12 to 45 in the IL-1lp polypeptide of amino acid positions 12 to 218 of Figure 19 (SEQ
15 ID NO:25).

In a still further aspect, the invention concerns an isolated nucleic acid molecule comprising (a) DNA encoding an IL-1lp polypeptide that retains at least one biologic activity of a native sequence IL-1lp, such as the IL-18R binding activity of a native sequence hIL-1Ra1, or the IL-1R binding activity of a native sequence hIL-1Ra3 or mL-1Ra3, and which IL-1lp
20 polypeptide has at least at or about 80% positives, or at least at or about 85% positives, or at least at or about 90% positives, or at least at or about 95% positives to an amino acid sequence selected from the group consisting of: (1) amino acid residues from at or about 37 to at or about 203, inclusive of Figure 2 (SEQ ID NO:5), (2) amino acid residues from at or about 15 to at or about 193, inclusive of Figure 3 (SEQ ID NO:7), (3) amino acid residues from at or about 34 to
25 at or about 155, inclusive of Figure 7 (SEQ ID NO:13), (4) amino acid residues from at or about 34 to at or about 155, inclusive of Figure 9 (SEQ ID NO:16), (5) amino acid residues from at or about 26 to at or about 207, inclusive of Figure 15 (SEQ ID NO:19), and (6) amino acid residues from at or about 46 to at or about 218, inclusive of Figure 19 (SEQ ID NO:25), or (b) the complement of the DNA of (a).

30 In a still further aspect, the invention concerns an isolated nucleic acid molecule comprising (a) DNA encoding an IL-1lp polypeptide that retains at least one biologic activity of a native sequence IL-1lp, such as the IL-18R binding activity of a native sequence hIL-1Ra1, or the IL-1R binding activity of a native sequence hIL-1Ra3 or mL-1Ra3, and which IL-1lp polypeptide has at least at or about 80% positives, or at least at or about 85% positives, or at
35 least at or about 90% positives, or at least at or about 95% positives to an amino acid sequence selected from the group consisting of: (1) amino acid residues from at or about 95 to at or about 134, inclusive of Figure 7 (SEQ ID NO:13), and (2) amino acid residues from at or about 95 to at or about 134, inclusive of Figure 9 (SEQ ID NO:16), or (b) the complement of the DNA of (a).

40 In a still further aspect, the invention concerns an isolated nucleic acid molecule comprising (a) DNA encoding an IL-1lp polypeptide that retains at least one biologic activity of a native sequence IL-1lp, such as the IL-18R binding activity of a native sequence hIL-1Ra1, or

the IL-1R binding activity of a native sequence hIL-1Ra3 or mIL-1Ra3, and which IL-1p polypeptide has at least at or about 80% positives, or at least at or about 85% positives, or at least at or about 90% positives, or at least at or about 95% positives to the amino acid sequence of amino acid residues from at or about 80 to at or about 155, inclusive of Figure 7 (SEQ ID NO:13), or (b) the complement of the DNA of (a).

In a still further aspect, the invention concerns an isolated nucleic acid molecule comprising (a) DNA encoding an IL-1lp polypeptide that retains at least one biologic activity of a native sequence IL-1lp, such as the IL-18R binding activity of a native sequence hIL-1Ra1, or the IL-1R binding activity of a native sequence hIL-1Ra3 or mIL-1Ra3, and which IL-1p polypeptide has at least at or about 80% positives, or at least at or about 85% positives, or at least at or about 90% positives, or at least at or about 95% positives to an amino acid sequence selected from the group consisting of: (1) amino acid residues from at or about 2 to at or about 155, inclusive of Figure 7 (SEQ ID NO:13), and (2) amino acid residues from at or about 2 to at or about 155, inclusive of Figure 9 (SEQ ID NO:16), or (b) the complement of the DNA of (a).

In a still further aspect, the invention concerns an isolated nucleic acid molecule comprising (a) DNA encoding an IL-1lp polypeptide having at least at or about 80% positives, or at least at or about 85% positives, or at least at or about 90% positives, or at least at or about 95% positives to an amino acid sequence selected from the group consisting of: (1) amino acid residues from at or about 95 to at or about 134, inclusive of Figure 7 (SEQ ID NO:13), and (2) amino acid residues from at or about 95 to at or about 134, inclusive of Figure 9 (SEQ ID NO:16), or (b) the complement of the DNA of (a).

In a still further aspect, the invention concerns an isolated nucleic acid molecule comprising (a) DNA encoding an IL-1lp polypeptide having at least at or about 80% positives, or at least at or about 85% positives, or at least at or about 90% positives, or at least at or about 95% positives to the amino acid sequence of amino acid residues from at or about 80 to at or about 155, inclusive of Figure 7 (SEQ ID NO:13), or (b) the complement of the DNA of (a).

In a still further aspect, the invention concerns an isolated nucleic acid molecule comprising (a) DNA encoding an IL-1lp polypeptide having at least at or about 80% positives, or at least at or about 85% positives, or at least at or about 90% positives, or at least at or about 95% positives to an amino acid sequence selected from the group consisting of: (1) amino acid residues from at or about 2 to at or about 155, inclusive of Figure 7 (SEQ ID NO:13), and (2) amino acid residues from at or about 2 to at or about 155, inclusive of Figure 9 (SEQ ID NO:16), or (b) the complement of the DNA of (a).

In a still further aspect, the invention concerns an isolated nucleic acid molecule comprising (a) DNA encoding an IL-1lp polypeptide that retains at least one biologic activity of a native sequence IL-1lp, such as the IL-18R binding activity of a native sequence hIL-1Ra1, hIL-1Ra1L, or hIL-1Ra1V, and which IL-1lp polypeptide has at least at or about 80% positives, or at least at or about 85% positives, or at least at or about 90% positives, or at least at or about 95% positives to an amino acid sequence selected from the group consisting of: (1) amino acid residues from at or about 1 to at or about 207, inclusive of Figure 15 (SEQ ID NO:19), and (2)

amino acid residues from at or about 1 to at or about 218, inclusive of Figure 19 (SEQ ID NO:25), or (b) the complement of the DNA of (a).

In another embodiment, the invention provides a vector comprising DNA encoding IL-1lp or its variants. The vector may comprise any of the isolated nucleic acid molecules hereinabove defined.

A host cell comprising such a vector is also provided. By way of example, the host cells may be CHO cells, *E. coli*, or yeast. A process for producing IL-1lp polypeptides is further provided and comprises culturing host cells under conditions suitable for expression of IL-1lp and recovering IL-1lp from the cell culture.

In another embodiment, the invention provides isolated IL-1lp polypeptide encoded by any of the isolated nucleic acid sequences hereinabove defined.

In another aspect, the invention provides isolated native sequence IL-1lp polypeptide, which in one embodiment, comprises an amino acid sequence selected from the group consisting of: (1) the amino acid sequence of residues from at or about 37 to at or about 203, inclusive of Figure 2 (SEQ ID NO:5), (2) the amino acid sequence of residues from at or about 15 to at or about 193, inclusive of Figure 3 (SEQ ID NO:7), (3) the amino acid sequence of residues from at or about 34 to at or about 155, inclusive of Figure 7 (SEQ ID NO:13), and (4) the amino acid sequence of residues from at or about 34 to at or about 155, inclusive of Figure 9 (SEQ ID NO:16).

In another aspect, the invention provides isolated native sequence IL-1lp polypeptide, which in one embodiment, comprises an amino acid sequence selected from the group consisting of: (1) the amino acid sequence of residues from at or about 1 to at or about 207, inclusive of Figure 15 (SEQ ID NO:19), (2) the amino acid sequence of residues from at or about 26 to at or about 207, inclusive of Figure 15 (SEQ ID NO:19), (3) the amino acid sequence of residues from at or about 1 to at or about 167, inclusive of Figure 16 (SEQ ID NO:21), (4) the amino acid sequence of residues from at or about 26 to at or about 167, inclusive of Figure 16 (SEQ ID NO:21), (5) the amino acid sequence of residues from at or about 1 to at or about 218, inclusive of Figure 19 (SEQ ID NO:25), (6) the amino acid sequence of residues from at or about 12 to at or about 218, inclusive of Figure 19 (SEQ ID NO:25), (7) the amino acid sequence of residues from at or about 37 to at or about 218, inclusive of Figure 19 (SEQ ID NO:25), and (8) the amino acid sequence of residues from at or about 46 to at or about 218, inclusive of Figure 19 (SEQ ID NO:25).

In another aspect, the invention concerns an isolated IL-1lp polypeptide that retains at least one biologic activity of a native sequence IL-1lp, such as the IL-1R binding activity of a native sequence hIL-1Ra3 or mIL-1Ra3, or the IL18R binding activity of a native sequence hIL-1Ra1, hIL-1Ra1L, or hIL-1Ra1V, and that consists of an amino acid sequence having at least at or about 80% sequence identity, or at least at or about 85% sequence identity, or at least at or about 90% sequence identity, or at least at or about 95% sequence identity to the sequence of amino acid residues from at or about 37 to at or about 203, inclusive of Figure 2 (SEQ ID NO:5), the sequence of amino acid residues from at or about 15 to at or about 193, inclusive of Figure 3 (SEQ ID NO:7), the sequence of amino acid residues from at or about 34 to at or about 155,

inclusive of Figure 7 (SEQ ID NO:13), the sequence of amino acid residues from at or about 34 to at or about 155, inclusive of Figure 9 (SEQ ID NO:16), the sequence of amino acid residues from at or about 26 to at or about 207, inclusive of Figure 15 (SEQ ID NO:19), or the sequence of amino acid residues from at or about 46 to at or about 218, inclusive of Figure 19 (SEQ ID NO:25).

In another aspect, the invention concerns an isolated IL-1lp polypeptide that retains at least one biologic activity of a native sequence IL-1lp, such as the IL-18R binding activity of a native sequence hIL-1Ra1, hIL-1Ra1L, or hIL-1Ra1V, and that consists of an amino acid sequence having at least at or about 80% sequence identity, or at least at or about 85% sequence identity, or at least at or about 90% sequence identity, or at least at or about 95% sequence identity to the sequence of amino acid residues from at or about 1 to at or about 207, inclusive of Figure 15 (SEQ ID NO:19), or the sequence of amino acid residues from at or about 1 to at or about 218, inclusive of Figure 19 (SEQ ID NO:25).

In another aspect, the invention provides an isolated IL-1lp selected from the group consisting of: (1) a polypeptide, such as an hIL-1Ra1 polypeptide, consisting of an amino acid sequence having a sequence identity of at least at or about 80%, or at least at or about 85%, or at least at or about 90%, or at least at or about 95%, to the sequence of amino acid residues from at or about 37 to at or about 63, inclusive of Figure 2 (SEQ ID NO:5); (2) a polypeptide, such as an hIL-1Ra1 polypeptide, consisting of an amino acid sequence having a sequence identity of at least at or about 80%, or at least at or about 85%, or at least at or about 90%, or at least at or about 95%, to the sequence of amino acid residues from at or about 15 to at or about 53, inclusive of Figure 3 (SEQ ID NO:7); (3) a polypeptide, such as an hIL-1Ra2 polypeptide, comprising the amino acid sequence of amino acid residues from at or about 1 to at or about 134, inclusive of Figure 5 (SEQ ID NO:10); (4) a polypeptide comprising the amino acid sequence of amino acid residues from at or about 10 to at or about 134, inclusive of Figure 5 (SEQ ID NO:10); (5) a polypeptide, such as an hIL-1Ra2 polypeptide, consisting of the amino acid sequence of amino acid residues from at or about 27 to at or about 134, inclusive of Figure 5 (SEQ ID NO:10); (6) a polypeptide, such as an hIL-1Ra2 fusion variant polypeptide, consisting of a native amino acid sequence of hIL-1Ra2 consisting of amino acid residues from at or about 27 to at or about 134, inclusive of Figure 5 (SEQ ID NO:10) fused at its N-terminus or C-terminus to a heterologous amino acid or amino acid sequence; (7) a polypeptide, such as an hIL-1Ra3 polypeptide, consisting of an amino acid sequence having a sequence identity of at least at or about 80%, or at least at or about 85%, or at least at or about 90%, or at least at or about 95%, to the sequence of amino acid residues from at or about 95 to at or about 134, inclusive of Figure 7 (SEQ ID NO:13); and (8) a polypeptide, such as a mL-1Ra3 polypeptide, consisting of an amino acid sequence having a sequence identity of at least at or about 80%, or at least at or about 85%, or at least at or about 90%, or at least at or about 95%, to the sequence of amino acid residues from at or about 95 to at or about 134, inclusive of Figure 9 (SEQ ID NO:16).

In another aspect, the invention provides an isolated IL-1lp selected from the group consisting of: (1) a polypeptide, such as an hIL-1Ra1L polypeptide, consisting of an amino acid sequence having a sequence identity of at least at or about 80%, or at least at or about 85%, or

at least at or about 90%, or at least at or about 95%, to the sequence of amino acid residues from at or about 27 to at or about 44, inclusive of Figure 15 (SEQ ID NO:19); (2) a polypeptide, such as an hIL-1Ra1S fusion variant polypeptide, consisting of a native amino acid sequence of hIL-1Ra1S consisting of amino acid residues from at or about 39 to at or about 167, inclusive of Figure 16 (SEQ ID NO:21) fused at its N-terminus or C-terminus to a heterologous amino acid or amino acid sequence; (3) a polypeptide, such as an hIL-1Ra1S fusion variant polypeptide, consisting of a native amino acid sequence of hIL-1Ra1S consisting of amino acid residues from at or about 47 to at or about 167, inclusive of Figure 16 (SEQ ID NO:21) fused at its N-terminus or C-terminus to a heterologous amino acid or amino acid sequence; (4) a polypeptide, such as an hIL-1Ra1S polypeptide, consisting of amino acid residues from at or about 39 to at or about 167, inclusive of Figure 16 (SEQ ID NO:21); (5) a polypeptide, such as an hIL-1Ra1S polypeptide, consisting of amino acid residues from at or about 47 to at or about 167, inclusive of Figure 16 (SEQ ID NO:21); and (6) a polypeptide, such as an hIL-1Ra1V polypeptide, consisting of an amino acid sequence having a sequence identity of at least at or about 80%, or at least at or about 85%, or at least at or about 90%, or at least at or about 95%, to the sequence of amino acid residues from at or about 37 to at or about 55, inclusive of Figure 19 (SEQ ID NO:25).

In another aspect, the invention concerns an isolated IL-1lp polypeptide that consists of an amino acid sequence having at least at or about 80% sequence identity, or at least at or about 85% sequence identity, or at least at or about 90% sequence identity, or at least at or about 95% sequence identity to the sequence of amino acid residues from at or about 37 to at or about 203, inclusive of Figure 2 (SEQ ID NO:5), the sequence of amino acid residues from at or about 15 to at or about 193, inclusive of Figure 3 (SEQ ID NO:7), the sequence of amino acid residues from at or about 34 to at or about 155, inclusive of Figure 7 (SEQ ID NO:13), the sequence of amino acid residues from at or about 34 to at or about 155, inclusive of Figure 9 (SEQ ID NO:16), the sequence of amino acid residues from at or about 26 to at or about 207, inclusive of Figure 15 (SEQ ID NO:19), or the sequence of amino acid residues from at or about 46 to at or about 218, inclusive of Figure 19 (SEQ ID NO:25).

In another aspect, the invention concerns an isolated IL-1lp polypeptide that consists of an amino acid sequence having at least at or about 80% sequence identity, or at least at or about 85% sequence identity, or at least at or about 90% sequence identity, or at least at or about 95% sequence identity to the sequence of amino acid residues from at or about 1 to at or about 207, inclusive of Figure 15 (SEQ ID NO:19), or the sequence of amino acid residues from at or about 1 to at or about 218, inclusive of Figure 19 (SEQ ID NO:25).

In another aspect, the invention provides an isolated polypeptide, such as an hIL-1Ra3 polypeptide, consisting of an amino acid sequence having a sequence identity of at least at or about 80%, or at least at or about 85%, or at least at or about 90%, or at least at or about 95%, to the sequence of amino acid residues from at or about 80 to at or about 155, inclusive of Figure 7 (SEQ ID NO:13).

In a further aspect, the invention concerns an isolated IL-1lp polypeptide that retains at least one biologic activity of a native sequence IL-1lp, such as the IL-1R binding activity of a native sequence hIL-1Ra3 or mIL-1Ra3, or the IL-18R binding activity of a native sequence hIL-

1Ra1, hIL-1Ra1L, or hIL-1Ra1V, and that consists of an amino acid sequence having at least at or about 80% sequence identity, or at least at or about 85% sequence identity, or at least at or about 90% sequence identity, or at least at or about 95% sequence identity to the amino acid sequence of an IL-1lp, such as a mature IL-1lp polypeptide, encoded by the cDNA insert in the vector deposited as ATCC Deposit No. 203588 (DNA85066-2534), ATCC Deposit No. 203587 (DNA96786-2534), ATCC Deposit No. 203589 (DNA96787-2534), ATCC Deposit No. 203590 (DNA92505-2534), ATCC Deposit No. 203846 (DNA102043-2534), or ATCC Deposit No. 203973 (DNA114876-2534). In a preferred embodiment, the IL-1lp polypeptide comprises the amino acid sequence of an IL-1lp, such as a mature IL-1lp polypeptide, encoded by the cDNA insert in the vector deposited as ATCC Deposit No. 203588 (DNA85066-2534), ATCC Deposit No. 203587 (DNA96786-2534), ATCC Deposit No. 203589 (DNA92929-2534), ATCC Deposit No. 203589 (DNA96787-2534), ATCC Deposit No. 203590 (DNA92505-2534), ATCC Deposit No. 203846 (DNA102043-2534), ATCC Deposit No. 203973 (DNA114876-2534), or ATCC Deposit No. 203855 (DNA102044-2534).

In a further aspect, the invention concerns an isolated IL-1lp polypeptide that retains at least one biologic activity of a native sequence IL-1lp, such as the IL-1R binding activity of a native sequence hIL-1Ra3 or mIL-1Ra3, or the IL-18R binding activity of a native sequence hIL-1Ra1, hIL-1Ra1L, or hIL-1Ra1V, and that consists of an amino acid sequence having at least at or about 80% sequence identity, or at least at or about 85% sequence identity, or at least at or about 90% sequence identity, or at least at or about 95% sequence identity to the entire amino acid sequence encoded by the longest open reading frame in the cDNA insert of a vector selected from the group consisting of the vectors deposited as ATCC Deposit No. 203588 (DNA85066-2534), ATCC Deposit No. 203587 (DNA96786-2534), ATCC Deposit No. 203589 (DNA96787-2534), ATCC Deposit No. 203590 (DNA92505-2534), ATCC Deposit No. 203846 (DNA102043-2534), and ATCC Deposit No. 203973 (DNA114876-2534). In a preferred embodiment, the IL-1lp polypeptide comprises the entire amino acid sequence encoded by the longest open reading frame in the cDNA insert of a vector selected from the group consisting of the vectors deposited as ATCC Deposit No. 203588 (DNA85066-2534), ATCC Deposit No. 203587 (DNA96786-2534), ATCC Deposit No. 203589 (DNA92929-2534), ATCC Deposit No. 203589 (DNA96787-2534), and ATCC Deposit No. 203590 (DNA92505-2534). In another preferred embodiment, the IL-1lp polypeptide comprises the entire amino acid sequence encoded by the longest open reading frame in the cDNA insert of a vector selected from the group consisting of the vectors deposited as ATCC Deposit No. 203846 (DNA102043-2534), ATCC Deposit No. 203855 (DNA102044-2534), and ATCC Deposit No. 203973 (DNA114876-2534).

In another aspect, the invention concerns an isolated IL-1lp polypeptide that retains at least one biologic activity of a native sequence IL-1lp, such as the IL-1R binding activity of a native sequence IL-1Ra3 or mIL-1Ra3, or the IL-18R binding activity of a native sequence hIL-1Ra1, hIL-1Ra1L, or hIL-1Ra1V, and that consists of an amino acid sequence having at least at or about 80% sequence identity, or at least at or about 85% sequence identity, or at least at or about 90% sequence identity, or at least at or about 95% sequence identity to an amino acid sequence selected from the group consisting of: (1) the entire amino acid sequence encoded by

the longest open reading frame in the cDNA insert in the vector deposited as ATCC Deposit No. 203588, (2) the entire amino acid sequence, or the entire amino acid sequence excluding the 36 N-terminal amino acid residues of such sequence, encoded by the longest open reading frame in the cDNA insert in the vector deposited as ATCC Deposit No. 203587, (3) the entire amino acid sequence, or the entire amino acid sequence excluding the N-terminal amino acid residue of such sequence, encoded by the longest open reading frame in the cDNA insert in the vector deposited as ATCC Deposit No. 203589, (4) the entire amino acid sequence, or the entire amino acid sequence excluding the N-terminal amino acid residue of such sequence, encoded by the longest open reading frame in the cDNA insert in the vector deposited as ATCC Deposit No. 203590, (5) the entire amino acid sequence, or the entire amino acid sequence excluding the N-terminal amino acid residue of such sequence, or the entire amino acid sequence excluding the 34 N-terminal amino acid residues of such sequence, encoded by the longest open reading frame in the cDNA insert in the vector deposited as ATCC Deposit No. 203846, and (6) the entire amino acid sequence, or the entire amino acid sequence excluding the N-terminal amino acid residue of such sequence, or the entire amino acid sequence excluding the 45 N-terminal amino acid residues of such sequence, encoded by the longest open reading frame in the cDNA insert in the vector deposited as ATCC Deposit No. 203973.

In a further aspect, the invention concerns an isolated IL-11p polypeptide that consists of an amino acid sequence having at least at or about 80% sequence identity, or at least at or about 85% sequence identity, or at least at or about 90% sequence identity, or at least at or about 95% sequence identity to the amino acid sequence of an IL-11p, such as a mature IL-11p polypeptide, encoded by the cDNA insert in the vector deposited as ATCC Deposit No. 203588 (DNA85066-2534), ATCC Deposit No. 203587 (DNA96786-2534), ATCC Deposit No. 203589 (DNA96787-2534), ATCC Deposit No. 203590 (DNA92505-2534), ATCC Deposit No. 203846 (DNA102043-2534), or ATCC Deposit No. 203973 (DNA114876-2534).

In a further aspect, the invention concerns an isolated IL-11p polypeptide that consists of an amino acid sequence having at least at or about 80% sequence identity, or at least at or about 85% sequence identity, or at least at or about 90% sequence identity, or at least at or about 95% sequence identity to the entire amino acid sequence encoded by the longest open reading frame in the cDNA insert of a vector selected from the group consisting of the vectors deposited as ATCC Deposit No. 203588 (DNA85066-2534), ATCC Deposit No. 203587 (DNA96786-2534), ATCC Deposit No. 203589 (DNA96787-2534), ATCC Deposit No. 203590 (DNA92505-2534), ATCC Deposit No. 203846 (DNA102043-2534), and ATCC Deposit No. 203973 (DNA114876-2534).

In another aspect, the invention concerns an isolated IL-11p polypeptide that consists of an amino acid sequence having at least at or about 80% sequence identity, or at least at or about 85% sequence identity, or at least at or about 90% sequence identity, or at least at or about 95% sequence identity to an amino acid sequence selected from the group consisting of: (1) the entire amino acid sequence encoded by the longest open reading frame in the cDNA insert in the vector deposited as ATCC Deposit No. 203588, (2) the entire amino acid sequence, or the entire amino acid sequence excluding the 36 N-terminal amino acid residues of such sequence, encoded by

the longest open reading frame in the cDNA insert in the vector deposited as ATCC Deposit No. 203587, (3) the entire amino acid sequence, or the entire amino acid sequence excluding the N-terminal amino acid residue of such sequence, encoded by the longest open reading frame in the cDNA insert in the vector deposited as ATCC Deposit No. 203589, (4) the entire amino acid sequence, or the entire amino acid sequence excluding the N-terminal amino acid residue of such sequence, encoded by the longest open reading frame in the cDNA insert in the vector deposited as ATCC Deposit No. 203590, (5) the entire amino acid sequence, or the entire amino acid sequence excluding the N-terminal amino acid residue of such sequence, or the entire amino acid sequence excluding the 34 N-terminal amino acid residues of such sequence, encoded by the longest open reading frame in the cDNA insert in the vector deposited as ATCC Deposit No. 203846, and (6) the entire amino acid sequence, or the entire amino acid sequence excluding the N-terminal amino acid residue of such sequence, or the entire amino acid sequence excluding the 45 N-terminal amino acid residues of such sequence, encoded by the longest open reading frame in the cDNA insert in the vector deposited as ATCC Deposit No. 203973.

In a preferred embodiment, the IL-11p polypeptide comprises an amino acid sequence selected from the group consisting of: (1) the entire amino acid sequence encoded by the longest open reading frame in the cDNA insert in the vector deposited as ATCC Deposit No. 203588, (2) the entire amino acid sequence, or the entire amino acid sequence excluding the 36 N-terminal amino acid residues of such sequence, encoded by the longest open reading frame in the cDNA insert in the vector deposited as ATCC Deposit No. 203587, (3) the entire amino acid sequence, or the entire amino acid sequence excluding the N-terminal amino acid residue of such sequence, encoded by the longest open reading frame in the cDNA insert in the vector deposited as ATCC Deposit No. 203589, (4) the entire amino acid sequence, or the entire amino acid sequence excluding the N-terminal amino acid residue of such sequence, encoded by the longest open reading frame in the cDNA insert in the vector deposited as ATCC Deposit No. 203590, (5) the entire amino acid sequence, or the entire amino acid sequence excluding the N-terminal amino acid residue of such sequence, or the entire amino acid sequence excluding the 34 N-terminal amino acid residues of such sequence, encoded by the longest open reading frame in the cDNA insert in the vector deposited as ATCC Deposit No. 203846, and (6) the entire amino acid sequence, or the entire amino acid sequence excluding the N-terminal amino acid residue of such sequence, or the entire amino acid sequence excluding the 45 N-terminal amino acid residues of such sequence, encoded by the longest open reading frame in the cDNA insert in the vector deposited as ATCC Deposit No. 203973.

In another aspect, the invention concerns an isolated IL-11p polypeptide that retains at least one biologic activity of a native sequence IL-11p, such as the IL-1R binding activity of a native sequence hIL-1Ra3 or mL-1Ra3, or the IL-18R binding activity of a native sequence hIL-1Ra1, hIL-1Ra1L, or hIL-1Ra1V, and that consists of an amino acid sequence having at least at or about 80% positives, or at least at or about 85% positives, or at least at or about 90% positives, or at least at or about 95% positives to the sequence of amino acid residues from at or about 37 to at or about 203, inclusive of Figure 2 (SEQ ID NO:5), the sequence of amino acid residues from at or about 15 to at or about 193, inclusive of Figure 3 (SEQ ID NO:7), the

sequence of amino acid residues from at or about 34 to at or about 155, inclusive of Figure 7 (SEQ ID NO:13), the sequence of amino acid residues from at or about 34 to at or about 155, inclusive of Figure 9 (SEQ ID NO:16), or the sequence of amino acid residues from at or about 46 to at or about 218, inclusive of Figure 19 (SEQ ID NO:25).

In another aspect, the invention concerns an isolated IL-1lp polypeptide that retains at least one biologic activity of a native sequence IL-1lp, such as the IL-18R binding activity of a native sequence hIL-1Ra1, hIL-1Ra1L, or hIL-1Ra1V, and that consists of an amino acid sequence having at least at or about 80% positives, or at least at or about 85% positives, or at least at or about 90% positives, or at least at or about 95% positives to the sequence of amino acid residues from at or about 1 to at or about 207, inclusive of Figure 15 (SEQ ID NO:19), or the sequence of amino acid residues from at or about 1 to at or about 218, inclusive of Figure 19 (SEQ ID NO:25).

In another aspect, the invention provides an isolated IL-1lp selected from the group consisting of: (1) a polypeptide, such as an hIL-1Ra1 polypeptide, consisting of an amino acid sequence having a % positives value of at least at or about 80%, or at least at or about 85%, or at least at or about 90%, or at least at or about 95%, to the sequence of amino acid residues from at or about 37 to at or about 63, inclusive of Figure 2 (SEQ ID NO:5); (2) a polypeptide, such as an hIL-1Ra1 polypeptide, consisting of an amino acid sequence having a % positives value of at least at or about 80%, or at least at or about 85%, or at least at or about 90%, or at least at or about 95%, to the sequence of amino acid residues from at or about 15 to at or about 53, inclusive of Figure 3 (SEQ ID NO:7); (3) a polypeptide, such as an hIL-1Ra2 polypeptide, comprising the amino acid sequence of amino acid residues from at or about 1 to at or about 134, inclusive of Figure 5 (SEQ ID NO:10); (4) a polypeptide comprising the amino acid sequence of amino acid residues from at or about 10 to at or about 134, inclusive of Figure 5 (SEQ ID NO:10); (5) a polypeptide, such as an hIL-1Ra2 polypeptide, consisting of the amino acid sequence of amino acid residues from at or about 27 to at or about 134, inclusive of Figure 5 (SEQ ID NO:10); (6) a polypeptide, such as an hIL-1Ra2 fusion variant polypeptide, consisting of a native amino acid sequence of hIL-1Ra2 consisting of amino acid residues from at or about 27 to at or about 134, inclusive of Figure 5 (SEQ ID NO:10) fused at its N-terminus or C-terminus to a heterologous amino acid or amino acid sequence; (7) a polypeptide, such as an hIL-1Ra3 polypeptide, consisting of an amino acid sequence having a % positives value of at least at or about 80%, or at least at or about 85%, or at least at or about 90%, or at least at or about 95%, to the sequence of amino acid residues from at or about 95 to at or about 134, inclusive of Figure 7 (SEQ ID NO:13); and (8) a polypeptide, such as a mL-1Ra3 polypeptide, consisting of an amino acid sequence having a % positives value of at least at or about 80%, or at least at or about 85%, or at least at or about 90%, or at least at or about 95%, to the sequence of amino acid residues from at or about 95 to at or about 134, inclusive of Figure 9 (SEQ ID NO:16).

In another aspect, the invention provides an isolated IL-1lp selected from the group consisting of: (1) a polypeptide, such as an hIL-1Ra1L polypeptide, consisting of an amino acid sequence having a % positives value of at least at or about 80%, or at least at or about 85%, or at least at or about 90%, or at least at or about 95%, to the sequence of amino acid residues

from at or about 27 to at or about 44, inclusive of Figure 15 (SEQ ID NO:19); (2) a polypeptide, such as an hIL-1Ra1S fusion variant polypeptide, consisting of a native amino acid sequence of hIL-1Ra1S consisting of amino acid residues from at or about 39 to at or about 167, inclusive of Figure 16 (SEQ ID NO:21) fused at its N-terminus or C-terminus to a heterologous amino acid or amino acid sequence; (3) a polypeptide, such as an hIL-1Ra1S fusion variant polypeptide, consisting of a native amino acid sequence of hIL-1Ra1S consisting of amino acid residues from at or about 47 to at or about 167, inclusive of Figure 16 (SEQ ID NO:21) fused at its N-terminus or C-terminus to a heterologous amino acid or amino acid sequence; (4) a polypeptide, such as an hIL-1Ra1S polypeptide, consisting of amino acid residues from at or about 39 to at or about 167, inclusive of Figure 16 (SEQ ID NO:21); (5) a polypeptide, such as an hIL-1Ra1S polypeptide, consisting of amino acid residues from at or about 47 to at or about 167, inclusive of Figure 16 (SEQ ID NO:21); and (6) a polypeptide, such as an hIL-1Ra1V polypeptide, consisting of an amino acid sequence having a % positives value of at least at or about 80%, or at least at or about 85%, or at least at or about 90%, or at least at or about 95%, to the sequence of amino acid residues from at or about 37 to at or about 55, inclusive of Figure 19 (SEQ ID NO:25).

In another aspect, the invention concerns an isolated IL-1lp polypeptide that consists of an amino acid sequence having at least at or about 80% positives, or at least at or about 85% positives, or at least at or about 90% positives, or at least at or about 95% positives to the sequence of amino acid residues from at or about 37 to at or about 203, inclusive of Figure 2 (SEQ ID NO:5), the sequence of amino acid residues from at or about 15 to at or about 193, inclusive of Figure 3 (SEQ ID NO:7), the sequence of amino acid residues from at or about 34 to at or about 155, inclusive of Figure 7 (SEQ ID NO:13), the sequence of amino acid residues from at or about 34 to at or about 155, inclusive of Figure 9 (SEQ ID NO:16), or the sequence of amino acid residues from at or about 46 to at or about 218, inclusive of Figure 19 (SEQ ID NO:25).

In another aspect, the invention concerns an isolated IL-1lp polypeptide that consists of an amino acid sequence having at least at or about 80% positives, or at least at or about 85% positives, or at least at or about 90% positives, or at least at or about 95% positives to the sequence of amino acid residues from at or about 1 to at or about 207, inclusive of Figure 15 (SEQ ID NO:19), or the sequence of amino acid residues from at or about 1 to at or about 218, inclusive of Figure 19 (SEQ ID NO:25).

In a further aspect, the invention concerns an isolated IL-1lp polypeptide that retains at least one biologic activity of a native sequence IL-1lp, such as the IL-1R binding activity of a native sequence hIL-1Ra3 or mL-1Ra3, or the IL-18R binding activity of a native sequence hIL-1Ra1, hIL-1Ra1L, or hIL-1Ra1V, and that consists of an amino acid sequence that has at least at or about 80% positives, or at least at or about 85% positives, or at least at or about 90% positives, or at least at or about 95% positives to the amino acid sequence of an IL-1lp, such as a mature IL-1lp polypeptide, encoded by the cDNA insert in the vector deposited as ATCC Deposit No. 203588 (DNA85066-2534), ATCC Deposit No. 203587 (DNA96786-2534), ATCC Deposit No. 203589 (DNA96787-2534), ATCC Deposit No. 203590 (DNA92505-2534), ATCC Deposit No. 203846 (DNA102043-2534), or ATCC Deposit No. 203973 (DNA114876-2534).

In a further aspect, the invention concerns an isolated IL-11p polypeptide that retains at least one biologic activity of a native sequence IL-11p, such as the IL-1R binding activity of a native sequence hIL-1Ra3 or mL-1Ra3, or the IL-18R binding activity of a native sequence hIL-1Ra1, hIL-1Ra1L, or hIL-1Ra1V, and that consists of an amino acid sequence that has at least 5 at or about 80% positives, or at least at or about 85% positives, or at least at or about 90% positives, or at least at or about 95% positives to the entire amino acid sequence encoded by the longest open reading frame in the cDNA insert of a vector selected from the group consisting of the vectors deposited as ATCC Deposit No. 203588 (DNA85066-2534), ATCC Deposit No. 203587 (DNA96786-2534), ATCC Deposit No. 203589 (DNA96787-2534), ATCC Deposit No. 203590 (DNA92505-2534), ATCC Deposit No. 203846 (DNA102043-2534), and ATCC Deposit No. 203973 (DNA114876-2534).

In another aspect, the invention concerns an isolated IL-11p polypeptide that retains at least one biologic activity of a native sequence IL-11p, such as the IL-1R binding activity of a native sequence IL-1Ra3 or mL-1Ra3, or the IL-18R binding activity of a native sequence hIL-1Ra1, hIL-1Ra1L, or hIL-1Ra1V, and that consists of an amino acid sequence having at least at or about 80% positives, or at least at or about 85% positives, or at least at or about 90% positives, or at least at or about 95% positives to an amino acid sequence selected from the group consisting of: (1) the entire amino acid sequence encoded by the longest open reading frame in the cDNA insert in the vector deposited as ATCC Deposit No. 203588, (2) the entire amino acid sequence, or the entire amino acid sequence excluding the 36 N-terminal amino acid residues of such sequence, encoded by the longest open reading frame in the cDNA insert in the vector deposited as ATCC Deposit No. 203587, (3) the entire amino acid sequence, or the entire amino acid sequence excluding the N-terminal amino acid residue of such sequence, encoded by the longest open reading frame in the cDNA insert in the vector deposited as ATCC Deposit No. 203589, (4) the entire amino acid sequence, or the entire amino acid sequence excluding the N-terminal amino acid residue of such sequence, encoded by the longest open reading frame in the cDNA insert in the vector deposited as ATCC Deposit No. 203590, (5) the entire amino acid sequence, or the entire amino acid sequence excluding the N-terminal amino acid residue of such sequence, or the entire amino acid sequence excluding the 34 N-terminal amino acid residues of such sequence, encoded by the longest open reading frame in the cDNA insert in the vector deposited as ATCC Deposit No. 203846, and (6) the entire amino acid sequence, or the entire amino acid sequence excluding the N-terminal amino acid residue of such sequence, or the entire amino acid sequence excluding the 45 N-terminal amino acid residues of such sequence, encoded by the longest open reading frame in the cDNA insert in the vector deposited as ATCC Deposit No. 203973.

In a further aspect, the invention concerns an isolated IL-11p polypeptide that consists of an amino acid sequence that has at least at or about 80% positives, or at least at or about 85% positives, or at least at or about 90% positives, or at least at or about 95% positives to the amino acid sequence of an IL-11p, such as a mature IL-11p polypeptide, encoded by the cDNA insert in the vector deposited as ATCC Deposit No. 203588 (DNA85066-2534), ATCC Deposit No. 203587 (DNA96786-2534), ATCC Deposit No. 203589 (DNA96787-2534), ATCC Deposit No. 203590 (DNA92505-2534), ATCC Deposit No. 203846 (DNA102043-2534), and ATCC Deposit No. 203973 (DNA114876-2534).

203590 (DNA92505-2534), ATCC Deposit No. 203846 (DNA102043-2534), or ATCC Deposit No. 203973 (DNA114876-2534).

In a further aspect, the invention concerns an isolated IL-11p polypeptide that consists of an amino acid sequence that has at least at or about 80% positives, or at least at or about 85% positives, or at least at or about 90% positives, or at least at or about 95% positives to the entire amino acid sequence encoded by the longest open reading frame in the cDNA insert of a vector selected from the group consisting of the vectors deposited as ATCC Deposit No. 203588 (DNA85066-2534), ATCC Deposit No. 203587 (DNA96786-2534), ATCC Deposit No. 203589 (DNA96787-2534), ATCC Deposit No. 203590 (DNA92505-2534), ATCC Deposit No. 203846 (DNA102043-2534), and ATCC Deposit No. 203973 (DNA114876-2534).

In another aspect, the invention concerns an isolated IL-11p polypeptide that consists of an amino acid sequence having at least at or about 80% positives, or at least at or about 85% positives, or at least at or about 90% positives, or at least at or about 95% positives to an amino acid sequence selected from the group consisting of: (1) the entire amino acid sequence encoded by the longest open reading frame in the cDNA insert in the vector deposited as ATCC Deposit No. 203588, (2) the entire amino acid sequence, or the entire amino acid sequence excluding the 36 N-terminal amino acid residues of such sequence, encoded by the longest open reading frame in the cDNA insert in the vector deposited as ATCC Deposit No. 203587, (3) the entire amino acid sequence, or the entire amino acid sequence excluding the N-terminal amino acid residue of such sequence, encoded by the longest open reading frame in the cDNA insert in the vector deposited as ATCC Deposit No. 203589, (4) the entire amino acid sequence, or the entire amino acid sequence excluding the N-terminal amino acid residue of such sequence, encoded by the longest open reading frame in the cDNA insert in the vector deposited as ATCC Deposit No. 203590, (5) the entire amino acid sequence, or the entire amino acid sequence excluding the N-terminal amino acid residue of such sequence, or the entire amino acid sequence excluding the 34 N-terminal amino acid residues of such sequence, encoded by the longest open reading frame in the cDNA insert in the vector deposited as ATCC Deposit No. 203846, and (6) the entire amino acid sequence, or the entire amino acid sequence excluding the N-terminal amino acid residue of such sequence, or the entire amino acid sequence excluding the 45 N-terminal amino acid residues of such sequence, encoded by the longest open reading frame in the cDNA insert in the vector deposited as ATCC Deposit No. 203973.

In another aspect, the invention provides an isolated polypeptide, such as an hIL-1Ra3 polypeptide, consisting of an amino acid sequence having a % positives value of at least at or about 80%, or at least at or about 85%, or at least at or about 90%, or at least at or about 95%, to the sequence of amino acid residues from at or about 80 to at or about 155, inclusive of Figure 7 (SEQ ID NO:13).

In yet another aspect, the invention concerns an isolated IL-11p polypeptide comprising an amino acid sequence selected from the group consisting of: (1) amino acid residues from at or about 37 to at or about 63, inclusive of Figure 2 (SEQ ID NO:5); (2) amino acid residues from at or about 15 to at or about 53, inclusive of Figure 3 (SEQ ID NO:7); (3) amino acid residues from at or about 80 to at or about 155, inclusive of Figure 7 (SEQ ID NO:13); and (4) amino acid

residues from at or about 95 to at or about 155, inclusive of Figure 9 (SEQ ID NO:16), or a fragment of such IL-11p polypeptide that coincides with a stretch of at least about 10 contiguous amino acids in such amino acid sequence, wherein the IL-11p polypeptide or fragment thereof is sufficient to provide a binding site for an anti-IL-11p antibody. Preferably, the IL-11p fragment retains at least one biologic activity of a native sequence IL-11p polypeptide.

In yet another aspect, the invention concerns an isolated IL-11p polypeptide comprising an amino acid sequence selected from the group consisting of: (1) amino acid residues from at or about 26 to at or about 44, inclusive of Figure 15 (SEQ ID NO:19); (2) amino acid residues from at or about 26 to at or about 78, inclusive of Figure 19 (SEQ ID NO:25); and (3) amino acid residues from at or about 46 to at or about 89, inclusive of Figure 19 (SEQ ID NO:25), or a fragment of such IL-11p polypeptide that coincides with a stretch of at least about 10 contiguous amino acids in such amino acid sequence, wherein the IL-11p polypeptide or fragment thereof is sufficient to provide a binding site for an anti-IL-11p antibody. Preferably, the IL-11p fragment retains at least one biologic activity of a native sequence IL-11p polypeptide.

In a further aspect, the invention provides an isolated IL-11p polypeptide selected from the group consisting of: (1) a polypeptide, such as an hIL-1Ra1 polypeptide, comprising amino acid residues from at or about 37 to at or about 63, inclusive of Figure 2 (SEQ ID NO:5); (2) a polypeptide, such as an hIL-1Ra1 polypeptide, comprising amino acid residues from at or about 15 to at or about 53, inclusive of Figure 3 (SEQ ID NO:7); (3) a polypeptide, such as an hIL-1Ra3 polypeptide, comprising amino acid residues from at or about 95 to at or about 134, inclusive of Figure 7 (SEQ ID NO:13); and (4) a polypeptide, such as a mL-1Ra3 polypeptide, comprising amino acid residues from at or about 95 to at or about 134, inclusive of Figure 9 (SEQ ID NO:16).

In a further aspect, the invention provides an isolated IL-11p polypeptide selected from the group consisting of: (1) a polypeptide, such as an hIL-1Ra1 polypeptide, comprising amino acid residues from at or about 37 to at or about 203, inclusive of Figure 2 (SEQ ID NO:5); (2) a polypeptide, such as an hIL-1Ra1 polypeptide, comprising amino acid residues from at or about 15 to at or about 193, inclusive of Figure 3 (SEQ ID NO:7); (3) a polypeptide, such as an hIL-1Ra3 polypeptide, comprising amino acid residues from at or about 34 to at or about 155, inclusive of Figure 7 (SEQ ID NO:13); and (3) a polypeptide, such as a mL-1Ra3 polypeptide, comprising amino acid residues from at or about 34 to at or about 155, inclusive of Figure 9 (SEQ ID NO:16).

In a further aspect, the invention provides an isolated polypeptide, such as an hIL-1Ra3 polypeptide, comprising amino acid residues from at or about 80 to at or about 155, inclusive of Figure 7 (SEQ ID NO:13).

In a further aspect, the invention provides an isolated IL-11p polypeptide selected from the group consisting of: (1) a polypeptide, such as an hIL-1Ra1L polypeptide, comprising amino acid residues from at or about 26 to at or about 44, inclusive of Figure 15 (SEQ ID NO:19); (2) a polypeptide, such as an hIL-1Ra1L polypeptide, comprising amino acid residues from at or about 1 to at or about 44, inclusive of Figure 15 (SEQ ID NO:19); (3) a polypeptide, such as an hIL-1Ra1S polypeptide, comprising amino acid residues from at or about 1 to at or about 38, inclusive of Figure 16 (SEQ ID NO:21); (4) a polypeptide, such as an hIL-1Ra1V polypeptide,

comprising amino acid residues from at or about 37 to at or about 55, inclusive of Figure 19 (SEQ ID NO:25); (5) a polypeptide, such as an hIL-1Ra1V polypeptide, comprising amino acid residues from at or about 12 to at or about 55, inclusive of Figure 19 (SEQ ID NO:25); and (6) a polypeptide, such as an hIL-1Ra1V polypeptide, comprising amino acid residues from at or about 1 to at or about 55, inclusive of Figure 19 (SEQ ID NO:25).

In a further aspect, the invention provides an isolated IL-11p polypeptide selected from the group consisting of: (1) a polypeptide, such as an hIL-1Ra1L fusion variant polypeptide, consisting of a native amino acid sequence of hIL-1Ra1L consisting of amino acid residues from at or about 1 to at or about 207, inclusive of Figure 15 (SEQ ID NO:19) fused at its N-terminus or C-terminus to a heterologous amino acid or amino acid sequence; (2) a polypeptide, such as an hIL-1Ra1L fusion variant polypeptide, consisting of a native amino acid sequence of hIL-1Ra1L consisting of amino acid residues from at or about 26 to at or about 207, inclusive of Figure 15 (SEQ ID NO:19) fused at its N-terminus or C-terminus to a heterologous amino acid or amino acid sequence; (3) a polypeptide, such as an hIL-1Ra1L polypeptide, consisting of amino acid residues from at or about 1 to at or about 207, inclusive of Figure 15 (SEQ ID NO:19); (4) a polypeptide, such as an hIL-1Ra1L polypeptide, consisting of amino acid residues from at or about 26 to at or about 207, inclusive of Figure 15 (SEQ ID NO:19); (5) a polypeptide, such as an hIL-1Ra1V fusion variant polypeptide, consisting of a native amino acid sequence of hIL-1Ra1V consisting of amino acid residues from at or about 1 to at or about 218, inclusive of Figure 19 (SEQ ID NO:25) fused at its N-terminus or C-terminus to a heterologous amino acid or amino acid sequence; (6) a polypeptide, such as an hIL-1Ra1V fusion variant polypeptide, consisting of a native amino acid sequence of hIL-1Ra1V consisting of amino acid residues from at or about 12 to at or about 218, inclusive of Figure 19 (SEQ ID NO:25) fused at its N-terminus or C-terminus to a heterologous amino acid or amino acid sequence; (7) a polypeptide, such as an hIL-1Ra1V fusion variant polypeptide, consisting of a native amino acid sequence of hIL-1Ra1V consisting of amino acid residues from at or about 37 to at or about 218, inclusive of Figure 19 (SEQ ID NO:25) fused at its N-terminus or C-terminus to a heterologous amino acid or amino acid sequence; (8) a polypeptide, such as an hIL-1Ra1V fusion variant polypeptide, consisting of a native amino acid sequence of hIL-1Ra1V consisting of amino acid residues from at or about 46 to at or about 218, inclusive of Figure 19 (SEQ ID NO:25) fused at its N-terminus or C-terminus to a heterologous amino acid or amino acid sequence; (9) a polypeptide, such as an hIL-1Ra1V polypeptide, consisting of the amino acid sequence of amino acid residues from at or about 1 to at or about 218, inclusive of Figure 19 (SEQ ID NO:25); (10) a polypeptide, such as an hIL-1Ra1V polypeptide, consisting of the amino acid sequence of amino acid residues from at or about 12 to at or about 218, inclusive of Figure 19 (SEQ ID NO:25); (11) a polypeptide, such as an hIL-1Ra1V polypeptide, consisting of the amino acid sequence of amino acid residues from at or about 37 to at or about 218, inclusive of Figure 19 (SEQ ID NO:25); and (12) a polypeptide, such as an hIL-1Ra1V polypeptide, consisting of the amino acid sequence of amino acid residues from at or about 46 to at or about 218, inclusive of Figure 19 (SEQ ID NO:25).

In a further aspect, the invention provides an isolated IL-11p polypeptide selected from the group consisting of: (1) a polypeptide, such as an hIL-1Ra1L polypeptide, comprising amino

acid residues from at or about 1 to at or about 207, inclusive of Figure 15 (SEQ ID NO:19); (2) a polypeptide, such as an hIL-1Ra1L polypeptide, comprising amino acid residues from at or about 26 to at or about 207, inclusive of Figure 15 (SEQ ID NO:19); (3) a polypeptide, such as an hIL-1Ra1S polypeptide, comprising amino acid residues from at or about 1 to at or about 167, inclusive of Figure 16 (SEQ ID NO:21); (4) a polypeptide, such as an hIL-1Ra1S polypeptide, comprising amino acid residues from at or about 26 to at or about 167, inclusive of Figure 16 (SEQ ID NO:21); (5) a polypeptide, such as an hIL-1Ra1V polypeptide, comprising amino acid residues from at or about 1 to at or about 218, inclusive of Figure 19 (SEQ ID NO:25); and (6) a polypeptide, such as an hIL-1Ra1V polypeptide, comprising amino acid residues from at or about 12 to at or about 218, inclusive of Figure 19 (SEQ ID NO:25).

In a further aspect, the invention provides an isolated IL-1lp polypeptide selected from the group consisting of: (1) a polypeptide, such as an hIL-1Ra1V polypeptide, comprising amino acid residues from at or about 37 to at or about 218, inclusive of Figure 19 (SEQ ID NO:25); and (2) a polypeptide, such as an hIL-1Ra1V polypeptide, comprising amino acid residues from at or about 46 to at or about 218, inclusive of Figure 19 (SEQ ID NO:25).

In a further aspect, the invention provides an isolated IL-1lp polypeptide selected from the group consisting of: (1) a polypeptide consisting of the amino acid sequence of amino acid residues 10 to 134, inclusive of Figure 5 (SEQ ID NO:10) fused at its N-terminus or C-terminus to a heterologous amino acid or amino acid sequence; (2) a polypeptide consisting of the amino acid sequence of amino acid residues 10 to 134, inclusive of Figure 5 (SEQ ID NO:10); (3) a polypeptide, such as an hIL-1Ra3 polypeptide, consisting of a native amino acid sequence of hIL-1Ra3 consisting of amino acid residues 2 to 155, inclusive of Figure 7 (SEQ ID NO:13) fused at its N-terminus or C-terminus to a heterologous amino acid or amino acid sequence; (4) a polypeptide, such as an hIL-1Ra3 polypeptide, consisting of the amino acid sequence of amino acid residues from 2 to 155, inclusive of Figure 7 (SEQ ID NO:13); (5) a polypeptide, such as a mL-1Ra3 polypeptide, consisting of a native amino acid sequence of mL-1Ra3 consisting of amino acid residues 2 to 155, inclusive of Figure 9 (SEQ ID NO:16) fused at its N-terminus or C-terminus to a heterologous amino acid or amino acid sequence; (6) a polypeptide, such as a mL-1Ra3 polypeptide, consisting of the amino acid sequence of amino acid residues 2 to 155, inclusive of Figure 9 (SEQ ID NO:16).

In a further aspect, the invention provides an isolated IL-1lp polypeptide selected from the group consisting of: (1) a polypeptide comprising the amino acid sequence of amino acid residues 10 to 134, inclusive of Figure 5 (SEQ ID NO:10); (2) a polypeptide, such as an hIL-1Ra3 polypeptide, comprising the amino acid sequence of amino acid residues 2 to 155, inclusive of Figure 7 (SEQ ID NO:13); and (3) a polypeptide, such as a mL-1Ra3 polypeptide, comprising the amino acid sequence of amino acid residues from 2 to 155, inclusive of Figure 9 (SEQ ID NO:16).

In a still further aspect, the invention provides an isolated IL-1lp polypeptide that is the same as a mature polypeptide encoded by the cDNA insert of a vector selected from the group consisting of the vectors deposited as ATCC Deposit Nos. 203588, 203587, 203586, 203589, and 203590.

In a still further aspect, the invention provides an isolated IL-11p polypeptide that is the same as a mature polypeptide encoded by the cDNA insert of a vector selected from the group consisting of the vectors deposited as ATCC Deposit Nos. 203846, 203855 and 203973.

In a still further aspect, the invention provides an isolated polypeptide comprising the entire amino acid sequence encoded by the longest open reading frame in the cDNA insert of a vector selected from the group consisting of the vectors deposited as ATCC Deposit Nos. 203586, 203589 and 203590.

In a still further aspect, the invention provides an isolated polypeptide comprising the entire amino acid sequence encoded by the longest open reading frame in the cDNA insert of a vector selected from the group consisting of the vectors deposited as ATCC Deposit Nos. 203846, 203855 and 203973.

In another aspect, the invention provides a polypeptide that is produced by (i) hybridizing a test DNA molecule under stringent conditions with (a) a DNA molecule encoding an amino acid sequence selected from the group consisting of: (1) amino acid residues from at or about 37 to at or about 203, inclusive of Figure 2 (SEQ ID NO:5), (2) amino acid residues from at or about 15 to at or about 193, inclusive of Figure 3 (SEQ ID NO:7), (3) amino acid residues from at or about 2 to at or about 155, inclusive of Figure 7 (SEQ ID NO:13), and (4) amino acid residues from at or about 2 to at or about 155, inclusive of Figure 9 (SEQ ID NO:16), or (b) the complement of the DNA molecule of (a), and if the test DNA molecule encodes an IL-11p polypeptide that retains at least one biologic activity of a native sequence IL-11p, such as the IL-1R binding activity of a native sequence hIL-1Ra3 or mL-1Ra3, or the IL-18R binding activity of a native sequence hIL-1Ra1, hIL-1Ra1L, or hIL-1Ra1V, and if the test DNA molecule has at least at or about an 80% sequence identity, or at least at or about an 85% sequence identity, or at least at or about a 90% sequence identity, or at least at or about a 95% sequence identity to the DNA molecule of (a) or (b), (ii) culturing a host cell comprising the test DNA molecule under conditions suitable for expression of the IL-11p polypeptide, and (iii) recovering the IL-11p polypeptide from the cell culture.

In a still further aspect, the invention provides a polypeptide that is produced by (i) hybridizing a test DNA molecule under stringent conditions with (a) a DNA molecule encoding an amino acid sequence selected from the group consisting of: (1) amino acid residues from at or about 26 to at or about 207, inclusive of Figure 15 (SEQ ID NO:19), (2) amino acid residues from at or about 1 to at or about 218, inclusive of Figure 19 (SEQ ID NO:25), (3) amino acid residues from at or about 12 to at or about 218, inclusive of Figure 19 (SEQ ID NO:25), (4) amino acid residues from at or about 37 to at or about 218, inclusive of Figure 19 (SEQ ID NO:25), and (5) amino acid residues from at or about 46 to at or about 218, inclusive of Figure 19 (SEQ ID NO:25), or (b) the complement of the DNA molecule of (a), and if the test DNA molecule encodes an IL-11p polypeptide that retains at least one biologic activity of a native sequence IL-11p, such as the IL-18R binding activity of a native sequence hIL-1Ra1, hIL-1Ra1L, or hIL-1Ra1V, and if the test DNA molecule has at least at or about an 80% sequence identity, or at least at or about an 85% sequence identity, or at least at or about a 90% sequence identity, or at least at or about a 95% sequence identity to the DNA molecule of (a) or (b), (ii) culturing a host cell comprising

the test DNA molecule under conditions suitable for expression of the IL-11p polypeptide, and (iii) recovering the IL-11p polypeptide from the cell culture.

A. Preparation of IL-11p

The description below relates primarily to production of IL-11p by culturing cells transformed or transfected with a vector containing IL-11p nucleic acid. It is, of course, contemplated that alternative methods, which are well known in the art, may be employed to prepare IL-11p. For instance, the IL-11p sequence, or portions thereof, may be produced by direct peptide synthesis using solid-phase techniques [see, e.g., Stewart *et al.*, *Solid-Phase Peptide Synthesis*, W.H. Freeman Co., San Francisco, CA (1969); Merrifield, *J. Am. Chem. Soc.*, 85:2149-2154 (1963)]. *In vitro* protein synthesis may be performed using manual techniques or by automation. Automated synthesis may be accomplished, for instance, using an Applied Biosystems Peptide Synthesizer (Foster City, CA) using manufacturer's instructions. Various portions of the IL-11p may be chemically synthesized separately and combined using chemical or enzymatic methods to produce the full-length IL-11p.

1. Isolation of DNA Encoding IL-11p

DNA encoding IL-11p may be obtained from a cDNA library prepared from tissue believed to possess the IL-11p mRNA and to express it at a detectable level. Accordingly, human IL-11p DNA can be conveniently obtained from a cDNA library prepared from human tissue, such as described in the Examples. The IL-11p-encoding gene may also be obtained from a genomic library or by oligonucleotide synthesis.

Libraries can be screened with probes (such as antibodies to the IL-11p or oligonucleotides of at least about 20-80 bases) designed to identify the gene of interest or the protein encoded by it. Screening the cDNA or genomic library with the selected probe may be conducted using standard procedures, such as described in Sambrook *et al.*, *Molecular Cloning: A Laboratory Manual* (New York: Cold Spring Harbor Laboratory Press, 1989). An alternative means to isolate the gene encoding IL-11p is to use PCR methodology [Sambrook *et al.*, *supra*; Dieffenbach *et al.*, *PCR Primer: A Laboratory Manual* (Cold Spring Harbor Laboratory Press, 1995)].

The Examples below describe techniques for screening a cDNA library. The oligonucleotide sequences selected as probes should be of sufficient length and sufficiently unambiguous that false positives are minimized. The oligonucleotide is preferably labeled such that it can be detected upon hybridization to DNA in the library being screened. Methods of labeling are well known in the art, and include the use of radiolabels like ³²P-labeled ATP, biotinylation or enzyme labeling. Hybridization conditions, including moderate stringency and high stringency, are provided in Sambrook *et al.*, *supra*.

Sequences identified in such library screening methods can be compared and aligned to other known sequences deposited and available in public databases such as GenBank or other private sequence databases. Sequence identity (at either the amino acid or nucleotide level) within defined regions of the molecule or across the full-length sequence can be determined

through sequence alignment using computer software programs such as BLAST, BLAST2, ALIGN-2, DNASTar, and INHERIT which employ various algorithms to measure homology.

Nucleic acid having protein coding sequence may be obtained by screening selected cDNA or genomic libraries using the deduced amino acid sequence disclosed herein for the first time, and, if necessary, using conventional primer extension procedures as described in Sambrook *et al.*, *supra*, to detect precursors and processing intermediates of mRNA that may not have been reverse-transcribed into cDNA.

2. Selection and Transformation of Host Cells

Host cells are transfected or transformed with expression or cloning vectors described herein for IL-1p production and cultured in conventional nutrient media modified as appropriate for inducing promoters, selecting transformants, or amplifying the genes encoding the desired sequences. The culture conditions, such as media, temperature, pH and the like, can be selected by the skilled artisan without undue experimentation. In general, principles, protocols, and practical techniques for maximizing the productivity of cell cultures can be found in *Mammalian Cell Biotechnology: A Practical Approach*, M. Butler, ed. (IRL Press, 1991) and Sambrook *et al.*, *supra*.

Methods of transfection are known to the ordinarily skilled artisan, for example, CaPO₄ and electroporation. Depending on the host cell used, transformation is performed using standard techniques appropriate to such cells. The calcium treatment employing calcium chloride, as described in Sambrook *et al.*, *supra*, or electroporation is generally used for prokaryotes or other cells that contain substantial cell-wall barriers. Infection with *Agrobacterium tumefaciens* is used for transformation of certain plant cells, as described by Shaw *et al.*, *Gene*, **23**:315 (1983) and WO 89/05859 published 29 June 1989. For mammalian cells without such cell walls, the calcium phosphate precipitation method of Graham and van der Eb, *Virology*, **52**:456-457 (1978) can be employed. General aspects of mammalian cell host system transformations have been described in U.S. Patent No. 4,399,216. Transformations into yeast are typically carried out according to the method of Van Solingen *et al.*, *J. Bact.*, **130**:946 (1977) and Hsiao *et al.*, *Proc. Natl. Acad. Sci. (USA)*, **76**:3829 (1979). However, other methods for introducing DNA into cells, such as by nuclear microinjection, electroporation, bacterial protoplast fusion with intact cells, or polycations, *e.g.*, polybrene, polyornithine, may also be used. For various techniques for transforming mammalian cells, see Keown *et al.*, *Methods in Enzymology*, **185**:527-537 (1990) and Mansour *et al.*, *Nature*, **336**:348-352 (1988).

Suitable host cells for cloning or expressing the DNA in the vectors herein include prokaryote, yeast, or higher eukaryote cells. Suitable prokaryotes include but are not limited to eubacteria, such as Gram-negative or Gram-positive organisms, for example, Enterobacteriaceae such as *E. coli*. Various *E. coli* strains are publicly available, such as *E. coli* K12 strain MM294 (ATCC 31,446); *E. coli* X1776 (ATCC 31,537); *E. coli* strain W3110 (ATCC 27,325) and K5 772 (ATCC 53,635).

In addition to prokaryotes, eukaryotic microbes such as filamentous fungi or yeast are suitable cloning or expression hosts for IL-11p-encoding vectors. *Saccharomyces cerevisiae* is a commonly used lower eukaryotic host microorganism.

Suitable host cells for the expression of glycosylated IL-11p are derived from multicellular organisms. Examples of invertebrate cells include insect cells such as *Drosophila* S2 and *Spodoptera* Sf9, as well as plant cells. Examples of useful mammalian host cell lines include Chinese hamster ovary (CHO) and COS cells. More specific examples include monkey kidney CV1 line transformed by SV40 (COS-7, ATCC CRL 1651); human embryonic kidney line (293 or 293 cells subcloned for growth in suspension culture, Graham *et al.*, *J. Gen Virol.*, 36:59 (1977)); Chinese hamster ovary cells/-DHFR (CHO, Urlaub and Chasin, *Proc. Natl. Acad. Sci. USA*, 77:4216 (1980)); mouse sertoli cells (TM4, Mather, *Biol. Reprod.*, 23:243-251 (1980)); human lung cells (W138, ATCC CCL 75); human liver cells (Hep G2, HB 8065); and mouse mammary tumor (MMT 060562, ATCC CCL51). The selection of the appropriate host cell is deemed to be within the skill in the art.

3. Selection and Use of a Replicable Vector

The nucleic acid (e.g., cDNA or genomic DNA) encoding IL-11p may be inserted into a replicable vector for cloning (amplification of the DNA) or for expression. Various vectors are publicly available. The vector may, for example, be in the form of a plasmid, cosmid, viral particle, or phage. The appropriate nucleic acid sequence may be inserted into the vector by a variety of procedures. In general, DNA is inserted into an appropriate restriction endonuclease site(s) using techniques known in the art. Vector components generally include, but are not limited to, one or more of a signal sequence, an origin of replication, one or more marker genes, an enhancer element, a promoter, and a transcription termination sequence. Construction of suitable vectors containing one or more of these components employs standard ligation techniques which are known to the skilled artisan.

The IL-11p may be produced recombinantly not only directly, but also as a fusion polypeptide with a heterologous polypeptide, which may be a signal sequence or other polypeptide having a specific cleavage site at the N-terminus of the mature protein or polypeptide. In general, the signal sequence may be a component of the vector, or it may be a part of the IL-11p-encoding DNA that is inserted into the vector. The signal sequence may be a prokaryotic signal sequence selected, for example, from the group of the alkaline phosphatase, penicillinase, lpp, or heat-stable enterotoxin II leaders. For yeast secretion the signal sequence may be, e.g., the yeast invertase leader, alpha factor leader (including *Saccharomyces* and *Kluyveromyces* α -factor leaders, the latter described in U.S. Patent No. 5,010,182), or acid phosphatase leader, the *C. albicans* glucoamylase leader (EP 362,179 published 4 April 1990), or the signal described in WO 90/13646 published 15 November 1990. In mammalian cell expression, mammalian signal sequences may be used to direct secretion of the protein, such as signal sequences from secreted polypeptides of the same or related species, as well as viral secretory leaders.

Both expression and cloning vectors contain a nucleic acid sequence that enables the vector to replicate in one or more selected host cells. Such sequences are well known for a variety of bacteria, yeast, and viruses. The origin of replication from the plasmid pBR322 is suitable for most Gram-negative bacteria, the 2 μ plasmid origin is suitable for yeast, and various viral origins (SV40, polyoma, adenovirus, VSV or BPV) are useful for cloning vectors in mammalian cells.

Expression and cloning vectors will typically contain a selection gene, also termed a selectable marker. Typical selection genes encode proteins that (a) confer resistance to antibiotics or other toxins, e.g., ampicillin, neomycin, methotrexate, or tetracycline, (b) complement auxotrophic deficiencies, or (c) supply critical nutrients not available from complex media, e.g., the gene encoding D-alanine racemase for *Bacilli*.

An example of suitable selectable markers for mammalian cells are those that enable the identification of cells competent to take up the IL-11p-encoding nucleic acid, such as DHFR or thymidine kinase. An appropriate host cell when wild-type DHFR is employed is the CHO cell line deficient in DHFR activity, prepared and propagated as described by Urlaub *et al.*, *Proc. Natl. Acad. Sci. USA*, 77:4216 (1980). A suitable selection gene for use in yeast is the *trp1* gene present in the yeast plasmid YRp7 [Stinchcomb *et al.*, *Nature*, 282:39 (1979); Kingsman *et al.*, *Gene*, 7:141 (1979); Tschemper *et al.*, *Gene*, 10:157 (1980)]. The *trp1* gene provides a selection marker for a mutant strain of yeast lacking the ability to grow in tryptophan, for example, ATCC No. 44076 or PEP4-1 [Jones, *Genetics*, 85:12 (1977)].

Expression and cloning vectors usually contain a promoter operably linked to the IL-11p-encoding nucleic acid sequence to direct mRNA synthesis. Promoters recognized by a variety of potential host cells are well known. Promoters suitable for use with prokaryotic hosts include the β -lactamase and lactose promoter systems [Chang *et al.*, *Nature*, 275:615 (1978); Goeddel *et al.*, *Nature*, 281:544 (1979)], alkaline phosphatase, a tryptophan (*trp*) promoter system [Goeddel, *Nucleic Acids Res.*, 8:4057 (1980); EP 36,776], and hybrid promoters such as the *tac* promoter [deBoer *et al.*, *Proc. Natl. Acad. Sci. USA*, 80:21-25 (1983)]. Promoters for use in bacterial systems also will contain a Shine-Dalgarno (S.D.) sequence operably linked to the DNA encoding IL-11p.

Examples of suitable promoting sequences for use with yeast hosts include the promoters for 3-phosphoglycerate kinase [Hitzeman *et al.*, *J. Biol. Chem.*, 255:2073 (1980)] or other glycolytic enzymes [Hess *et al.*, *J. Adv. Enzyme Reg.*, 7:149 (1968); Holland, *Biochemistry*, 17:4900 (1978)], such as enolase, glyceraldehyde-3-phosphate dehydrogenase, hexokinase, pyruvate decarboxylase, phosphofructokinase, glucose-6-phosphate isomerase, 3-phosphoglycerate mutase, pyruvate kinase, triosephosphate isomerase, phosphoglucose isomerase, and glucokinase.

Other yeast promoters, which are inducible promoters having the additional advantage of transcription controlled by growth conditions, are the promoter regions for alcohol dehydrogenase 2, isocytochrome C, acid phosphatase, degradative enzymes associated with nitrogen metabolism, metallothionein, glyceraldehyde-3-phosphate dehydrogenase, and enzymes

responsible for maltose and galactose utilization. Suitable vectors and promoters for use in yeast expression are further described in EP 73,657.

IL-11p transcription from vectors in mammalian host cells is controlled, for example, by promoters obtained from the genomes of viruses such as polyoma virus, fowlpox virus (UK 2,211,504 published 5 July 1989), adenovirus (such as Adenovirus 2), bovine papilloma virus, avian sarcoma virus, cytomegalovirus, a retrovirus, hepatitis-B virus and Simian Virus 40 (SV40), from heterologous mammalian promoters, *e.g.*, the actin promoter or an immunoglobulin promoter, and from heat-shock promoters, provided such promoters are compatible with the host cell systems.

Transcription of a DNA encoding the IL-11p by higher eukaryotes may be increased by inserting an enhancer sequence into the vector. Enhancers are cis-acting elements of DNA, usually about from 10 to 300 bp, that act on a promoter to increase its transcription. Many enhancer sequences are now known from mammalian genes (globin, elastase, albumin, α -fetoprotein, and insulin). Typically, however, one will use an enhancer from a eukaryotic cell virus. Examples include the SV40 enhancer on the late side of the replication origin (bp 100-270), the cytomegalovirus early promoter enhancer, the polyoma enhancer on the late side of the replication origin, and adenovirus enhancers. The enhancer may be spliced into the vector at a position 5' or 3' to the IL-11p coding sequence, but is preferably located at a site 5' from the promoter.

Expression vectors used in eukaryotic host cells (yeast, fungi, insect, plant, animal, human, or nucleated cells from other multicellular organisms) will also contain sequences necessary for the termination of transcription and for stabilizing the mRNA. Such sequences are commonly available from the 5' and, occasionally 3', untranslated regions of eukaryotic or viral DNAs or cDNAs. These regions contain nucleotide segments transcribed as polyadenylated fragments in the untranslated portion of the mRNA encoding IL-11p.

Still other methods, vectors, and host cells suitable for adaptation to the synthesis of IL-11p in recombinant vertebrate cell culture are described in Gething *et al.*, *Nature*, 293:620-625 (1981); Mantei *et al.*, *Nature*, 281:40-46 (1979); EP 117,060; and EP 117,058.

4. Detecting Gene Amplification/Expression

Gene amplification and/or expression may be measured in a sample directly, for example, by conventional Southern blotting, Northern blotting to quantitate the transcription of mRNA [Thomas, *Proc. Natl. Acad. Sci. USA*, 77:5201-5205 (1980)], dot blotting (DNA analysis), or *in situ* hybridization, using an appropriately labeled probe, based on the sequences provided herein. Alternatively, antibodies may be employed that can recognize specific duplexes, including DNA duplexes, RNA duplexes, and DNA-RNA hybrid duplexes or DNA-protein duplexes. The antibodies in turn may be labeled and the assay may be carried out where the duplex is bound to a surface, so that upon the formation of duplex on the surface, the presence of antibody bound to the duplex can be detected.

Gene expression, alternatively, may be measured by immunological methods, such as immunohistochemical staining of cells or tissue sections and assay of cell culture or body fluids,

to quantitate directly the expression of gene product. Antibodies useful for immunohistochemical staining and/or assay of sample fluids may be either monoclonal or polyclonal, and may be prepared in any mammal. Conveniently, the antibodies may be prepared against a native sequence IL-11p polypeptide or against a synthetic peptide based on the DNA sequences provided herein or against exogenous sequence fused to IL-11p DNA and encoding a specific antibody epitope.

5. Purification of Polypeptide

Forms of IL-11p may be recovered from culture medium or from host cell lysates. If membrane-bound, it can be released from the membrane using a suitable detergent solution (e.g. Triton-X 100) or by enzymatic cleavage. Cells employed in expression of IL-11p can be disrupted by various physical or chemical means, such as freeze-thaw cycling, sonication, mechanical disruption, or cell lysing agents. It may be desired to purify IL-11p from recombinant cell proteins or polypeptides. The following procedures are exemplary of suitable purification procedures: by fractionation on an ion-exchange column; ethanol precipitation; reverse phase HPLC; chromatography on silica or on a cation-exchange resin such as DEAE; chromatofocusing; SDS-PAGE; ammonium sulfate precipitation; gel filtration using, for example, Sephadex G-75; protein A Sepharose columns to remove contaminants such as IgG; and metal chelating columns to bind epitope-tagged forms of the IL-11p. Various methods of protein purification may be employed and such methods are known in the art and described for example in Deutscher, *Methods in Enzymology*, 182 (1990); Scopes, *Protein Purification: Principles and Practice*, Springer-Verlag, New York (1982). The purification step(s) selected will depend, for example, on the nature of the production process used and the particular IL-11p produced.

B. Activity Assays for IL-11p Variants

The biological activity or activities of a particular IL-11p variant polypeptide can be characterized using a variety of in vitro assays known in the art. For example, the ability of an hIL-1Ra3 variant polypeptide or a mL-1Ra3 variant polypeptide to bind IL-1R can be assayed using a radioimmunoprecipitation assay wherein IL-1R extracellular domain (ECD) fused to the Fc region of human immunoglobulin G (IL-1R ECD-Fc) (which can be prepared, e.g., as described in Examples 9 and 10 below) is incubated in solution with radiolabeled hIL-1Ra3 variant polypeptide or mL-1Ra3 variant polypeptide to form labeled complexes, followed by immunoprecipitation of the labeled complexes with goat anti-human IgG Fc and quantitation of radioactivity in the precipitate. In another example, an hIL-1Ra3 variant polypeptide-FLAG tag fusion protein-encoding DNA and an IL-1R ECD-Fc encoding DNA can be coexpressed in a host cell and secreted into the cell's culture medium, followed by immunoprecipitation of culture supernatant with protein G-sepharose and identification of bound hIL-1Ra3 variant polypeptide-FLAG tag fusion protein by immunoblotting with anti-FLAG monoclonal antibody, essentially as described in Example 9 below.

In another embodiment, the ability of an hIL-1Ra3 variant polypeptide or a mL-1Ra3 variant polypeptide to inhibit the binding of IL-1 to IL-1R can be assayed using a competitive binding assay. For example, a radioimmunoprecipitation assay can be employed wherein IL-1R ECD-Fc is incubated in solution of radiolabeled IL-1 with or without unlabeled hIL-1Ra3 variant

polypeptide or unlabeled mIL-1Ra3 variant polypeptide to form labeled or unlabeled complexes, followed by immunoprecipitation of complexes with anti-human IgG Fc and quantitation of radioactivity in the precipitate. If the presence of unlabeled hIL-1Ra3 variant polypeptide or unlabeled mIL-1Ra3 variant polypeptide in the incubation solution diminishes the radioactivity measured in the resulting immunoprecipitate, the hIL-1Ra3 variant polypeptide or mIL-1Ra3 variant polypeptide in question qualifies as an inhibitor of IL-1 binding to IL-1R. In yet another embodiment, IL-1R ECD-Fc and an hIL-1Ra3 variant-FLAG tag fusion protein or mIL-1Ra3 variant-FLAG tag fusion protein are obtained by recombinant expression in separate cell cultures (essentially as described in Example 10 below), IL-1 and IL-1R ECD-Fc are admixed together with or without the hIL-1Ra3 variant-FLAG tag fusion protein or mIL-1Ra3 variant-FLAG tag fusion protein and incubated in solution, the incubation solution is immunoprecipitated with protein G-sepharose, and the bound hIL-1Ra3 variant-FLAG tag fusion protein or mIL-1Ra3 variant-FLAG tag fusion protein is identified by immunoblotting with anti-FLAG monoclonal antibody. If the presence of IL-1 in the incubation solution diminishes the signal detected by anti-FLAG immunoblotting, the hIL-1Ra3 variant polypeptide or mIL-1Ra3 variant polypeptide in question qualifies as an inhibitor of IL-1 binding to IL-1R.

Similarly, the biological activity or activities of a particular hIL-1Ra1 variant polypeptide can be determined by using a variety of in vitro assays known in the art. For example, the ability of an hIL-1Ra1 variant polypeptide to bind IL-18R can be assayed using a radioimmunoprecipitation assay wherein IL-18R extracellular domain (ECD) fused to the Fc region of human immunoglobulin G (IL-18R ECD-Fc) (which can be prepared, e.g., as described in Examples 9 and 10 below) is incubated in solution with radiolabeled hIL-1Ra1 variant polypeptide to form labeled complex, followed by immunoprecipitation of the labeled complex with goat anti-human IgG Fc and quantitation of radioactivity in the precipitate. In another example, an hIL-1Ra1 variant polypeptide-FLAG tag fusion protein-encoding DNA and an IL-18R ECD-Fc encoding DNA can be coexpressed in a host cell and secreted into the cell's culture medium, followed by immunoprecipitation of culture supernatant with protein G-sepharose and identification of bound hIL-1Ra1 variant polypeptide-FLAG tag fusion protein by immunoblotting with anti-FLAG monoclonal antibody, essentially as described in Example 9 below.

In another embodiment, the ability of an hIL-1Ra1 variant polypeptide to inhibit the binding of IL-18 to IL-18R can be assayed using a competitive binding assay. For example, a radioimmunoprecipitation assay can be employed wherein IL-18R ECD-Fc is incubated in solution of radiolabeled IL-18 with or without unlabeled hIL-1Ra1 variant polypeptide to form labeled or unlabeled complexes, followed by immunoprecipitation of complexes with anti-human IgG Fc and quantitation of radioactivity in the precipitate. If the presence of unlabeled hIL-1Ra1 variant polypeptide in the incubation solution diminishes the radioactivity measured in the resulting immunoprecipitate, the hIL-1Ra1 variant polypeptide in question qualifies as an inhibitor of IL-18 binding to IL-18R. In yet another embodiment, IL-18R ECD-Fc and an hIL-1Ra1 variant-FLAG tag fusion protein are obtained by recombinant expression in separate cell cultures (essentially as described in Example 10 below), IL-18 and IL-18R ECD-Fc are admixed together with or without the hIL-1Ra1 variant-FLAG tag fusion protein and incubated in

solution, the incubation solution is immunoprecipitated with protein G-sepharose, and the bound hIL-1Ra1 variant-FLAG tag fusion protein is identified by immunoblotting with anti-FLAG monoclonal antibody. If the presence of IL-18 in the incubation solution diminishes the signal detected by anti-FLAG immunoblotting, the hIL-1Ra1 variant polypeptide in question qualifies as an inhibitor of IL-18 binding to IL-18R.

C. Uses for IL-1lp

Nucleotide sequences (or their complement) encoding IL-1lp have various applications in the art of molecular biology, including uses as hybridization probes, in chromosome and gene mapping and in the generation of anti-sense RNA and DNA. IL-1lp nucleic acid will also be useful for the preparation of IL-1lp polypeptides by the recombinant techniques described herein.

The full-length native sequence IL-1lp genes of Figure 1 (SEQ ID NO:1), Figure 2 (SEQ ID NO:4), Figure 3 (SEQ ID NO:6), Figure 5 (SEQ ID NO:9), Figure 7 (SEQ ID NO:12), Figure 9 (SEQ ID NO:15), Figure 15 (SEQ ID NO:18), and Figure 16 (SEQ ID NO:20), and Figure 19 (SEQ ID NO:24), or portions thereof, may be used as hybridization probes for a cDNA library to isolate the full-length IL-1lp gene or to isolate still other genes (for instance, those encoding naturally-occurring variants of IL-1lp or IL-1lp from other species) which have a desired sequence identity to the IL-1lp sequence disclosed in Figure 1 (SEQ ID NO:1), Figure 2 (SEQ ID NO:4), Figure 3 (SEQ ID NO:6), Figure 5 (SEQ ID NO:9), Figure 7 (SEQ ID NO:12), Figure 9 (SEQ ID NO:15), Figure 15 (SEQ ID NO:18), Figure 16 (SEQ ID NO:20), or Figure 19 (SEQ ID NO:24). Optionally, the length of the probes will be about 20 to about 50 bases. The hybridization probes may be derived from the nucleotide sequence of Figure 1 (SEQ ID NO:1), Figure 2 (SEQ ID NO:4), Figure 3 (SEQ ID NO:6), Figure 5 (SEQ ID NO:9), Figure 7 (SEQ ID NO:12), Figure 9 (SEQ ID NO:15), Figure 15 (SEQ ID NO:18), Figure 16 (SEQ ID NO:20) or Figure 19 (SEQ ID NO:24), or from genomic sequences including promoters, enhancer elements and introns of native sequence IL-1lp. By way of example, a screening method will comprise isolating the coding region of the IL-1lp gene using the known DNA sequence to synthesize a selected probe of about 40 bases. Hybridization probes may be labeled by a variety of labels, including radionucleotides such as ³²P or ³⁵S, or enzymatic labels such as alkaline phosphatase coupled to the probe via avidin/biotin coupling systems. Labeled probes having a sequence complementary to that of the IL-1lp gene of the present invention can be used to screen libraries of human cDNA, genomic DNA or mRNA to determine which members of such libraries the probe hybridizes to. Hybridization techniques are described in further detail in the Examples below.

The probes may also be employed in PCR techniques to generate a pool of sequences for identification of closely related IL-1lp coding sequences.

Nucleotide sequences encoding an IL-1lp can also be used to construct hybridization probes for mapping the gene which encodes that IL-1lp and for the genetic analysis of individuals with genetic disorders. The nucleotide sequences provided herein may be mapped to a chromosome and specific regions of a chromosome using known techniques, such as *in situ* hybridization, linkage analysis against known chromosomal markers, and hybridization screening with libraries.

When the coding sequences for IL-11p encode a protein which binds to another protein (example, where the IL-11p binds to an IL-1 receptor or IL-18 receptor), the IL-11p can be used in assays to identify the other proteins or molecules involved in the binding interaction. By such methods, inhibitors of the receptor/ligand binding interaction can be identified. Proteins involved in such binding interactions can also be used to screen for peptide or small molecule inhibitors or agonists of the binding interaction. Screening assays can be designed to find lead compounds that mimic the biological activity of a native IL-11p or a receptor for IL-11p. Such screening assays will include assays amenable to high-throughput screening of chemical libraries, making them particularly suitable for identifying small molecule drug candidates.

Small molecules contemplated include synthetic organic or inorganic compounds. The assays can be performed in a variety of formats, including protein-protein binding assays, biochemical screening assays, immunoassays and cell based assays, which are well characterized in the art.

Nucleic acids which encode IL-11p or its modified forms can also be used to generate either transgenic animals or "knock out" animals which, in turn, are useful in the development and screening of therapeutically useful reagents. A transgenic animal (e.g., a mouse or rat) is an animal having cells that contain a transgene, which transgene was introduced into the animal or an ancestor of the animal at a prenatal, e.g., an embryonic stage. A transgene is a DNA which is integrated into the genome of a cell from which a transgenic animal develops. In one embodiment, cDNA encoding IL-11p can be used to clone genomic DNA encoding IL-11p in accordance with established techniques and the genomic sequences used to generate transgenic animals that contain cells which express DNA encoding IL-11p. Methods for generating transgenic animals, particularly animals such as mice or rats, have become conventional in the art and are described, for example, in U.S. Patent Nos. 4,736,866 and 4,870,009. Typically, particular cells would be targeted for IL-11p transgene incorporation with tissue-specific enhancers. Transgenic animals that include a copy of a transgene encoding IL-11p introduced into the germ line of the animal at an embryonic stage can be used to examine the effect of increased expression of DNA encoding IL-11p. Such animals can be used as tester animals for reagents thought to confer protection from, for example, pathological conditions associated with its overexpression. In accordance with this facet of the invention, an animal is treated with the reagent and a reduced incidence of the pathological condition, compared to untreated animals bearing the transgene, would indicate a potential therapeutic intervention for the pathological condition.

Alternatively, non-human homologues of IL-11p can be used to construct an IL-11p "knock out" animal which has a defective or altered gene encoding IL-11p as a result of homologous recombination between the endogenous gene encoding IL-11p and altered genomic DNA encoding IL-11p introduced into an embryonic cell of the animal. For example, cDNA encoding IL-11p can be used to clone genomic DNA encoding IL-11p in accordance with established techniques. A portion of the genomic DNA encoding IL-11p can be deleted or replaced with another gene, such as a gene encoding a selectable marker which can be used to monitor integration. Typically, several kilobases of unaltered flanking DNA (both at the 5' and 3' ends) are included in the vector [see e.g., Thomas and Capecchi, *Cell*, 51:503 (1987) for a description

of homologous recombination vectors]. The vector is introduced into an embryonic stem cell line (e.g., by electroporation) and cells in which the introduced DNA has homologously recombined with the endogenous DNA are selected [see e.g., Li et al., *Cell*, 69:915 (1992)]. The selected cells are then injected into a blastocyst of an animal (e.g., a mouse or rat) to form aggregation chimeras [see e.g., Bradley, in *Teratocarcinomas and Embryonic Stem Cells: A Practical Approach*, E. J. Robertson, ed. (IRL, Oxford, 1987), pp. 113-152]. A chimeric embryo can then be implanted into a suitable pseudopregnant female foster animal and the embryo brought to term to create a "knock out" animal. Progeny harboring the homologously recombined DNA in their germ cells can be identified by standard techniques and used to breed animals in which all cells of the animal contain the homologously recombined DNA. Knockout animals can be characterized for instance, for their ability to defend against certain pathological conditions and for their development of pathological conditions due to absence of the IL-11p polypeptide.

Nucleic acid encoding the IL-11p polypeptides may also be used in gene therapy. In gene therapy applications, genes are introduced into cells in order to achieve *in vivo* synthesis of a therapeutically effective genetic product, for example for replacement of a defective gene. "Gene therapy" includes both conventional gene therapy where a lasting effect is achieved by a single treatment, and the administration of gene therapeutic agents, which involves the one time or repeated administration of a therapeutically effective DNA or mRNA. Antisense RNAs and DNAs can be used as therapeutic agents for blocking the expression of certain genes *in vivo*. It has already been shown that short antisense oligonucleotides can be imported into cells where they act as inhibitors, despite their low intracellular concentrations caused by their restricted uptake by the cell membrane. (Zamecnik et al., *Proc. Natl. Acad. Sci. USA* 83, 4143-4146 [1986]). The oligonucleotides can be modified to enhance their uptake, e.g. by substituting their negatively charged phosphodiester groups by uncharged groups.

There are a variety of techniques available for introducing nucleic acids into viable cells. The techniques vary depending upon whether the nucleic acid is transferred into cultured cells *in vitro*, or *in vivo* in the cells of the intended host. Techniques suitable for the transfer of nucleic acid into mammalian cells *in vitro* include the use of liposomes, electroporation, microinjection, cell fusion, DEAE-dextran, the calcium phosphate precipitation method, etc. The currently preferred *in vivo* gene transfer techniques include transfection with viral (typically retroviral) vectors and viral coat protein-liposome mediated transfection (Dzau et al., *Trends in Biotechnology* 11, 205-210 [1993]). In some situations it is desirable to provide the nucleic acid source with an agent that targets the target cells, such as an antibody specific for a cell surface membrane protein or the target cell, a ligand for a receptor on the target cell, etc. Where liposomes are employed, proteins which bind to a cell surface membrane protein associated with endocytosis may be used for targeting and/or to facilitate uptake, e.g. capsid proteins or fragments thereof tropic for a particular cell type, antibodies for proteins which undergo internalization in cycling, proteins that target intracellular localization and enhance intracellular half-life. The technique of receptor-mediated endocytosis is described, for example, by Wu et al., *J. Biol. Chem.* 262, 4429-4432 (1987); and Wagner et al., *Proc. Natl. Acad. Sci. USA* 87, 3410-

3414 (1990). For review of gene marking and gene therapy protocols see Anderson *et al.*, *Science* 256, 808-813 (1992).

The IL-11p polypeptides of the present invention can be formulated according to known methods to prepare pharmaceutically useful compositions, whereby the IL-11p product hereof is combined in admixture with a pharmaceutically acceptable carrier vehicle. Therapeutic formulations are prepared for storage by mixing the active ingredient having the desired degree of purity with optional physiologically acceptable carriers, excipients or stabilizers (*Remington's Pharmaceutical Sciences* 16th edition, Osol, A. Ed. (1980)), in the form of lyophilized formulations or aqueous solutions. Acceptable carriers, excipients or stabilizers are nontoxic to recipients at the dosages and concentrations employed, and include buffers such as phosphate, citrate and other organic acids; antioxidants including ascorbic acid; low molecular weight (less than about 10 residues) polypeptides; proteins, such as serum albumin, gelatin or immunoglobulins; hydrophilic polymers such as polyvinylpyrrolidone, amino acids such as glycine, glutamine, asparagine, arginine or lysine; monosaccharides, disaccharides and other carbohydrates including glucose, mannose, or dextrans; chelating agents such as EDTA; sugar alcohols such as mannitol or sorbitol; salt-forming counterions such as sodium; and/or nonionic surfactants such as Tween, Plurionics or PEG.

The formulations to be used for *in vivo* administration must be sterile. This is readily accomplished by filtration through sterile filtration membranes, prior to or following lyophilization and reconstitution.

Therapeutic compositions herein generally are placed into a container having a sterile access port, for example, an intravenous solution bag or vial having a stopper pierceable by a hypodermic injection needle.

The route of administration is in accord with known methods, e.g. injection or infusion by intravenous, intraperitoneal, intracerebral, intramuscular, intraocular, intraarterial or intralesional routes, topical administration, or by sustained release systems.

Dosages and desired drug concentrations of pharmaceutical compositions of the present invention may vary depending on the particular use envisioned. The determination of the appropriate dosage or route of administration is well within the skill of an ordinary physician.

Animal experiments provide reliable guidance for the determination of effective doses for human therapy. Interspecies scaling of effective doses can be performed following the principles laid down by Mordenti, J. and Chappell, W. "The use of interspecies scaling in toxicokinetics" *In* Toxicokinetics and New Drug Development, Yacobi *et al.*, Eds., Pergamon Press, New York 1989, pp. 42-96.

An "effective amount" of the IL-11p to be employed therapeutically will depend, for example, upon the therapeutic objectives, the route of administration, and the condition of the patient. Accordingly, it will be necessary for the therapist to titer the dosage and modify the route of administration as required to obtain the optimal therapeutic effect. Typically, the clinician will administer the IL-11p until a dosage is reached that achieves the desired effect. The progress of this therapy is easily monitored by conventional assays.

In one embodiment, the invention provides a method for treating an IL-1-mediated disorder comprising administering to a mammal, such as human, in need of such treatment an effective amount of an IL-1lp, such as a native sequence IL-1lp.

In another embodiment, the invention provides a method for treating an IL-1-mediated disorder comprising administering to a mammal, such as human, in need of such treatment an effective amount of an IL-1lp selected from the group consisting of hIL-1Ra1, hIL-1Ra1L, hIL-1Ra1V, hIL-1Ra1S, hIL-1Ra2, hIL-1Ra3, and mL-1Ra3.

In another embodiment, the invention provides a method for treating an IL-1-mediated disorder comprising administering to a mammal, such as human, in need of such treatment an effective amount of an hIL-1lp, such as a native sequence hIL-1lp, e.g. native sequence hIL-1Ra3.

In another embodiment, the invention provides a method for treating an IL-18-mediated disorder comprising administering to a mammal, such as human, in need of such treatment an effective amount of an IL-1lp, such as a native sequence IL-1lp.

In another embodiment, the invention provides a method for treating an IL-18-mediated disorder comprising administering to a mammal, such as human, in need of such treatment an effective amount of an IL-1lp selected from the group consisting of hIL-1Ra1, hIL-1Ra1L, hIL-1Ra1V, hIL-1Ra1S, hIL-1Ra2, hIL-1Ra3, and mL-1Ra3.

In another embodiment, the invention provides a method for treating an IL-18-mediated disorder comprising administering to a mammal, such as human, in need of such treatment an effective amount of an hIL-1lp, such as a native sequence hIL-1lp, e.g. native sequence hIL-1Ra1.

In another embodiment, the invention provides a method for treating an IL-18-mediated disorder comprising administering to a mammal, such as human, in need of such treatment an effective amount of an hIL-1Ra1L, such as a native sequence hIL-1Ra1L, or an effective amount of an hIL-1Ra1V, such as a native sequence hIL-1Ra1V, or an effective amount of an hIL-1Ra1S, such as a native sequence hIL-1Ra1S.

In one embodiment, the invention provides a method for treating an inflammatory disorder comprising administering to a mammal, such as human, in need of such treatment an effective amount of an IL-1lp, such as a native sequence IL-1lp.

In another embodiment, the invention provides a method for treating an inflammatory disorder comprising administering to a mammal, such as human, in need of such treatment an effective amount of an IL-1lp selected from the group consisting of hIL-1Ra1, hIL-1Ra1L, hIL-1Ra1V, hIL-1Ra1S, hIL-1Ra2, hIL-1Ra3, and mL-1Ra3.

In another embodiment, the invention provides a method for treating an inflammatory disorder comprising administering to a mammal, such as human, in need of such treatment an effective amount of an hIL-1lp, such as a native sequence hIL-1lp, e.g. native sequence hIL-1Ra1 or hIL-1Ra3.

In another embodiment, the invention provides a method for treating an inflammatory disorder comprising administering to a mammal, such as human, in need of such treatment an effective amount of an hIL-1Ra1L, such as a native sequence hIL-1Ra1L, or an effective amount of an hIL-1Ra1V, such as a native sequence hIL-1Ra1V, or an effective amount of an hIL-1Ra1S, such as a native sequence hIL-1Ra1S.

In another embodiment, the invention provides a method for treating asthma comprising administering to a mammal, such as human, in need of such treatment an effective amount of an IL-11p, such as a native sequence IL-11p.

In another embodiment, the invention provides a method for treating asthma comprising administering to a mammal, such as human, in need of such treatment an effective amount of an IL-11p selected from the group consisting of hIL-1Ra1, hIL-1Ra1L, hIL-1Ra1V, hIL-1Ra1S, hIL-1Ra2, hIL-1Ra3, and mL-1Ra3.

In another embodiment, the invention provides a method for treating asthma comprising administering to a mammal, such as human, in need of such treatment an effective amount of an hIL-11p, such as a native sequence hIL-11p, e.g. native sequence hIL-1Ra1 or hIL-1Ra3.

In another embodiment, the invention provides a method for treating asthma comprising administering to a mammal, such as human, in need of such treatment an effective amount of an hIL-1Ra1L, such as a native sequence hIL-1Ra1L, or an effective amount of an hIL-1Ra1V, such as a native sequence hIL-1Ra1V, or an effective amount of an hIL-1Ra1S, such as a native sequence hIL-1Ra1S.

In another embodiment, the invention provides a method for treating rheumatoid arthritis comprising administering to a mammal, such as human, in need of such treatment an effective amount of an IL-11p, such as a native sequence IL-11p.

In another embodiment, the invention provides a method for treating rheumatoid arthritis comprising administering to a mammal, such as human, in need of such treatment an effective amount of an IL-11p selected from the group consisting of hIL-1Ra1, hIL-1Ra1L, hIL-1Ra1V, hIL-1Ra1S, hIL-1Ra2, hIL-1Ra3, and mL-1Ra3.

In another embodiment, the invention provides a method for treating rheumatoid arthritis comprising administering to a mammal, such as human, in need of such treatment an effective amount of an hIL-11p, such as a native sequence hIL-11p, e.g. native sequence hIL-1Ra1 or hIL-1Ra3.

In another embodiment, the invention provides a method for treating rheumatoid arthritis comprising administering to a mammal, such as human, in need of such treatment an effective amount of an hIL-1Ra1L, such as a native sequence hIL-1Ra1L, or an effective amount of an hIL-1Ra1V, such as a native sequence hIL-1Ra1V, or an effective amount of an hIL-1Ra1S, such as a native sequence hIL-1Ra1S.

In another embodiment, the invention provides a method for treating osteoarthritis comprising administering to a mammal, such as human, in need of such treatment an effective amount of an IL-11p, such as a native sequence IL-11p.

In another embodiment, the invention provides a method for treating osteoarthritis comprising administering to a mammal, such as human, in need of such treatment an effective amount of an IL-11p selected from the group consisting of hIL-1Ra1, hIL-1Ra1L, hIL-1Ra1V, hIL-1Ra1S, hIL-1Ra2, hIL-1Ra3, and mL-1Ra3.

In another embodiment, the invention provides a method for treating osteoarthritis comprising administering to a mammal, such as human, in need of such treatment an effective

amount of an hIL-11p, such as a native sequence hIL-11p, e.g. native sequence hIL-1Ra1 or hIL-1Ra3.

In another embodiment, the invention provides a method for treating osteoarthritis comprising administering to a mammal, such as human, in need of such treatment an effective amount of an hIL-1Ra1L, such as a native sequence hIL-1Ra1L, or an effective amount of an hIL-1Ra1V, such as a native sequence hIL-1Ra1V, or an effective amount of an hIL-1Ra1S, such as a native sequence hIL-1Ra1S.

In another embodiment, the invention provides a method for treating sepsis comprising administering to a mammal, such as human, in need of such treatment an effective amount of an IL-11p, such as a native sequence IL-11p.

In another embodiment, the invention provides a method for treating sepsis comprising administering to a mammal, such as human, in need of such treatment an effective amount of an IL-11p selected from the group consisting of hIL-1Ra1, hIL-1Ra1L, hIL-1Ra1V, hIL-1Ra1S, hIL-1Ra2, hIL-1Ra3, and mLIL-1Ra3.

In another embodiment, the invention provides a method for treating sepsis comprising administering to a mammal, such as human, in need of such treatment an effective amount of an hIL-11p, such as a native sequence hIL-11p, e.g. native sequence hIL-1Ra1 or hIL-1Ra3.

In another embodiment, the invention provides a method for treating sepsis comprising administering to a mammal, such as human, in need of such treatment an effective amount of an hIL-1Ra1L, such as a native sequence hIL-1Ra1L, or an effective amount of an hIL-1Ra1V, such as a native sequence hIL-1Ra1V, or an effective amount of an hIL-1Ra1S, such as a native sequence hIL-1Ra1S.

In another embodiment, the invention provides a method for treating acute lung injury comprising administering to a mammal, such as human, in need of such treatment an effective amount of an IL-11p, such as a native sequence IL-11p.

In another embodiment, the invention provides a method for treating acute lung injury comprising administering to a mammal, such as human, in need of such treatment an effective amount of an IL-11p selected from the group consisting of hIL-1Ra1, hIL-1Ra1L, hIL-1Ra1V, hIL-1Ra1S, hIL-1Ra2, hIL-1Ra3, and mLIL-1Ra3.

In another embodiment, the invention provides a method for treating acute lung injury comprising administering to a mammal, such as human, in need of such treatment an effective amount of an hIL-11p, such as a native sequence hIL-11p, e.g. native sequence hIL-1Ra1 or hIL-1Ra3.

In another embodiment, the invention provides a method for treating acute lung injury comprising administering to a mammal, such as human, in need of such treatment an effective amount of an hIL-1Ra1L, such as a native sequence hIL-1Ra1L, or an effective amount of an hIL-1Ra1V, such as a native sequence hIL-1Ra1V, or an effective amount of an hIL-1Ra1S, such as a native sequence hIL-1Ra1S.

In another embodiment, the invention provides a method for treating adult respiratory distress syndrome comprising administering to a mammal, such as human, in need of such treatment an effective amount of an IL-11p, such as a native sequence IL-11p.

In another embodiment, the invention provides a method for treating adult respiratory distress syndrome comprising administering to a mammal, such as human, in need of such treatment an effective amount of an IL-1lp selected from the group consisting of hIL-1Ra1, hIL-1Ra1L, hIL-1Ra1V, hIL-1Ra1S, hIL-1Ra2, hIL-1Ra3, and mL-1Ra3.

5 In another embodiment, the invention provides a method for treating adult respiratory distress syndrome comprising administering to a mammal, such as human, in need of such treatment an effective amount of an hIL-1lp, such as a native sequence hIL-1lp, e.g. native sequence hIL-1Ra1 or hIL-1Ra3.

10 In another embodiment, the invention provides a method for treating adult respiratory distress syndrome comprising administering to a mammal, such as human, in need of such treatment an effective amount of an hIL-1Ra1L, such as a native sequence hIL-1Ra1L, or an effective amount of an hIL-1Ra1V, such as a native sequence hIL-1Ra1V, or an effective amount of an hIL-1Ra1S, such as a native sequence hIL-1Ra1S.

15 In another embodiment, the invention provides a method for treating idiopathic pulmonary fibrosis comprising administering to a mammal, such as human, in need of such treatment an effective amount of an IL-1lp, such as a native sequence IL-1lp.

20 In another embodiment, the invention provides a method for treating idiopathic pulmonary fibrosis comprising administering to a mammal, such as human, in need of such treatment an effective amount of an IL-1lp selected from the group consisting of hIL-1Ra1, hIL-1Ra1L, hIL-1Ra1V, hIL-1Ra1S, hIL-1Ra2, hIL-1Ra3, and mL-1Ra3.

25 In another embodiment, the invention provides a method for treating idiopathic pulmonary fibrosis comprising administering to a mammal, such as human, in need of such treatment an effective amount of an hIL-1lp, such as a native sequence hIL-1lp, e.g. native sequence hIL-1Ra1 or hIL-1Ra3.

30 In another embodiment, the invention provides a method for treating idiopathic pulmonary fibrosis comprising administering to a mammal, such as human, in need of such treatment an effective amount of an hIL-1Ra1L, such as a native sequence hIL-1Ra1L, or an effective amount of an hIL-1Ra1V, such as a native sequence hIL-1Ra1V, or an effective amount of an hIL-1Ra1S, such as a native sequence hIL-1Ra1S.

35 In another embodiment, the invention provides a method for treating an ischemic reperfusion disease, such as surgical tissue reperfusion injury, stroke, myocardial ischemia, or acute myocardial infarction, comprising administering to a mammal, such as human, in need of such treatment an effective amount of an IL-1lp, such as a native sequence IL-1lp.

40 In another embodiment, the invention provides a method for treating an ischemic reperfusion disease, such as surgical tissue reperfusion injury, stroke, myocardial ischemia, or acute myocardial infarction, comprising administering to a mammal, such as human, in need of such treatment an effective amount of an IL-1lp selected from the group consisting of hIL-1Ra1, hIL-1Ra1L, hIL-1Ra1V, hIL-1Ra1S, hIL-1Ra2, hIL-1Ra3, and mL-1Ra3.

In another embodiment, the invention provides a method for treating an ischemic reperfusion disease, such as surgical tissue reperfusion injury, stroke, myocardial ischemia, or acute myocardial infarction, comprising administering to a mammal, such as human, in need

of such treatment an effective amount of an hIL-1lp, such as a native sequence hIL-1lp, e.g. native sequence hIL-1Ra1 or hIL-1Ra3.

In another embodiment, the invention provides a method for treating an ischemic reperfusion disease, such as surgical tissue reperfusion injury, stroke, myocardial ischemia, or acute myocardial infarction, comprising administering to a mammal, such as human, in need of such treatment an effective amount of an hIL-1Ra1L, such as a native sequence hIL-1Ra1L, or an effective amount of an hIL-1Ra1V, such as a native sequence hIL-1Ra1V, or an effective amount of an hIL-1Ra1S, such as a native sequence hIL-1Ra1S.

In another embodiment, the invention provides a method for treating psoriasis comprising administering to a mammal, such as human, in need of such treatment an effective amount of an IL-1lp, such as a native sequence IL-1lp.

In another embodiment, the invention provides a method for treating psoriasis comprising administering to a mammal, such as human, in need of such treatment an effective amount of an IL-1lp selected from the group consisting of hIL-1Ra1, hIL-1Ra1L, hIL-1Ra1V, hIL-1Ra1S, hIL-1Ra2, hIL-1Ra3, and mL-1Ra3.

In another embodiment, the invention provides a method for treating psoriasis comprising administering to a mammal, such as human, in need of such treatment an effective amount of an hIL-1lp, such as a native sequence hIL-1lp, e.g. native sequence hIL-1Ra1 or hIL-1Ra3.

In another embodiment, the invention provides a method for treating psoriasis comprising administering to a mammal, such as human, in need of such treatment an effective amount of an hIL-1Ra1L, such as a native sequence hIL-1Ra1L, or an effective amount of an hIL-1Ra1V, such as a native sequence hIL-1Ra1V, or an effective amount of an hIL-1Ra1S, such as a native sequence hIL-1Ra1S.

In another embodiment, the invention provides a method for treating graft-versus-host disease (GVHD) comprising administering to a mammal, such as human, in need of such treatment an effective amount of an IL-1lp, such as a native sequence IL-1lp.

In another embodiment, the invention provides a method for treating graft-versus-host disease (GVHD) comprising administering to a mammal, such as human, in need of such treatment an effective amount of an IL-1lp selected from the group consisting of hIL-1Ra1, hIL-1Ra1L, hIL-1Ra1V, hIL-1Ra1S, hIL-1Ra2, hIL-1Ra3, and mL-1Ra3.

In another embodiment, the invention provides a method for treating graft-versus-host disease (GVHD) comprising administering to a mammal, such as human, in need of such treatment an effective amount of an hIL-1lp, such as a native sequence hIL-1lp, e.g. native sequence hIL-1Ra1 or hIL-1Ra3.

In another embodiment, the invention provides a method for treating graft-versus-host disease (GVHD) comprising administering to a mammal, such as human, in need of such treatment an effective amount of an hIL-1Ra1L, such as a native sequence hIL-1Ra1L, or an effective amount of an hIL-1Ra1V, such as a native sequence hIL-1Ra1V, or an effective amount of an hIL-1Ra1S, such as a native sequence hIL-1Ra1S.

In another embodiment, the invention provides a method for treating an inflammatory bowel disease such as ulcerative colitis, comprising administering to a mammal, such as human, in need of such treatment an effective amount of an IL-1lp, such as a native sequence IL-1lp.

In another embodiment, the invention provides a method for treating an inflammatory bowel disease such as ulcerative colitis, comprising administering to a mammal, such as human, in need of such treatment an effective amount of an IL-1lp selected from the group consisting of hIL-1Ra1, hIL-1Ra1L, hIL-1Ra1V, hIL-1Ra1S, hIL-1Ra2, hIL-1Ra3, and mL-1Ra3.

In another embodiment, the invention provides a method for treating an inflammatory bowel disease such as ulcerative colitis, comprising administering to a mammal, such as human, in need of such treatment an effective amount of an hIL-1lp, such as a native sequence hIL-1lp, e.g. native sequence hIL-1Ra1 or hIL-1Ra3.

In another embodiment, the invention provides a method for treating an inflammatory bowel disease such as ulcerative colitis, comprising administering to a mammal, such as human, in need of such treatment an effective amount of an hIL-1Ra1L, such as a native sequence hIL-1Ra1L, or an effective amount of an hIL-1Ra1V, such as a native sequence hIL-1Ra1V, or an effective amount of an hIL-1Ra1S, such as a native sequence hIL-1Ra1S.

D. Anti-IL-1lp Antibodies

The present invention further provides anti-IL-1lp antibodies. Exemplary antibodies include polyclonal, monoclonal, humanized, bispecific, and heteroconjugate antibodies.

1. Polyclonal Antibodies

The anti-IL-1lp antibodies may comprise polyclonal antibodies. Methods of preparing polyclonal antibodies are known to the skilled artisan. Polyclonal antibodies can be raised in a mammal, for example, by one or more injections of an immunizing agent and, if desired, an adjuvant. Typically, the immunizing agent and/or adjuvant will be injected in the mammal by multiple subcutaneous or intraperitoneal injections. The immunizing agent may include the IL-1lp polypeptide or a fusion protein thereof. It may be useful to conjugate the immunizing agent to a protein known to be immunogenic in the mammal being immunized. Examples of such immunogenic proteins include but are not limited to keyhole limpet hemocyanin, serum albumin, bovine thyroglobulin, and soybean trypsin inhibitor. Examples of adjuvants which may be employed include Freund's complete adjuvant and MPL-TDM adjuvant (monophosphoryl Lipid A, synthetic trehalose dicorynomycolate). The immunization protocol may be selected by one skilled in the art without undue experimentation.

2. Monoclonal Antibodies

The anti-IL-1lp antibodies may, alternatively, be monoclonal antibodies. Monoclonal antibodies may be prepared using hybridoma methods, such as those described by Kohler and Milstein, Nature, 256:495 (1975). In a hybridoma method, a mouse, hamster, or other appropriate host animal, is typically immunized with an immunizing agent to elicit lymphocytes that produce or are capable of producing antibodies that will specifically bind to the immunizing agent. Alternatively, the lymphocytes may be immunized *in vitro*.

The immunizing agent will typically include the IL-1lp polypeptide or a fusion protein thereof. Generally, either peripheral blood lymphocytes ("PBLs") are used if cells of human origin

are desired, or spleen cells or lymph node cells are used if non-human mammalian sources are desired. The lymphocytes are then fused with an immortalized cell line using a suitable fusing agent, such as polyethylene glycol, to form a hybridoma cell [Goding, *Monoclonal Antibodies: Principles and Practice*, Academic Press, (1986) pp. 59-103]. Immortalized cell lines are usually transformed mammalian cells, particularly myeloma cells of rodent, bovine and human origin. Usually, rat or mouse myeloma cell lines are employed. The hybridoma cells may be cultured in a suitable culture medium that preferably contains one or more substances that inhibit the growth or survival of the unfused, immortalized cells. For example, if the parental cells lack the enzyme hypoxanthine guanine phosphoribosyl transferase (HGPRT or HPRT), the culture medium for the hybridomas typically will include hypoxanthine, aminopterin, and thymidine ("HAT medium"), which substances prevent the growth of HGPRT-deficient cells.

Preferred immortalized cell lines are those that fuse efficiently, support stable high level expression of antibody by the selected antibody-producing cells, and are sensitive to a medium such as HAT medium. More preferred immortalized cell lines are murine myeloma lines, which can be obtained, for instance, from the Salk Institute Cell Distribution Center, San Diego, California and the American Type Culture Collection, Rockville, Maryland. Human myeloma and mouse-human heteromyeloma cell lines also have been described for the production of human monoclonal antibodies [Kozbor, *J. Immunol.*, **133**:3001 (1984); Brodeur *et al.*, *Monoclonal Antibody Production Techniques and Applications*, Marcel Dekker, Inc., New York, (1987) pp. 51-63].

The culture medium in which the hybridoma cells are cultured can then be assayed for the presence of monoclonal antibodies directed against IL-11p. Preferably, the binding specificity of monoclonal antibodies produced by the hybridoma cells is determined by immunoprecipitation or by an *in vitro* binding assay, such as radioimmunoassay (RIA) or enzyme-linked immunosorbant assay (ELISA). Such techniques and assays are known in the art. The binding affinity of the monoclonal antibody can, for example, be determined by the Scatchard analysis of Munson and Pollard, *Anal. Biochem.*, **107**:220 (1980).

After the desired hybridoma cells are identified, the clones may be subcloned by limiting dilution procedures and grown by standard methods [Goding, *supra*]. Suitable culture media for this purpose include, for example, Dulbecco's Modified Eagle's Medium and RPMI-1640 medium. Alternatively, the hybridoma cells may be grown *in vivo* as ascites in a mammal.

The monoclonal antibodies secreted by the subclones may be isolated or purified from the culture medium or ascites fluid by conventional immunoglobulin purification procedures such as, for example, protein A-Sepharose, hydroxylapatite chromatography, gel electrophoresis, dialysis, or affinity chromatography.

The monoclonal antibodies may also be made by recombinant DNA methods, such as those described in U.S. Patent No. 4,816,567. DNA encoding the monoclonal antibodies of the invention can be readily isolated and sequenced using conventional procedures (e.g., by using oligonucleotide probes that are capable of binding specifically to genes encoding the heavy and light chains of murine antibodies). The hybridoma cells of the invention serve as a preferred

source of such DNA. Once isolated, the DNA may be placed into expression vectors, which are then transfected into host cells such as simian COS cells, Chinese hamster ovary (CHO) cells, or myeloma cells that do not otherwise produce immunoglobulin protein, to obtain the synthesis of monoclonal antibodies in the recombinant host cells. The DNA also may be modified, for example, by substituting the coding sequence for human heavy and light chain constant domains in place of the homologous murine sequences [U.S. Patent No. 4,816,567; Morrison et al., supra] or by covalently joining to the immunoglobulin coding sequence all or part of the coding sequence for a non-immunoglobulin polypeptide. Such a non-immunoglobulin polypeptide can be substituted for the constant domains of an antibody of the invention, or can be substituted for the variable domains of one antigen-combining site of an antibody of the invention to create a chimeric bivalent antibody.

The antibodies may be monovalent antibodies. Methods for preparing monovalent antibodies are well known in the art. For example, one method involves recombinant expression of immunoglobulin light chain and modified heavy chain. The heavy chain is truncated generally at any point in the Fc region so as to prevent heavy chain crosslinking. Alternatively, the relevant cysteine residues are substituted with another amino acid residue or are deleted so as to prevent crosslinking.

In vitro methods are also suitable for preparing monovalent antibodies. Digestion of antibodies to produce fragments thereof, particularly, Fab fragments, can be accomplished using routine techniques known in the art.

3. Human and Humanized Antibodies

The anti-IL-11p antibodies of the invention may further comprise humanized antibodies or human antibodies. Humanized forms of non-human (e.g., murine) antibodies are chimeric immunoglobulins, immunoglobulin chains or fragments thereof (such as Fv, Fab, Fab', F(ab')₂ or other antigen-binding subsequences of antibodies) which contain minimal sequence derived from non-human immunoglobulin. Humanized antibodies include human immunoglobulins (recipient antibody) in which residues from a complementary determining region (CDR) of the recipient are replaced by residues from a CDR of a non-human species (donor antibody) such as mouse, rat or rabbit having the desired specificity, affinity and capacity. In some instances, Fv framework residues of the human immunoglobulin are replaced by corresponding non-human residues. Humanized antibodies may also comprise residues which are found neither in the recipient antibody nor in the imported CDR or framework sequences. In general, the humanized antibody will comprise substantially all of at least one, and typically two, variable domains, in which all or substantially all of the CDR regions correspond to those of a non-human immunoglobulin and all or substantially all of the FR regions are those of a human immunoglobulin consensus sequence. The humanized antibody optimally also will comprise at least a portion of an immunoglobulin constant region (Fc), typically that of a human immunoglobulin [Jones *et al.*, *Nature*, 321: 522-525 (1986); Riechmann *et al.*, *Nature*, 332: 323-329 (1988); and Presta, *Curr. Op. Struct. Biol.*, 2: 593-596 (1992)].

Methods for humanizing non-human antibodies are well known in the art. Generally, a humanized antibody has one or more amino acid residues introduced into it from a source which

is non-human. These non-human amino acid residues are often referred to as "import" residues, which are typically taken from an "import" variable domain. Humanization can be essentially performed following the method of Winter and co-workers [Jones *et al.*, *Nature*, 321: 522-525 (1986); Riechmann *et al.*, *Nature*, 332: 323-327 (1988); Verhoeven *et al.*, *Science*, 239: 1534-1536 (1988)], by substituting rodent CDRs or CDR sequences for the corresponding sequences of a human antibody. Accordingly, such "humanized" antibodies are chimeric antibodies (U.S. Patent No. 4,816,567), wherein substantially less than an intact human variable domain has been substituted by the corresponding sequence from a non-human species. In practice, humanized antibodies are typically human antibodies in which some CDR residues and possibly some FR residues are substituted by residues from analogous sites in rodent antibodies.

Human antibodies can also be produced using various techniques known in the art, including phage display libraries [Hoogenboom and Winter, *J. Mol. Biol.*, 227: 381 (1991); Marks *et al.*, *J. Mol. Biol.*, 222: 581 (1991)]. The techniques of Cole *et al.* and Boerner *et al.* are also available for the preparation of human monoclonal antibodies (Cole *et al.*, *Monoclonal Antibodies and Cancer Therapy*, Alan R. Liss, p. 77 (1985) and Boerner *et al.*, *J. Immunol.*, 147(1): 86-95 (1991)]. Similarly, human antibodies can be made by introducing of human immunoglobulin loci into transgenic animals, *e.g.*, mice in which the endogenous immunoglobulin genes have been partially or completely inactivated. Upon challenge, human antibody production is observed, which closely resembles that seen in humans in all respects, including gene rearrangement, assembly, and antibody repertoire. This approach is described, for example, in U.S. Patent Nos. 5,545,807; 5,545,806; 5,569,825; 5,625,126; 5,633,425; 5,661,016, and in the following scientific publications: Marks *et al.*, *Bio/Technology* 10, 779-783 (1992); Lonberg *et al.*, *Nature* 368: 856-859 (1994); Morrison, *Nature* 368, 812-13 (1994); Fishwild *et al.*, *Nature Biotechnology* 14, 845-51 (1996); Neuberger, *Nature Biotechnology* 14, 826 (1996); Lonberg and Huszar, *Intern. Rev. Immunol.* 13 65-93 (1995).

4. Bispecific Antibodies

Bispecific antibodies are monoclonal, preferably human or humanized, antibodies that have binding specificities for at least two different antigens. In the present case, one of the binding specificities is for the IL-1p, the other one is for any other antigen, and preferably for a cell-surface protein or receptor or receptor subunit.

Methods for making bispecific antibodies are known in the art. Traditionally, the recombinant production of bispecific antibodies is based on the co-expression of two immunoglobulin heavy-chain/light-chain pairs, where the two heavy chains have different specificities [Milstein and Cuello, *Nature*, 305:537-539 (1983)]. Because of the random assortment of immunoglobulin heavy and light chains, these hybridomas (quadromas) produce a potential mixture of ten different antibody molecules, of which only one has the correct bispecific structure. The purification of the correct molecule is usually accomplished by affinity chromatography steps. Similar procedures are disclosed in WO 93/08829, published 13 May 1993, and in Traunecker *et al.*, *EMBO J.*, 10: 3655-3659 (1991).

Antibody variable domains with the desired binding specificities (antibody-antigen combining sites) can be fused to immunoglobulin constant domain sequences. The fusion preferably is with an immunoglobulin heavy-chain constant domain, comprising at least part of the hinge, CH2, and CH3 regions. It is preferred to have the first heavy-chain constant region (CH1) containing the site necessary for light-chain binding present in at least one of the fusions. DNAs encoding the immunoglobulin heavy-chain fusions and, if desired, the immunoglobulin light chain, are inserted into separate expression vectors, and are co-transfected into a suitable host organism. For further details of generating bispecific antibodies see, for example, Suresh et al., Methods in Enzymology, 121:210 (1986).

5. Heteroconjugate Antibodies

Heteroconjugate antibodies are also within the scope of the present invention. Heteroconjugate antibodies are composed of two covalently joined antibodies. Such antibodies have, for example, been proposed to target immune system cells to unwanted cells [U.S. Patent No. 4,676,980], and for treatment of HIV infection [WO 91/00360; WO 92/200373; EP 03089].

It is contemplated that the antibodies may be prepared *in vitro* using known methods in synthetic protein chemistry, including those involving crosslinking agents. For example, immunotoxins may be constructed using a disulfide exchange reaction or by forming a thioether bond. Examples of suitable reagents for this purpose include iminothiolate and methyl-4-mercaptobutyrimide and those disclosed, for example, in U.S. Patent No. 4,676,980.

E. Uses for anti-IL-1p Antibodies

The anti-IL-1p antibodies of the invention have various utilities. For example, anti-IL-1p antibodies may be used in diagnostic assays for IL-1p, *e.g.*, detecting its expression in specific cells, tissues, or serum. Various diagnostic assay techniques known in the art may be used, such as competitive binding assays, direct or indirect sandwich assays and immunoprecipitation assays conducted in either heterogeneous or homogeneous phases [Zola, *Monoclonal Antibodies: A Manual of Techniques*, CRC Press, Inc. (1987) pp. 147-158]. The antibodies used in the diagnostic assays can be labeled with a detectable moiety. The detectable moiety should be capable of producing, either directly or indirectly, a detectable signal. For example, the detectable moiety may be a radioisotope, such as ^3H , ^{14}C , ^{32}P , ^{35}S , or ^{125}I , a fluorescent or chemiluminescent compound, such as fluorescein isothiocyanate, rhodamine, or luciferin, or an enzyme, such as alkaline phosphatase, beta-galactosidase or horseradish peroxidase. Any method known in the art for conjugating the antibody to the detectable moiety may be employed, including those methods described by Hunter *et al.*, Nature, 144: 945 (1962); David *et al.*, Biochemistry, 13: 1014 (1974); Pain *et al.*, J. Immunol. Meth., 40: 219 (1981); and Nygren, J. Histochem. and Cytochem., 30: 407 (1982).

Anti-IL-1p antibodies also are useful for the affinity purification of IL-1p from recombinant cell culture or natural sources. In this process, the antibodies against IL-1p are immobilized on a suitable support, such as Sephadex resin or filter paper, using methods well known in the art. The immobilized antibody then is contacted with a sample containing the IL-1p to be purified, and thereafter the support is washed with a suitable solvent that will remove

substantially all the material in the sample except the IL-1lp, which is bound to the immobilized antibody. Finally, the support is washed with another suitable solvent that will release the IL-1lp from the antibody.

In addition, anti-IL-1lp antibodies are useful as therapeutic agents for targeting of native IL-1lp in IL-1lp-mediated disease conditions, e.g. disease states characterized by pathologic IL-1 or IL-18 agonist or agonist-like activity of the native IL-1lp. In the treatment and prevention of a native IL-1lp-mediated disorder with the anti-IL-1lp antibody of the invention, the antibody composition will be formulated, dosed, and administered in a fashion consistent with good medical practice. Factors for consideration in this context include the particular disorder being treated, the particular mammal being treated, the clinical condition of the individual patient, the cause of the disorder, the site of delivery of the antibody, the particular type of antibody, the method of administration, the scheduling of administration, and other factors known to medical practitioners. The "effective amount" or "therapeutically effective amount" of antibody to be administered will be governed by such considerations, and is the minimum amount necessary to prevent, ameliorate, or treat the native IL-1lp-mediated disorder, including treating inflammatory diseases and reducing inflammatory responses. Such amount is preferably below the amount that is toxic to the host or renders the host significantly more susceptible to infections.

As a general proposition, the initial pharmaceutically effective amount of the antibody or antibody fragment administered parenterally per dose will be in the range of about 0.1 to 50 mg/kg of patient body weight per day, with the typical initial range of antibody used being 0.3 to 20 mg/kg/day, more preferably 0.3 to 15 mg/kg/day.

In one embodiment, using systemic administration, the initial pharmaceutically effective amount will be in the range of about 2 to 5 mg/kg/day.

For methods of the invention using administration by inhalation, the initial pharmaceutically effective amount will be in the range of about 1 microgram (μ g)/kg/day to 100 mg/kg/day.

In one embodiment, the invention provides a method for treating an IL-1lp-mediated inflammatory disorder comprising administering to a mammal, such as human, in need of such treatment an effective amount of an anti-IL-1lp antibody.

In another embodiment, the invention provides a method for treating an hIL-1lp-mediated asthmatic disorder comprising administering to a human in need of such treatment an effective amount of an anti-hIL-1lp antibody.

In another embodiment, the invention provides a method for treating an hIL-1Ra1-mediated asthmatic disorder comprising administering to a human in need of such treatment an effective amount of an anti-hIL-1Ra1 antibody.

In another embodiment, the invention provides a method for treating an IL-1lp-mediated rheumatoid arthritic disorder comprising administering to a mammal, such as human, in need of such treatment an effective amount of an anti-IL-1lp antibody.

In another embodiment, the invention provides a method for treating an hIL-11p-mediated rheumatoid arthritic disorder comprising administering to a human in need of such treatment an effective amount of an anti-hIL-11p antibody.

5 In another embodiment, the invention provides a method for treating an hIL-1Ra1-mediated rheumatoid arthritic disorder comprising administering to a human in need of such treatment an effective amount of an anti-hIL-1Ra1 antibody.

In another embodiment, the invention provides a method for treating an IL-11p-mediated osteoarthritic disorder comprising administering to a mammal, such as human, in need of such treatment an effective amount of an anti-IL-11p antibody.

10 In another embodiment, the invention provides a method for treating an hIL-11p-mediated osteoarthritic disorder comprising administering to a human in need of such treatment an effective amount of an anti-hIL-11p antibody.

15 In another embodiment, the invention provides a method for treating an hIL-1Ra1-mediated osteoarthritic disorder comprising administering to a human in need of such treatment an effective amount of an anti-hIL-1Ra1 antibody.

In another embodiment, the invention provides a method for treating an IL-11p-mediated septic disorder comprising administering to a mammal, such as human, in need of such treatment an effective amount of an anti-IL-11p antibody.

20 In another embodiment, the invention provides a method for treating an hIL-11p-mediated septic disorder comprising administering to a human in need of such treatment an effective amount of an anti-hIL-11p antibody.

In another embodiment, the invention provides a method for treating an hIL-1Ra1-mediated septic disorder comprising administering to a human in need of such treatment an effective amount of an anti-hIL-1Ra1 antibody.

25 In another embodiment, the invention provides a method for treating IL-11p-mediated acute lung injury comprising administering to a mammal, such as human, in need of such treatment an effective amount of an anti-IL-11p antibody.

30 In another embodiment, the invention provides a method for treating hIL-11p-mediated acute lung injury comprising administering to a human in need of such treatment an effective amount of an anti-hIL-11p antibody.

In another embodiment, the invention provides a method for treating hIL-1Ra1-mediated acute lung injury comprising administering to a human in need of such treatment an effective amount of an anti-hIL-1Ra1 antibody.

35 In another embodiment, the invention provides a method for treating IL-11p-mediated adult respiratory distress syndrome comprising administering to a mammal, such as human, in need of such treatment an effective amount of an anti-IL-11p antibody.

In another embodiment, the invention provides a method for treating hIL-11p-mediated adult respiratory distress syndrome comprising administering to a human in need of such treatment an effective amount of an anti-hIL-11p antibody.

In another embodiment, the invention provides a method for treating hIL-1Ra1-mediated adult respiratory distress syndrome comprising administering to a human in need of such treatment an effective amount of an anti-hIL-1Ra1 antibody.

In another embodiment, the invention provides a method for treating IL-1lp-mediated idiopathic pulmonary fibrosis comprising administering to a mammal, such as human, in need of such treatment an effective amount of an anti-IL-1lp antibody.

In another embodiment, the invention provides a method for treating hIL-1lp-mediated idiopathic pulmonary fibrosis comprising administering to a human in need of such treatment an effective amount of an anti-hIL-1lp antibody.

In another embodiment, the invention provides a method for treating hIL-1Ra1-mediated idiopathic pulmonary fibrosis comprising administering to a human in need of such treatment an effective amount of an anti-hIL-1Ra1 antibody.

In another embodiment, the invention provides a method for treating an IL-1lp-mediated ischemic reperfusion disease, such as surgical tissue reperfusion injury, stroke, myocardial ischemia, or acute myocardial infarction, comprising administering to a mammal, such as human, in need of such treatment an effective amount of an anti-IL-1lp antibody.

In another embodiment, the invention provides a method for treating an hIL-1lp-mediated ischemic reperfusion disease, such as surgical tissue reperfusion injury, stroke, myocardial ischemia, or acute myocardial infarction, comprising administering to a human in need of such treatment an effective amount of an anti-hIL-1lp antibody.

In another embodiment, the invention provides a method for treating an hIL-1Ra1-mediated ischemic reperfusion disease, such as surgical tissue reperfusion injury, stroke, myocardial ischemia, or acute myocardial infarction, comprising administering to a human in need of such treatment an effective amount of an anti-hIL-1Ra1 antibody.

In another embodiment, the invention provides a method for treating an IL-1lp-mediated psoriatic disorder comprising administering to a mammal, such as human, in need of such treatment an effective amount of an anti-IL-1lp antibody.

In another embodiment, the invention provides a method for treating an hIL-1lp-mediated psoriatic disorder comprising administering to a human in need of such treatment an effective amount of an anti-hIL-1lp antibody.

In another embodiment, the invention provides a method for treating an hIL-1Ra1-mediated psoriatic disorder comprising administering to a human in need of such treatment an effective amount of an anti-hIL-1Ra1 antibody.

In another embodiment, the invention provides a method for treating an IL-1lp-mediated graft-versus-host disease (GVHD) comprising administering to a mammal, such as human, in need of such treatment an effective amount of an anti-IL-1lp antibody.

In another embodiment, the invention provides a method for treating an hIL-1lp-mediated graft-versus-host disease (GVHD) comprising administering to a human in need of such treatment an effective amount of an anti-hIL-1lp antibody.

M 27.07.00

In another embodiment, the invention provides a method for treating an hIL-1Ra1-mediated graft-versus-host disease (GVHD) comprising administering to a human in need of such treatment an effective amount of an anti-hIL-1Ra1 antibody.

In another embodiment, the invention provides a method for treating an IL-1Ip-mediated inflammatory bowel disease such as ulcerative colitis, comprising administering to a mammal, such as human, in need of such treatment an effective amount of an anti-IL-1Ip antibody.

In another embodiment, the invention provides a method for treating an hIL-1Ip-mediated inflammatory bowel disease such as ulcerative colitis, comprising administering to a human in need of such treatment an effective amount of an anti-hIL-1Ip antibody.

In another embodiment, the invention provides a method for treating an hIL-1Ra1-mediated inflammatory bowel disease such as ulcerative colitis, comprising administering to a human in need of such treatment an effective amount of an anti-hIL-1Ra1 antibody.

The following examples are offered for illustrative purposes only, and are not intended to limit the scope of the present invention in any way.

All patent and literature references cited in the present specification are hereby incorporated by reference in their entirety.

EXAMPLES

Commercially available reagents referred to in the examples were used according to manufacturer's instructions unless otherwise indicated. The source of those cells identified in the following examples, and throughout the specification, by ATCC accession numbers is the American Type Culture Collection, Manassas, VA.

EXAMPLE 1

Isolation of DNA encoding hIL-1Ra1 and mIL-1Ra3

A public expressed sequence tag (EST) DNA database (Genbank) was searched with human interleukin-1 receptor antagonist (hIL-1Ra) sequence, also known as secretory human interleukin-1 receptor antagonist ("sIL-1Ra") sequence, and a human EST designated AI014548 (which corresponds to nucleotides 145-629 of Figure 4, SEQ ID NO:8), and a murine EST designated W08205 (Figure 10, SEQ ID NO:17), were identified, which showed homology with the known protein hIL-1Ra (sIL-1Ra).

EST clones AI014548 and W08205 were purchased from Research Genetics (Huntsville, AL) and the cDNA inserts were obtained and sequenced in their entirety.

The entire nucleotide sequence of the clone AI014548, designated DNA85066, is shown in Figure 1 (SEQ ID NO:1). Clone DNA85066 contains a single open reading frame that is interrupted by an apparent intronic sequence. The intron is bounded by splice junctions at nucleotide positions 181 to 186 (splice donor site) and nucleotide positions 430 to 432 (splice acceptor site) (Fig. 1; SEQ ID NO:1).

A virtual processed nucleotide sequence (Fig. 3; SEQ ID NO:6), designated DNA94618, was derived by removing the apparent intronic sequence from clone DNA85066. Clone DNA94618 contains a single open reading frame with an apparent translational initiation site at nucleotide positions 103-105, and a stop codon at nucleotide positions 682-684 (Fig. 3; SEQ ID NO:6). The predicted polypeptide precursor (hIL-1Ra1) (Fig. 3; SEQ ID NO:7) is 193 amino acids long. The putative signal sequence extends from amino acid positions 1 to 14. A putative

M 27.07.00

cAMP- and cGMP-dependent protein kinase phosphorylation site is located at amino acid positions 33-36. Putative N-myristoylation sites are located at amino acid positions 50-55 and 87-92.

Clone DNA85066 (designated as DNA85066-2534) has been deposited with ATCC and was assigned ATCC deposit no. 203588. The full-length hIL-1Ra1 protein shown in Figure 3 (SEQ ID NO:7) has an estimated molecular weight of about 21,822 daltons and a pI of about 8.9.

Based on a sequence alignment analysis of the full-length sequence (SEQ ID NO:7), hIL-1Ra1 shows significant amino acid sequence identity to hIL-1Ra (sIL-1Ra) and hIL-1Ra β proteins.

The entire nucleotide sequence of the clone W08205, designated DNA92505, is shown in Figure 9 (SEQ ID NO:15). Clone DNA92505 contains a single open reading frame with an apparent translational initiation site at nucleotide positions 145-147, and a stop codon at nucleotide positions 610-612 (Fig. 9; SEQ ID NO:15). The predicted polypeptide precursor (mIL-1Ra3) (Fig. 9; SEQ ID NO:16) is 155 amino acids long. The putative signal sequence extends from amino acid positions 1-33. Putative N-myristoylation sites are located at amino acid positions 29-34, 60-65, 63-68, 91-96 and 106-111. An interleukin-1-like sequence is located at amino acid positions 111-131.

Clone DNA92505 (designated as DNA92505-2534) was deposited with ATCC and was assigned ATCC deposit no. 203590. The full length mIL-1Ra3 protein shown in Figure 9 (SEQ ID NO:16) has an estimated molecular weight of about 17,134 daltons and a pI of about 4.8.

Based on a sequence alignment analysis of the full-length sequence (SEQ ID NO:16), mIL-1Ra3 shows significant amino acid sequence identity to mIL-1Ra, hIL-1Ra, hIL-1Ra (sIL-1Ra) and hIL-1Ra β proteins.

EXAMPLE 2

Isolation of DNA encoding hIL-1ra2 and hIL-1Ra3

A expressed sequence tag (EST) DNA database (LIFESEQ®, Incyte Pharmaceuticals, Palo Alto, CA) was searched with human interleukin-1 receptor antagonist (hIL-1Ra) sequence, also known as secretory human interleukin-1 receptor antagonist ("sIL-1Ra") sequence, and the ESTs, designated 1433156 (Figure 5, SEQ ID NO:11) and 5120028 (Figure 8, SEQ ID NO:14), were identified, which showed homology with the hIL-1Ra known protein.

EST clones 1433156 and 5120028 were purchased from Incyte Pharmaceuticals (Palo Alto, CA) and the cDNA inserts were obtained and sequenced in their entireties.

The entire nucleotide sequence of the clone 1433156, designated DNA92929, is shown in Figure 5 (SEQ ID NO:9). Clone DNA92929 contains a single open reading frame with an apparent translational initiation site at nucleotide positions 96-98, and a stop codon at nucleotide positions 498-500 (Fig. 5; SEQ ID NO:9). The predicted polypeptide precursor (hIL-1Ra2) (Fig. 5; SEQ ID NO:10) is 134 amino acids long. A putative signal sequence extends from amino acid positions 1-26.

Clone DNA92929 (designated as DNA92929-2534) was deposited with ATCC and was assigned ATCC deposit no. 203586. The full-length hIL-1Ra2 protein shown in Figure 5 (SEQ ID NO:10) has an estimated molecular weight of about 14,927 daltons and a pI of about 4.8.

Based on a sequence alignment analysis of the full-length sequence (SEQ ID NO:10), hIL-1Ra2 shows significant amino acid sequence identity to hIL-1Ra β protein. hIL-1Ra2 is believed to be a splice variant of hIL-1Ra β .

The entire nucleotide sequence of the clone 5120028, designated DNA96787, is shown in Figure 7 (SEQ ID NO:12). Clone DNA96787 contains a single open reading frame with an apparent translational initiation site at nucleotide positions 1-3, and a stop codon at nucleotide positions 466-468 (Fig. 7; SEQ ID NO:12). The predicted polypeptide precursor (hIL-1Ra3) (Fig. 7; SEQ ID NO:13) is 155 amino acids long. A putative signal sequence extends from amino acid positions 1-33. Putative N-myristoylation sites are located at amino acid positions 29-34, 60-65, 63-68, 73-78, 91-96 and 106-111. An interleukin-1-like sequence is located at amino acid positions 111-131.

It is believed that the predicted 155 amino acid polypeptide of hIL-1Ra3 behaves as a mature sequence (without a presequence that is removed in post-translational processing) in certain animal cells. It is also believed that other animal cells recognize and remove one or more signal peptide(s) extending from amino acid positions 1 to about 33. As shown in Example 14 below, transiently transfected CHO host cells secrete a form of hIL-1Ra3 that only lacks the N-terminal methionine in the sequence of Figure 7 (SEQ ID NO:13).

Clone DNA96787 (designated as DNA96787-2534) was deposited with ATCC and was assigned ATCC deposit no. 203589. The full length hIL-1Ra3 protein shown in Figure 7 (SEQ ID NO:13) has an estimated molecular weight of about 16,961 daltons and a pI of about 4.9.

Based on a sequence alignment analysis of the full-length sequence (SEQ ID NO:13), hIL-1Ra3 shows significant amino acid sequence identity to hIL-1Ra and hIL-1Ra (sIL-1Ra) proteins.

EXAMPLE 3

Northern Blot Analysis

Expression of hIL-1Ra3 mRNA in human tissues and mIL-1Ra3 mRNA in mouse tissues was examined by Northern blot analysis. Human and mouse multiple tissue northern (RNA) blots and mouse embryo blots were purchased from Clontech and probed with corresponding cDNA according to the manufacturer's instructions.

As shown in Fig. 11, hIL-1Ra3 mRNA (2.7 kb) were detected only in human placenta and mIL-1Ra3 mRNA transcripts (1.4 kb and 2.5 kb) were detected only in the day-17 mouse embryo.

EXAMPLE 4

Use of IL-1lp as a hybridization probe

The following method describes use of a nucleotide sequence encoding IL-1lp as a hybridization probe.

DNA comprising the coding sequence of full-length IL-1lp (as shown in Figures 3, 5, 7, 9, 15, 16 and 19; SEQ ID NOS:6, 9, 12, 15, 18, 20 and 24) is employed as a probe to screen for homologous DNAs (such as those encoding naturally-occurring variants of IL-1lp) in human tissue cDNA libraries or human tissue genomic libraries.

Hybridization and washing of filters containing either library DNAs is performed under the following high stringency conditions. Hybridization of radiolabeled IL-1lp-derived probe to

the filters is performed in a solution of 50% formamide, 5x SSC, 0.1% SDS, 0.1% sodium pyrophosphate, 50 mM sodium phosphate, pH 6.8, 2x Denhardt's solution, and 10% dextran sulfate at 42°C for 20 hours. Washing of the filters is performed in an aqueous solution of 0.1x SSC and 0.1% SDS at 42°C.

DNAs having a desired sequence identity with the DNA encoding full-length native sequence IL-1lp can then be identified using standard techniques known in the art.

EXAMPLE 5

Expression of IL-1lp in *E. coli*

This example illustrates preparation of an unglycosylated form of IL-1lp by recombinant expression in *E. coli*.

The DNA sequence encoding an IL-1lp is initially amplified using selected PCR primers. The primers should contain restriction enzyme sites which correspond to the restriction enzyme sites on the selected expression vector. A variety of expression vectors may be employed. An example of a suitable vector is pBR322 (derived from *E. coli*; see Bolivar *et al.*, *Gene*, 2:95 (1977)) which contains genes for ampicillin and tetracycline resistance. The vector is digested with restriction enzyme and dephosphorylated. The PCR amplified sequences are then ligated into the vector. The vector will preferably include sequences which encode for an antibiotic resistance gene, a trp promoter, a polyhis leader (including the first six STII codons, polyhis sequence, and enterokinase cleavage site), the IL-1lp coding region, lambda transcriptional terminator, and an argU gene.

The ligation mixture is then used to transform a selected *E. coli* strain using the methods described in Sambrook *et al.*, *supra*. Transformants are identified by their ability to grow on LB plates and antibiotic resistant colonies are then selected. Plasmid DNA can be isolated and confirmed by restriction analysis and DNA sequencing.

Selected clones can be grown overnight in liquid culture medium such as LB broth supplemented with antibiotics. The overnight culture may subsequently be used to inoculate a larger scale culture. The cells are then grown to a desired optical density, during which the expression promoter is turned on.

After culturing the cells for several more hours, the cells can be harvested by centrifugation. The cell pellet obtained by the centrifugation can be solubilized using various agents known in the art, and the solubilized IL-1lp protein can then be purified using a metal chelating column under conditions that allow tight binding of the protein.

EXAMPLE 6

Expression of IL-1lp in mammalian cells

This example illustrates preparation of a potentially glycosylated form of IL-1lp by recombinant expression in mammalian cells.

The vector, pRK5 (see EP 307,247, published March 15, 1989), is employed as the expression vector. Optionally, the IL-1lp DNA is ligated into pRK5 with selected restriction enzymes to allow insertion of the IL-1lp DNA using ligation methods such as described in Sambrook *et al.*, *supra*. The resulting vector is called pRK5-IL-1lp.

In one embodiment, the selected host cells may be 293 cells. Human 293 cells (ATCC CCL 1573) are grown to confluence in tissue culture plates in medium such as DMEM supplemented with fetal calf serum and optionally, nutrient components and/or antibiotics.

About 10 µg pRK5-IL-11p DNA is mixed with about 1 µg DNA encoding the VA RNA gene [Thimmapaya *et al.*, *Cell*, 31:543 (1982)] and dissolved in 500 µl of 1 mM Tris-HCl, 0.1 mM EDTA, 0.227 M CaCl₂. To this mixture is added, dropwise, 500 µl of 50 mM HEPES (pH 7.35), 280 mM NaCl, 1.5 mM NaPO₄, and a precipitate is allowed to form for 10 minutes at 25°C. The precipitate is suspended and added to the 293 cells and allowed to settle for about four hours at 37°C. The culture medium is aspirated off and 2 ml of 20% glycerol in PBS is added for 30 seconds. The 293 cells are then washed with serum free medium, fresh medium is added and the cells are incubated for about 5 days.

Approximately 24 hours after the transfections, the culture medium is removed and replaced with culture medium (alone) or culture medium containing 200 µCi/ml ³⁵S-cysteine and 200 µCi/ml ³⁵S-methionine. After a 12 hour incubation, the conditioned medium is collected, concentrated on a spin filter, and loaded onto a 15% SDS gel. The processed gel may be dried and exposed to film for a selected period of time to reveal the presence of IL-11p polypeptide. The cultures containing transfected cells may undergo further incubation (in serum free medium) and the medium is tested in selected bioassays.

In an alternative technique, IL-11p may be introduced into 293 cells transiently using the dextran sulfate method described by Somparyrac *et al.*, *Proc. Natl. Acad. Sci. USA*, 12: 7575 (1981). 293 cells are grown to maximal density in a spinner flask and 700 µg pRK5-IL-11p DNA is added. The cells are first concentrated from the spinner flask by centrifugation and washed with PBS. The DNA-dextran precipitate is incubated on the cell pellet for four hours. The cells are treated with 20% glycerol for 90 seconds, washed with tissue culture medium, and re-introduced into the spinner flask containing tissue culture medium, 5 µg/ml bovine insulin and 0.1 µg/ml bovine transferrin. After about four days, the conditioned media is centrifuged and filtered to remove cells and debris. The sample containing expressed IL-11p can then be concentrated and purified by any selected method, such as dialysis and/or column chromatography.

In another embodiment, IL-11p can be expressed in CHO cells. The pRK5-IL-11p can be transfected into CHO cells using known reagents such as CaPO₄ or DEAE-dextran. As described above, the cell cultures can be incubated, and the medium replaced with culture medium (alone) or medium containing a radiolabel such as ³⁵S-methionine. After determining the presence of IL-11p polypeptide, the culture medium may be replaced with serum free medium. Preferably, the cultures are incubated for about 6 days, and then the conditioned medium is harvested. The medium containing the expressed IL-11p can then be concentrated and purified by any selected method.

Epitope-tagged IL-11p may also be expressed in host CHO cells. The IL-11p may be subcloned out of the pRK5 vector. The subclone insert can undergo PCR to fuse in frame with a selected epitope tag such as a poly-his tag into a Baculovirus expression vector. The poly-his

tagged IL-11p insert can then be subcloned into a SV40 driven vector containing a selection marker such as DHFR for selection of stable clones. Finally, the CHO cells can be transfected (as described above) with the SV40 driven vector. Labeling may be performed, as described above, to verify expression. The culture medium containing the expressed poly-His tagged IL-11p can then be concentrated and purified by any selected method, such as by Ni²⁺-chelate affinity chromatography.

EXAMPLE 7

Expression of IL-11p in Yeast

The following method describes recombinant expression of IL-11p in yeast.

First, yeast expression vectors are constructed for intracellular production or secretion of IL-11p from the ADH2/GAPDH promoter. DNA encoding IL-11p and the promoter is inserted into suitable restriction enzyme sites in the selected plasmid to direct intracellular expression of IL-11p. For secretion, DNA encoding IL-11p can be cloned into the selected plasmid, together with DNA encoding the ADH2/GAPDH promoter, a native IL-11p signal peptide or other mammalian signal peptide, or, for example, a yeast alpha-factor or invertase secretory signal/leader sequence, and linker sequences (if needed) for expression of IL-11p.

Yeast cells, such as yeast strain AB110, can then be transformed with the expression plasmids described above and cultured in selected fermentation media. The transformed yeast supernatants can be analyzed by precipitation with 10% trichloroacetic acid and separation by SDS-PAGE, followed by staining of the gels with Coomassie Blue stain.

Recombinant IL-11p can subsequently be isolated and purified by removing the yeast cells from the fermentation medium by centrifugation and then concentrating the medium using selected cartridge filters. The concentrate containing IL-11p may further be purified using selected column chromatography resins.

EXAMPLE 8

Expression of IL-11p in Baculovirus-Infected Insect Cells

The following method describes recombinant expression of IL-11p in Baculovirus-infected insect cells.

The sequence coding for IL-11p is fused upstream of an epitope tag contained within a baculovirus expression vector. Such epitope tags include poly-his tags and immunoglobulin tags (like Fc regions of IgG). A variety of plasmids may be employed, including plasmids derived from commercially available plasmids such as pVL1393 (Novagen). Briefly, the sequence encoding IL-11p or the desired portion of the coding sequence of IL-11p (such as the sequence encoding the mature protein) is amplified by PCR with primers complementary to the 5' and 3' regions. The 5' primer may incorporate flanking (selected) restriction enzyme sites. The product is then digested with those selected restriction enzymes and subcloned into the expression vector.

Recombinant baculovirus is generated by co-transfecting the above plasmid and BaculoGold™ virus DNA (Pharmingen) into *Spodoptera frugiperda* ("Sf9") cells (ATCC CRL 1711) using lipofectin (commercially available from GIBCO-BRL). After 4 - 5 days of incubation at 28°C, the released viruses are harvested and used for further amplifications. Viral infection and

protein expression are performed as described by O'Reilley *et al.*, *Baculovirus expression vectors: A Laboratory Manual*, Oxford: Oxford University Press (1994).

Expressed poly-his tagged IL-1lp can then be purified, for example, by Ni²⁺-chelate affinity chromatography as follows. Extracts are prepared from recombinant virus-infected Sf9 cells as described by Rupert *et al.*, *Nature*, **362**: 175-179 (1993). Briefly, Sf9 cells are washed, resuspended in sonication buffer (25 mL Hepes, pH 7.9; 12.5 mM MgCl₂; 0.1 mM EDTA; 10% glycerol; 0.1% NP-40; 0.4 M KCl), and sonicated twice for 20 seconds on ice. The sonicates are cleared by centrifugation, and the supernatant is diluted 50-fold in loading buffer (50 mM phosphate, 300 mM NaCl, 10% glycerol, pH 7.8) and filtered through a 0.45 μ m filter. A Ni²⁺-NTA agarose column (commercially available from Qiagen) is prepared with a bed volume of 5 mL, washed with 25 mL of water and equilibrated with 25 mL of loading buffer. The filtered cell extract is loaded onto the column at 0.5 mL per minute. The column is washed to baseline A₂₈₀ with loading buffer, at which point fraction collection is started. Next, the column is washed with a secondary wash buffer (50 mM phosphate; 300 mM NaCl, 10% glycerol, pH 6.0), which elutes nonspecifically bound protein. After reaching A₂₈₀ baseline again, the column is developed with a 0 to 500 mM imidazole gradient in the secondary wash buffer. One mL fractions are collected and analyzed by SDS-PAGE and silver staining or Western blot with Ni²⁺-NTA-conjugated to alkaline phosphatase (Qiagen). Fractions containing the eluted His₁₀-tagged IL-1lp are pooled and dialyzed against loading buffer.

Alternatively, purification of the IgG tagged (or Fc tagged) IL-1lp can be performed using known chromatography techniques, including for instance, Protein A or protein G column chromatography.

EXAMPLE 9

IL-18 Receptor and IL-1 Receptor Binding of hIL-1Ra1

To facilitate the characterization of hIL-1Ra1, a PCR fragment containing the partial ORF of clone DNA85066 (Figure 1; SEQ ID NO:3) was cloned into pCMV1FLAG (IBI Kodak, described in Pan *et al.*, *Science*, **276**: 111-113) as an in-frame fusion to a NH₂-terminal preprotrypsin leader sequence and FLAG tag encoded by the vector. The entire cDNA insert of the recombinant pCMV1FLAG vector clone (designated clone DNA96786) was sequenced (Figure 2; SEQ ID NO:4). The cDNAs encoding the extracellular domain of hIL1R and hIL18R (formerly known as hIL1Rrp) were obtained by polymerase chain reaction (PCR) and cloned into a modified pCMV1FLAG vector that allowed for in-frame fusion with the Fc portion of human immunoglobulin G.

Human embryonic kidney 293 cells were grown in high glucose DMEM (Genentech, Inc). The cells were seeded at 3-4 X10⁶ per plate (100 mm) and co-transfected with pCMV1FLAG-hIL-1Ra1 and pCMV1FLAG-IL1R-ECD-Fc or pCMV1FLAG-IL18R-ECD-Fc by means of calcium phosphate precipitation. The media were changed 12 hours post transfection. The resultant conditioned media (10 ml each) were harvested after a further 70-74 hour incubation, clarified by centrifugation, aliquoted and stored at -70°C. The receptor-Fc and ligand complex from 1.5 ml conditioned medium was immunoprecipitated with protein G-Sepharose, washed three times with buffer containing 50 mM Hepes, pH7.0, 150 mM NaCl, 1 mM EDTA, 1% NP-40, and a

protease inhibitor cocktail (BMB) and resolved on a 10-20% SDS-PAGE gel. The bound ligand was identified by immunoblotting using anti-FLAG monoclonal antibody (BMB).

As shown in Figure 13A, the secreted FLAGhIL-1Ra1 fusion protein bound to IL-18R ECD and did not bind to IL-1R ECD, which indicates that hIL-1Ra1 could be an agonist or antagonist of IL-18R.

EXAMPLE 10

IL-1 Receptor and IL-18 Receptor Binding of mIL-1Ra3 and hIL-1Ra3

cDNA encoding mIL-1Ra3 (DNA92505 shown in Figure 9; SEQ ID NO:15) was cloned into pRK7 with a carboxy-terminal FLAG-tag. The resulting expression construct was transfected into human embryonic kidney 293 cells by means of calcium phosphate precipitation. 84-90 hours post transfection, the conditioned media containing secreted FLAGmIL-1Ra3 fusion protein was harvested. Conditioned media containing secreted IL-18R-Fc and IL-1R-Fc proteins were prepared as described in Example 9 above, with the exception that the 293 cells were transfected with either pCMV1FLAG-IL1R-ECD-Fc or pCMV1FLAG-IL18R-ECD-Fc alone (without pCMV1FLAG-IL-1Ra1 cotransfection).

For in vitro binding assays, IL-1R-Fc or IL-18R-Fc from 0.5 ml of the conditioned medium was immobilized to protein G-agarose and then mixed with 1.2 ml conditioned medium containing FLAGmIL-1Ra3. The receptor-ligand complexes were washed and resolved on an 10-20% SDS-PAGE gel and the bound ligand was detected by immunoblotting using anti-FLAG monoclonal antibody (Boehringer Mannheim).

As shown in Figure 14, FLAGmIL-1Ra3 fusion protein bound to IL-1R ECD and did not bind to IL-18R ECD. Since the amino acid sequence of mIL-1Ra3 is related to that of the known interleukin-1 receptor antagonist protein (IL-1Ra), mIL-1Ra3 is believed to be a novel IL-1 receptor antagonist.

cDNA encoding hIL-1Ra3 (DNA96787 shown in Figure 7; SEQ ID NO:12) was cloned into pRK7 with a carboxy-terminal FLAG tag to form pRK7hIL-1Ra3-FLAG. pCMV1FLAG-IL1R-ECD-Fc and pCMV1FLAG-IL18R-ECD-Fc were obtained as described in Example 9 above. Similarly, cDNA encoding DR6 was cloned into the modified pCMV1FLAG vector of Example 9 to form pCMV1FLAG-DR6-Fc, encoding DR6 fused to the Fc portion of human immunoglobulin G. Conditioned media containing (1) a combination of secreted FLAGhIL-1Ra3 and FLAG-DR6-Fc (2) a combination of secreted FLAGhIL-1Ra3 and FLAG-IL1R-ECD-Fc or (3) a combination of secreted FLAGhIL-1Ra3 and FLAG-IL18R-ECD-Fc were prepared by cotransfecting Human 293 cells with (1) pRK7hIL-1Ra3-FLAG and pCMV1FLAG-DR6-Fc (2) pRK7hIL-1Ra3-FLAG and pCMV1FLAG-IL1R-ECD-Fc or (3) pRK7hIL-1Ra3-FLAG and pCMV1FLAG-IL18R-ECD-Fc, culturing the transfectant cells and harvesting culture media essentially as described in Example 9 above. The receptor-Fc and ligand complex from each conditioned medium was immunoprecipitated with protein G-Sepharose or anti-FLAG monoclonal antibody, and immunoprecipitates were resolved by gel electrophoresis and immunoblotting with anti-FLAG monoclonal antibody essentially as described in Example 9 above.

As shown in Figure 13B, FLAGhIL-1Ra3 fusion protein bound to IL-1R-ECD-Fc and did not bind to IL-18R-ECD-Fc or DR6-Fc. Since the amino acid sequence of hIL-1Ra3 is related to

that of the known interleukin-1 receptor antagonist protein (IL-1Ra), hIL-3Ra3 is believed to be a novel IL-1 receptor antagonist.

EXAMPLE 11

Preparation of Antibodies that Bind IL-1 β

This example illustrates preparation of monoclonal antibodies which can specifically bind IL-1 β .

Techniques for producing the monoclonal antibodies are known in the art and are described, for instance, in Goding, *supra*. Immunogens that may be employed include purified IL-1 β , fusion proteins containing IL-1 β , and cells expressing recombinant IL-1 β on the cell surface. Selection of the immunogen can be made by the skilled artisan without undue experimentation.

Mice, such as Balb/c, are immunized with the IL-1 β immunogen emulsified in complete Freund's adjuvant and injected subcutaneously or intraperitoneally in an amount from 1-100 micrograms. Alternatively, the immunogen is emulsified in MPL-TDM adjuvant (Ribi Immunochemical Research, Hamilton, MT) and injected into the animal's hind foot pads. The immunized mice are then boosted 10 to 12 days later with additional immunogen emulsified in the selected adjuvant. Thereafter, for several weeks, the mice may also be boosted with additional immunization injections. Serum samples may be periodically obtained from the mice by retro-orbital bleeding for testing in ELISA assays to detect anti-IL-1 β antibodies.

After a suitable antibody titer has been detected, the animals "positive" for antibodies can be injected with a final intravenous injection of IL-1 β . Three to four days later, the mice are sacrificed and the spleen cells are harvested. The spleen cells are then fused (using 35% polyethylene glycol) to a selected murine myeloma cell line such as P3X63AgU.1, available from ATCC, No. CRL 1597. The fusions generate hybridoma cells which can then be plated in 96 well tissue culture plates containing HAT (hypoxanthine, aminopterin, and thymidine) medium to inhibit proliferation of non-fused cells, myeloma hybrids, and spleen cell hybrids.

The hybridoma cells will be screened in an ELISA for reactivity against IL-1 β . Determination of "positive" hybridoma cells secreting the desired monoclonal antibodies against IL-1 β is within the skill in the art.

The positive hybridoma cells can be injected intraperitoneally into syngeneic Balb/c mice to produce ascites containing the anti-IL-1 β monoclonal antibodies. Alternatively, the hybridoma cells can be grown in tissue culture flasks or roller bottles. Purification of the monoclonal antibodies produced in the ascites can be accomplished using ammonium sulfate precipitation, followed by gel exclusion chromatography. Alternatively, affinity chromatography based upon binding of antibody to protein A or protein G can be employed.

EXAMPLE 12

Isolation of DNA encoding hIL-1Ra1L, hIL-1Ra1V and hIL-1Ra1S

Several intron-containing cDNA clones related to the hIL-1Ra1 intron-containing clone DNA85066 (Figure 2) (SEQ ID NO:4) were isolated from a human testis cDNA library and fully sequenced. The intron-containing cDNA sequences were used to determine a full-length open reading frame (ORF) with the GENESCAN program (<http://CCR->

M 27.07.00

081.mit.edu/GENESCAN.html). The ORF-encoding sequence was used to design two DNA primers, ggc gga tcc aaa atg ggc tct gag gac tgg g (SEQ ID NO:29) (1Ra1016) and gcg gaa ttc taa tcg ctg acc tca ctg ggg (SEQ ID NO:30) (1Ra1017). The 1Ra1016 and 1Ra1017 primers were synthesized and used to clone cDNA from human fetal skin and SK-lu-1 cell cDNA libraries using polymerase chain reaction (PCR). Several PCR products were isolated and sequenced. Two full length cDNA clones (designated DNA102043 and DNA102044) from PCR products were found to encode hIL-1Ra1 isoforms.

The entire nucleotide sequence of clone DNA102043 is shown in Figure 15 (SEQ ID NO:18). Clone DNA102043 contains a single open reading frame with an apparent translational initiation site at nucleotide positions 4-6, and a stop codon at nucleotide positions 625-627 (Figure 15; SEQ ID NO:18). The predicted polypeptide precursor (designated hIL-1Ra1L) (Fig. 15; SEQ ID NO:19) is 207 amino acids long. The putative signal sequence extends from amino acid positions 1-34.

Clone DNA102043 (designated DNA102043-2534) was deposited with ATCC and was assigned ATCC deposit no. 203846. The full-length hIL-1Ra1L protein shown in Figure 15 (SEQ ID NO:19) has an estimated molecular weight of about 23,000 daltons and a pI of about 6.08.

Based on a sequence alignment analysis of the full length sequence (SEQ ID NO:19), hIL-1Ra1L shows significant amino acid sequence identity to hIL-1Ra β and TANGO-77 protein. In addition, a portion of the DNA sequence of clone DNA102043 (Figure 15) (SEQ ID NO:18) was found to coincide with the DNA sequence of EST AI014548 (Figure 4) (SEQ ID NO:8) and with the complement of the DNA sequence of EST AI343258 (Figure 17) (SEQ ID NO:23).

The entire nucleotide sequence of clone DNA102044 is shown in Figure 16 (SEQ ID NO:20). Clone DNA102044 contains a single open reading frame with an apparent translational initiation site at nucleotide positions 4-6, and a stop codon at nucleotide positions 505-507 (Figure 16; SEQ ID NO:20). The predicted polypeptide (designated hIL-1Ra1S) (Fig. 16; SEQ ID NO:21) is 167 amino acids long, and it is believed to behave as a mature sequence (without a presequence that is removed in post-translational processing) in certain animal cells. In addition, it is believed that other animal cells recognize and remove in post-translational processing one or more signal peptide(s) contained in the sequence extending from amino acid positions 1 to about 46.

Clone DNA102044 (designated DNA102044-2534) was deposited with ATCC and was assigned ATCC deposit no. 203855. The full-length hIL-1Ra1S protein shown in Figure 16 (SEQ ID NO:21) has an estimated molecular weight of about 18,478 daltons and a pI of about 5.5.

Based on a sequence alignment analysis of the full length sequence (SEQ ID NO:21), hIL-1Ra1S appears to be an allelic variant of TANGO-77 protein and also shows significant amino acid sequence identity to hIL-1Ra β . In addition, a portion of the DNA sequence of clone DNA102044 (Figure 16) (SEQ ID NO:20) was found to coincide with the DNA sequence of EST AI014548 (Figure 4) (SEQ ID NO:8) and with the complement of the DNA sequence of EST AI343258 (Figure 17) (SEQ ID NO:23).

EST clone AI343258 was purchased from Research Genetics (Huntsville, AL) and the cDNA insert was obtained and sequenced in its entirety. The entire sequence of the clone

27.07.00

AI343258, designated DNA114876, is shown in Figure 19 (SEQ ID NO:24). Clone DNA114876 contains a single open reading frame (ORF) with an apparent translation initiation site at nucleotide positions 73-75 and a stop codon at nucleotide positions 726-728 (Figure 19; SEQ ID NO:24), encoding a predicted polypeptide precursor (hIL-1Ra1V) (Fig. 19; SEQ ID NO:25) that is 218 amino acids long. In addition, the ORF contains an alternate translation initiation site at nucleotide positions 106-108. The predicted polypeptide (also designated hIL-1Ra1V) for translation initiated at the alternate start codon is 207 amino acids in length (lacking the first eleven residues at the N-terminus of the 218 amino acid polypeptide). It is believed that the predicted 218 amino acid and 207 amino acid polypeptides behave as mature sequences (without a presequence that is removed in post-translational processing) in certain animal cells. It is also believed that other animal cells recognize and remove one or more signal peptide(s) extending from amino acid positions 1 to about 48 (a putative leader sequence in the 218 amino acid polypeptide) or from amino acid positions 12 to 36 (a putative leader sequence in the 207 amino acid polypeptide) in the amino acid sequence of Figure 19 (SEQ ID NO:25). As shown in Example 14 below, transiently transfected CHO host cells secrete unprocessed forms of hIL-1Ra1V and hIL-1Ra1L and a single processed form that results from the removal of a signal peptide extending from amino acid positions 1 to 45 in Figure 19 (SEQ ID NO:25) or the removal of a signal peptide extending from amino acid positions 1 to 34 of Figure 15 (SEQ ID NO:19). The processed form of hIL-1Ra1V and hIL-1Ra1L secreted by transiently transfected CHO host cells has the amino acid sequence of amino acid residues 35 to 207 of Figure 15 (SEQ ID NO:19) and amino acid residues 46 to 218 of Figure 19 (SEQ ID NO:25).

Clone DNA114876 (designated DNA114876-2534) was deposited with ATCC and was assigned ATCC deposit no. 203973. The full length hIL-1Ra1V protein shown in Figure 19 (SEQ ID NO:25) has an estimated molecular weight of about 24,124 and a pI of about 6.1.

Based on a sequence alignment analysis of the full length sequence (SEQ ID NO:25), hIL-1Ra1V shows significant amino acid sequence identity to hIL-1Ra β . hIL-1Ra1V is believed to be an allelic variant of hIL-1Ra1L.

EXAMPLE 13

IL-18 Receptor and IL-1 Receptor Binding of hIL-1Ra1S

To facilitate the characterization of hIL-1Ra1S, a PCR fragment encoding amino acid residues 39-167 in the ORF of clone DNA102044 (Figure 16; SEQ ID NO:21) was cloned into pCMV1FLAG (IBI Kodak, described in Pan et al., *Science*, 276: 111-113) as an in-frame fusion to a NH₂-terminal preprotrypsin leader sequence and FLAG tag encoded by the vector to form plasmid pCMV1FLAG-IL-1Ra1S. Plasmid pCMV1FLAG-IL18R-ECD-Fc was obtained as described in Example 9 above.

Human embryonic kidney 293 cells were grown in high glucose DMEM (Genentech, Inc.). The cells were seeded at 3-4 X10⁶ per plate (100 mm) and co-transfected with pCMV1FLAG-hIL-1Ra1S and pCMV1FLAG-IL18R-ECD-Fc by means of calcium phosphate precipitation. The media were changed 12 hours post transfection. The resultant conditioned media (10 ml each) were harvested after a further 70-74 hour incubation, clarified by centrifugation, aliquoted and stored at -70°C. The receptor-Fc and ligand complex from 1.5 ml conditioned medium was

immunoprecipitated with protein G-Sepharose, washed three times with buffer containing 50 mM Hepes, pH7.0, 150 mM NaCl, 1 mM EDTA, 1% NP-40, and a protease inhibitor cocktail (BMB) and resolved on a 10-20% SDS-PAGE gel. The bound ligand was identified by immunoblotting using anti-FLAG monoclonal antibody (BMB).

The immunoblotting results indicated that the secreted FLAGhIL-1Ra1S fusion protein bound to IL-18R ECD. These data show that hIL-1Ra1S could be an agonist or antagonist of IL-18R.

EXAMPLE 14

hIL-1Ra1V, hIL-1Ra1L and hIL-1Ra3 Processing

cDNAs encoding full-length hIL-1Ra1V (amino acids 1-218 in the ORF of clone DNA114876 shown in Figure 19 (SEQ ID NO:25)), full length hIL-1Ra1L (amino acids 1-207 in the ORF of clone DNA102043 shown in Figure 15 (SEQ ID NO:19)), and full length hIL-1Ra3 (amino acids 1-155 in the ORF of clone DNA96787 shown in Figure 7 (SEQ ID NO:13)) were each cloned into a pRK7 expression vector as an in-frame fusion with a carboxy-terminal FLAG-tag sequence. In preparation for mammalian cell transient transfections, CHO DP12 cells were seeded at 4×10^6 cells per plate (100mm petri dish) in growth medium (PS20, 5% FBS, 1X GHT, 1X pen/strep, 1X L-glutamine) the day before transfection. On the day of transfection, cells were washed with PBS and fed with 10 ml serum-free transfection medium (PS20, 1X GHT). DNA-lipid transfection mixtures were prepared by adding stepwise into eppendorf tubes (1) 400 μ l transfection medium (PS20, 1X GHT); (2) 12 μ g DNA; (3) 10 μ g poly-lysine; and (4) 50 μ l Dospers liposomal transfection reagent (Boehringer Mannheim). The DNA-lipid mixtures were incubated for 15 minutes at room temperature and then added dropwise to cell culture plates. Cells were incubated overnight at 37°C. On the day after transfection, cells were washed with PBS, fed with 10 ml serum-free production medium (PS24, 10 mg/L insulin, 1X trace elements, 1.4 mg/L lipid EtOH), and placed in a 32°C incubator. After 5 days, the culture media containing the expressed proteins were harvested and cleared by centrifugation. For peptide sequencing of each expressed protein, 5-10 ml of the conditioned medium containing the expressed protein was incubated with monoclonal anti-FLAG antibody (Boehringer Mannheim) coupled to agarose beads. The immunoprecipitated FLAG-tag proteins were extensively washed with 1% NP-40 buffer (125 mM NaCl, 1 mM EDTA and 50 mM Tris-HCl, pH 7.4). The immunoprecipitates were run on a SDS polyacrylamide gel, the separated polypeptides on the gel were transferred to a PVDF membrane, the PVDF membrane was stained with Coomassie blue, and the corresponding protein bands were excised from the membrane. The amino-terminal protein sequences were obtained by conventional methods.

The processed N-terminal sequence of both of the hIL-1Ra1L and hIL-1Ra1V polypeptides was determined to be VHTSPKVKVN (SEQ ID NO:31). Approximately 50% of hIL-1Ra1L and hIL-1Ra1V material recovered from conditioned media exhibited the processed N-terminal sequence, indicating that the CHO host cells secreted a processed form corresponding to amino acid residues 35 to 207 in the amino acid sequence of Figure 15 (SEQ ID NO:19) and amino acid residues 46 to 218 in the amino acid sequence of Figure 19 (SEQ ID NO:25). The remaining 50% of the hIL-1Ra1L and hIL-1Ra1V material recovered from conditioned media exhibited an

M 27.07.00

unprocessed N-terminus, indicating that the CHO host cells also secreted unprocessed forms of hIL-1Ra1L and hIL-1Ra1V corresponding to amino acid residues 1 to 207 in the amino acid sequence of Figure 15 (SEQ ID NO:19) and to amino acid residues 1 to 218 in the amino acid sequence of Figure 19 (SEQ ID NO:25), respectively.

The processed N-terminal sequence of both of the hIL-1Ra3 and mL-1Ra3 polypeptides was determined to be VLSGALCFRM (SEQ ID NO:32). Approximately 100% of the hIL-1Ra3 and mL-1Ra3 material recovered from conditioned media exhibited the processed N-terminal sequence, indicating that the CHO host cells secreted processed forms of hIL-1Ra3 and mL-1Ra3 that lack the N-terminal methionine and correspond to amino acid residues 2 to 155 in the amino acid sequence of Figure 7 (SEQ ID NO:13) and amino acid residues 2 to 155 in the amino acid sequence of Figure 9 (SEQ ID NO:16), respectively.

Deposit of Material

The following materials have been deposited with the American Type Culture Collection, 10801 University Blvd., Manassas, VA 20110-2209, USA (ATCC):

<u>Material</u>	<u>ATCC Dep. No.</u>	<u>Deposit Date</u>
pSPORT1-based plasmid DNA92929-2534	203586	Jan. 12, 1999
pCMV-1Flag-pcmv5 plasmid DNA96786-2534	203587	Jan. 12, 1999
pT7T3D-Pac plasmid DNA85066-2534	203588	Jan. 12, 1999
pINCY-based plasmid DNA96787-2534	203589	Jan. 12, 1999
pT7T3D-Pac plasmid DNA92505-2534	203590	Jan. 12, 1999
pRK7-based plasmid DNA102043-2534	203846	March 16, 1999
pRK7-based plasmid DNA102044-2534	203855	March 16, 1999
pRK7-based plasmid DNA114876-2534	203973	April 27, 1999

These deposits were made under the provisions of the Budapest Treaty on the International Recognition of the Deposit of Microorganisms for the Purpose of Patent Procedure and the Regulations thereunder (Budapest Treaty). This assures maintenance of viable cultures of the deposits for 30 years from the date of deposit. The deposits will be made available by ATCC under the terms of the Budapest Treaty, and subject to an agreement between Genentech, Inc. and ATCC, which assures permanent and unrestricted availability of the progeny of the cultures of the deposits to the public upon issuance of the pertinent U.S. patent or upon laying open to the public of any U.S. or foreign patent application, whichever comes first, and assures availability of the progeny to one determined by the U.S. Commissioner of Patents and

Trademarks to be entitled thereto according to 35 USC §122 and the Commissioner's rules pursuant thereto (including 37 CFR §1.14 with particular reference to 886 OG 638).

5 The assignee of the present application has agreed that if a culture of the materials on deposit should die or be lost or destroyed when cultivated under suitable conditions, the materials will be promptly replaced on notification with another of the same. Availability of the deposited material is not to be construed as a license to practice the invention in contravention of the rights granted under the authority of any government in accordance with its patent laws.

10 The foregoing written specification is considered to be sufficient to enable one skilled in the art to practice the invention. The present invention is not to be limited in scope by the constructs deposited, since the deposited embodiment is intended as a single illustration of certain aspects of the invention and any constructs that are functionally equivalent are within the scope of this invention. The deposit of material herein does not constitute an admission that the written description herein contained is inadequate to enable the practice of any aspect of the invention, including the best mode thereof, nor is it to be construed as limiting the scope of the claims to the specific illustrations that it represents. Indeed, various modifications of the invention in addition to those shown and described herein will become apparent to those skilled in the art from the foregoing description and fall within the scope of the appended claims.

15

Table 2A

PRO	XXXXXXXXXXXXXXXXXX	(Length = 15 amino acids)
Comparison Protein	XXXXXXXXYYYYYYY	(Length = 12 amino acids)

% amino acid sequence identity =

(the number of identically matching amino acid residues between the two polypeptide sequences as determined by ALIGN-2) divided by (the total number of amino acid residues of the PRO polypeptide) =

5 divided by 15 = 33.3%

Table 2B

PRO	XXXXXXXXXX	(Length = 10 amino acids)
Comparison Protein	XXXXXXXXYYYYZZYZ	(Length = 15 amino acids)

% amino acid sequence identity =

(the number of identically matching amino acid residues between the two polypeptide sequences as determined by ALIGN-2) divided by (the total number of amino acid residues of the PRO polypeptide) =

5 divided by 10 = 50%

Table 2C

PRO-DNA	NNNNNNNNNNNNNNNN	(Length = 14 nucleotides)
Comparison DNA	NNNNNNLLLLLLLLLLL	(Length = 16 nucleotides)

% nucleic acid sequence identity =

(the number of identically matching nucleotides between the two nucleic acid sequences as determined by ALIGN-2) divided by (the total number of nucleotides of the PRO-DNA nucleic acid sequence) =

6 divided by 14 = 42.9%

Table 2D

PRO-DNA	NNNNNNNNNNNNNN	(Length = 12 nucleotides)
Comparison DNA	NNNNLLLVV	(Length = 9 nucleotides)

% nucleic acid sequence identity =

(the number of identically matching nucleotides between the two nucleic acid sequences as determined by ALIGN-2) divided by (the total number of nucleotides of the PRO-DNA nucleic acid sequence) =

4 divided by 12 = 33.3%

Table 3A

```

/*
*
* C-C increased from 12 to 15
5  * Z is average of EQ
  * B is average of ND
  * match with stop is _M; stop-stop = 0; J (joker) match = 0
  */
10 #define _M -8 /* value of a match with a stop */

int
  _day[26][26] = {
/* A */ { 2, 0, -2, 0, 0, -4, 1, -1, -1, 0, -1, -2, -1, 0, _M, 1, 0, -2, 1, 1, 0, 0, -6, 0, -3, 0},
/* B */ { 0, 3, -4, 3, 2, -5, 0, 1, -2, 0, 0, -3, -2, 2, _M, -1, 1, 0, 0, 0, 0, -2, -5, 0, -3, 1},
15 /* C */ { -2, 4, 15, -5, -5, -4, -3, -2, 0, -5, -6, -5, -4, _M, -3, -5, -4, 0, -2, 0, -2, -8, 0, 0, -5},
/* D */ { 0, 3, -5, 4, 3, -6, 1, 1, -2, 0, 0, -4, -3, 2, _M, -1, 2, -1, 0, 0, 0, -2, -7, 0, -4, 2},
/* E */ { 0, 2, -5, 3, 4, -5, 0, 1, -2, 0, 0, -3, -2, 1, _M, -1, 2, -1, 0, 0, 0, -2, -7, 0, -4, 3},
/* F */ { -4, -5, -4, -6, -5, 9, -5, -2, 1, 0, -5, 2, 0, -4, _M, -5, -5, -4, -3, -3, 0, -1, 0, 0, 7, -5},
20 /* G */ { 1, 0, -3, 1, 0, -5, 5, -2, -3, 0, -2, -4, -3, 0, _M, -1, -1, -3, 1, 0, 0, -1, -7, 0, -5, 0},
/* H */ { -1, 1, -3, 1, 1, -2, -2, 6, -2, 0, 0, -2, -2, 2, _M, 0, 3, 2, -1, -1, 0, -2, -3, 0, 0, 2},
/* I */ { -1, -2, -2, -2, -2, 1, -3, -2, 5, 0, -2, 2, 2, -2, _M, -2, -2, -2, -1, 0, 0, 4, -5, 0, -1, -2},
/* J */ { 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, _M, 0, 0, 0, 0, 0, 0, 0, 0, 0},
/* K */ { -1, 0, -5, 0, 0, -5, -2, 0, -2, 0, 5, -3, 0, 1, _M, -1, 1, 3, 0, 0, 0, -2, -3, 0, -4, 0},
/* L */ { -2, -3, -6, -4, -3, 2, -4, -2, 2, 0, -3, 6, 4, -3, _M, -3, -2, -3, -3, -1, 0, 2, -2, 0, -1, -2},
25 /* M */ { -1, -2, -5, -3, -2, 0, -3, -2, 2, 0, 0, 4, 6, -2, _M, -2, -1, 0, -2, -1, 0, 2, -4, 0, -2, -1},
/* N */ { 0, 2, -4, 2, 1, -4, 0, 2, -2, 0, 1, -3, -2, 2, _M, -1, 1, 0, 1, 0, 0, -2, -4, 0, -2, 1},
/* O */ { _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M},
/* P */ { 1, -1, -3, -1, -1, -5, -1, 0, -2, 0, -1, -3, -2, -1, _M, 6, 0, 0, 1, 0, 0, -1, -6, 0, -5, 0},
/* Q */ { 0, 1, -5, 2, 2, -5, -1, 3, -2, 0, 1, -2, -1, 1, _M, 0, 4, 1, -1, -1, 0, -2, -5, 0, -4, 3},
30 /* R */ { -2, 0, -4, -1, -1, -4, -3, 2, -2, 0, 3, -3, 0, 0, _M, 0, 1, 6, 0, -1, 0, -2, 2, 0, -4, 0},
/* S */ { 1, 0, 0, 0, 0, -3, 1, -1, -1, 0, 0, -3, -2, 1, _M, 1, -1, 0, 2, 1, 0, -1, -2, 0, -3, 0},
/* T */ { 1, 0, -2, 0, 0, -3, 0, -1, 0, 0, 0, -1, -1, 0, _M, 0, -1, -1, 1, 3, 0, 0, -5, 0, -3, 0},
/* U */ { 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, _M, 0, 0, 0, 0, 0, 0, 0, 0, 0},
/* V */ { 0, -2, -2, -2, -2, -1, -1, -2, 4, 0, -2, 2, 2, -2, _M, -1, -2, -2, -1, 0, 0, 4, -6, 0, -2, -2},
35 /* W */ { -6, -5, -8, -7, -7, 0, -7, -3, -5, 0, -3, -2, -4, -4, _M, -6, -5, 2, -2, -5, 0, -6, 17, 0, 0, -6},
/* X */ { 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, _M, 0, 0, 0, 0, 0, 0, 0, 0, 0},
/* Y */ { -3, -3, 0, -4, -4, 7, -5, 0, -1, 0, -4, -1, -2, -2, _M, -5, -4, -4, -3, -3, 0, -2, 0, 0, 10, -4},
/* Z */ { 0, 1, -5, 2, 3, -5, 0, 2, -2, 0, 0, -2, -1, 1, _M, 0, 3, 0, 0, 0, 0, -2, -6, 0, -4, 4}
};
40

```

Table 3B

```

/*
*/
#include <stdio.h>
#include <ctype.h>

#define MAXJMP      16      /* max jumps in a diag */
#define MAXGAP      24      /* don't continue to penalize gaps larger than this */
#define JMPS        1024    /* max jmps in an path */
#define MX          4       /* save if there's at least MX-1 bases since last jmp */

#define DMAT        3       /* value of matching bases */
#define DMIS        0       /* penalty for mismatched bases */
#define DINS0       8       /* penalty for a gap */
#define DINS1       1       /* penalty per base */
#define PINS0       8       /* penalty for a gap */
#define PINS1       4       /* penalty per residue */

struct jmp {
    short          n[MAXJMP]; /* size of jmp (neg for del) */
    unsigned short x[MAXJMP]; /* base no. of jmp in seq x */
}; /* limits seq to 2^16 -1 */

struct diag {
    int            score;      /* score at last jmp */
    long           offset;     /* offset of prev block */
    short          jmp;        /* current jmp index */
    struct jmp     jp;         /* list of jmps */
};

struct path {
    int            spc;        /* number of leading spaces */
    short          n[JMPS]; /* size of jmp (gap) */
    int            x[JMPS]; /* loc of jmp (last elem before gap) */
};

char *ofile; /* output file name */
char *namex[2]; /* seq names: getseqs() */
char *prog; /* prog name for err msgs */
char *seqx[2]; /* seqs: getseqs() */
int dmax; /* best diag: nw() */
int dmax0; /* final diag */
int dna; /* set if dna: main() */
int endgaps; /* set if penalizing end gaps */
int gapx, gapy; /* total gaps in seqs */
int len0, len1; /* seq lens */
int ngapx, ngapy; /* total size of gaps */
int smax; /* max score: nw() */
int *xbm; /* bitmap for matching */
long offset; /* current offset in jmp file */
struct diag dx; /* holds diagonals */
struct path pp[2]; /* holds path for seqs */

char *calloc(), *malloc(), *index(), *strcpy();
char *getseq(), *g_calloc();

```

Table 3C

```

/* Needleman-Wunsch alignment program
*
* usage: prog file1 file2
* where file1 and file2 are two dna or two protein sequences.
* The sequences can be in upper- or lower-case an may contain ambiguity
* Any lines beginning with ';', '>' or '<' are ignored
* Max file length is 65535 (limited by unsigned short x in the jmp struct)
* A sequence with 1/3 or more of its elements ACGTU is assumed to be DNA
* Output is in the file "align.out"
*
* The program may create a tmp file in /tmp to hold info about traceback.
* Original version developed under BSD 4.3 on a vax 8650
*/
#include "nw.h"
#include "day.h"

static _dbval[26] = {
1,14,2,13,0,0,4,11,0,0,12,0,3,15,0,0,0,5,6,8,8,7,9,0,10,0
};

static _pbval[26] = {
1, 2{(1<<(<('D'-'A'))){1<<(<('N'-'A'))}, 4, 8, 16, 32, 64,
128, 256, 0xFFFFFFF, 1<<10, 1<<11, 1<<12, 1<<13, 1<<14,
1<<15, 1<<16, 1<<17, 1<<18, 1<<19, 1<<20, 1<<21, 1<<22,
1<<23, 1<<24, 1<<25{(1<<(<('E'-'A'))){1<<(<('Q'-'A'))}
};

main(ac, av)                                main
{
    int      ac;
    char      *av[];

    prog = av[0];
    if (ac != 3) {
        fprintf(stderr, "usage: %s file1 file2\n", prog);
        fprintf(stderr, "where file1 and file2 are two dna or two protein sequences.\n");
        fprintf(stderr, "The sequences can be in upper- or lower-case\n");
        fprintf(stderr, "Any lines beginning with ';', '>' or '<' are ignored\n");
        fprintf(stderr, "Output is in the file \"align.out\"\n");
        exit(1);
    }
    namex[0] = av[1];
    namex[1] = av[2];
    seqx[0] = getseq(namex[0], &len0);
    seqx[1] = getseq(namex[1], &len1);
    xbm = (dna)? _dbval : _pbval;

    endgaps = 0;                                /* 1 to penalize endgaps */
    ofile = "align.out";                        /* output file */

    nw();                                /* fill in the matrix, get the possible jmps */
    readjmps();                            /* get the actual jmps */
    print();                                /* print stats, alignment */

    cleanup(0);                            /* unlink any tmp files */
}

```

Table 3D

```
/* do the alignment, return best score: main()
```

```
* dna: values in Fitch and Smith, PNAS, 80, 1382-1386, 1983
```

```
* pro: PAM 250 values
```

```
* When scores are equal, we prefer mismatches to any gap, prefer
```

```
* a new gap to extending an ongoing gap, and prefer a gap in seq
```

```
* to a gap in seq y.
```

```
*/
```

```
nw()
```

```
nw
```

```
{
```

```
    char      *px, *py;           /* seqs and ptrs */
    int        *ndely, *dely;     /* keep track of dely */
    int        ndelx, delx;       /* keep track of delx */
    int        *tmp;              /* for swapping row0, row1 */
    int        mis;               /* score for each type */
    int        ins0, ins1;        /* insertion penalties */
    register   id;                /* diagonal index */
    register   ij;                /* jmp index */
    register   *col0, *col1;      /* score for curr, last row */
    register   xx, yy;            /* index into seqs */
```

```
    dx = (struct diag *)g_calloc("to get diag", len0+len1+1, sizeof(struct diag));
```

```
    ndely = (int *)g_calloc("to get ndely", len1+1, sizeof(int));
```

```
    dely = (int *)g_calloc("to get dely", len1+1, sizeof(int));
```

```
    col0 = (int *)g_calloc("to get col0", len1+1, sizeof(int));
```

```
    col1 = (int *)g_calloc("to get col1", len1+1, sizeof(int));
```

```
    ins0 = (dna)? DINS0 : PINS0;
```

```
    ins1 = (dna)? DINS1 : PINS1;
```

```
    smax = -10000;
```

```
    if (endgaps) {
```

```
        for (col0[0] = dely[0] = -ins0, yy = 1; yy <= len1; yy++) {
```

```
            col0[yy] = dely[yy] = col0[yy-1] - ins1;
```

```
            ndely[yy] = yy;
```

```
        }
```

```
        col0[0] = 0;          /* Waterman Bull Math Biol 84 */
```

```
    }
```

```
    else
```

```
        for (yy = 1; yy <= len1; yy++)
```

```
            dely[yy] = -ins0;
```

```
    /* fill in match matrix
```

```
    */
```

```
    for (px = seqx[0], xx = 1; xx <= len0; px++, xx++) {
```

```
        /* initialize first entry in col
```

```
        */
```

```
        if (endgaps) {
```

```
            if (xx == 1)
```

```
                col1[0] = delx = -(ins0+ins1);
```

```
            else
```

```
                col1[0] = delx = col0[0] - ins1;
```

```
            ndelx = xx;
```

```
        }
```

```
        else {
```

```
            col1[0] = 0;
```

```
            delx = -ins0;
```

```
            ndelx = 0;
```

```
        }
```


Table 3E

...nw

```

for (py = seqx[1], yy = 1; yy <= len1; py++, yy++) {
    mis = col0[yy-1];
    if (dna)
        mis += (xbm["px-'A'"] & xbm["py-'A'"])? DMAT : DMIS;
    else
        mis += _day["px-'A'"] ["py-'A'"];

    /* update penalty for del in x seq;
     * favor new del over ongoing del
     * ignore MAXGAP if weighting endgaps
     */
    if (endgaps || ndely[yy] < MAXGAP) {
        if (col0[yy] - ins0 >= dely[yy]) {
            dely[yy] = col0[yy] - (ins0 + ins1);
            ndely[yy] = 1;
        } else {
            dely[yy] -= ins1;
            ndely[yy]++;
        }
    } else {
        if (col0[yy] - (ins0 + ins1) >= dely[yy]) {
            dely[yy] = col0[yy] - (ins0 + ins1);
            ndely[yy] = 1;
        } else
            ndely[yy]++;
    }

    /* update penalty for del in y seq;
     * favor new del over ongoing del
     */
    if (endgaps || ndelx < MAXGAP) {
        if (col1[yy-1] - ins0 >= delx) {
            delx = col1[yy-1] - (ins0 + ins1);
            ndelx = 1;
        } else {
            delx -= ins1;
            ndelx++;
        }
    }
    else {
        if (col1[yy-1] - (ins0 + ins1) >= delx) {
            delx = col1[yy-1] - (ins0 + ins1);
            ndelx = 1;
        } else
            ndelx++;
    }

    /* pick the maximum score; we're favoring
     * mis over any del and delx over dely
     */

```

Table 3F

...nw

```

id = xx - yy + len1 - 1;
if (mis >= delx && mis >= dely[yy])
    col1[yy] = mis;
else if (delx >= dely[yy]) {
    col1[yy] = delx;
    ij = dx[id].ijmp;
    if (dx[id].jp.n[0] && (!dna || (ndelx >= MAXJMP
    && xx > dx[id].jp.x[ij] + MX) || mis > dx[id].score + DINS0)) {
        dx[id].ijmp++;
        if (++ij >= MAXJMP) {

```

```

writejmps(id);
ij = dx[id].jmp = 0;
dx[id].offset = offset;
offset += sizeof(struct jmp) + sizeof(offset);
    }
    }
    dx[id].jp.n[ij] = ndelx;
    dx[id].jp.x[ij] = xx;
    dx[id].score = delx;
}
else {
    coll[yy] = dely[yy];
    ij = dx[id].jmp;
    if (dx[id].jp.n[0] && (!dna || (ndely[yy] >= MAXJMP
    && xx > dx[id].jp.x[ij]+MX) || mis > dx[id].score+DINS0)) {
        dx[id].jmp++;
        if (++ij >= MAXJMP) {
            writejmps(id);
            ij = dx[id].jmp = 0;
            dx[id].offset = offset;
            offset += sizeof(struct jmp) + sizeof(offset);
        }
        dx[id].jp.n[ij] = -ndely[yy];
        dx[id].jp.x[ij] = xx;
        dx[id].score = dely[yy];
    }
    if (xx == len0 && yy < len1) {
        /* last col
        */
        if (endgaps)
            coll[yy] -= ins0+ins1*(len1-yy);
        if (coll[yy] > smax) {
            smax = coll[yy];
            dmax = id;
        }
    }
    if (endgaps && xx < len0)
        coll[yy-1] -= ins0+ins1*(len0-xx);
    if (coll[yy-1] > smax) {
        smax = coll[yy-1];
        dmax = id;
    }
    tmp = col0; col0 = coll; coll = tmp;
}
(void) free((char *)ndely);
(void) free((char *)dely);
(void) free((char *)col0);
(void) free((char *)coll);
}

```

Table 3G

```

/*
 *
 * print() -- only routine visible outside this module
 *
 * static:
 * getmat() -- trace back best path, count matches; print()
 * pr_align() -- print alignment of described in array p[]; print()
 * dumpblock() -- dump a block of lines with numbers, stars; pr_align()
 * nums() -- put out a number line; dumpblock()
 * putline() -- put out a line (name, [num], seq, [num]): dumpblock()
 * stars() -- put a line of stars; dumpblock()
 * stripname() -- strip any path and prefix from a seqname
 */

#include "nw.h"

#define SPC 3
#define P_LINE 256 /* maximum output line */
#define P_SPC 3 /* space between name or num and seq */

extern _day[26][26];
int olen; /* set output line length */
FILE *fx; /* output file */

print()
{
    int lx, ly, firstgap, lastgap; /* overlap */

    if ((fx = fopen(ofile, "w")) == 0) {
        fprintf(stderr, "%s: can't write %s\n", prog, ofile);
        cleanup(1);
    }
    fprintf(fx, "<first sequence: %s (length = %d)\n", namex[0], len0);
    fprintf(fx, "<second sequence: %s (length = %d)\n", namex[1], len1);
    olen = 60;
    lx = len0;
    ly = len1;
    firstgap = lastgap = 0;
    if (dmax < len1 - 1) { /* leading gap in x */
        pp[0].spc = firstgap = len1 - dmax - 1;
        ly -= pp[0].spc;
    }
    else if (dmax > len1 - 1) { /* leading gap in y */
        pp[1].spc = firstgap = dmax - (len1 - 1);
        lx -= pp[1].spc;
    }
    if (dmax0 < len0 - 1) { /* trailing gap in x */
        lastgap = len0 - dmax0 - 1;
        lx -= lastgap;
    }
    else if (dmax0 > len0 - 1) { /* trailing gap in y */
        lastgap = dmax0 - (len0 - 1);
        ly -= lastgap;
    }
    getmat(lx, ly, firstgap, lastgap);
    pr_align();
}

```

print

Table 3H

```

/*
 * trace back the best path, count matches
 */
5  static
   getmat(lx, ly, firstgap, lastgap)                                getmat
   {
       int      lx, ly;                                           /* "core" (minus endgaps) */
       int      firstgap, lastgap;                                /* leading trailing overlap */

10      int      nm, i0, i1, siz0, siz1;
       char      outx[32];
       double     pct;
       register   n0, n1;
15      register char *p0, *p1;

       /* get total matches, score
        */
       i0 = i1 = siz0 = siz1 = 0;
       p0 = seqx[0] + pp[1].spc;
       p1 = seqx[1] + pp[0].spc;
       n0 = pp[1].spc + 1;
       n1 = pp[0].spc + 1;

20      nm = 0;
       while ( *p0 && *p1 ) {
           if (siz0) {
               p1++;
               n1++;
               siz0--;
30           }
           else if (siz1) {
               p0++;
               n0++;
               siz1--;
35           }
           else {
               if (x[bm[*p0-'A']&x[bm[*p1-'A']])
                   nm++;
               if (n0++ == pp[0].x[i0])
                   siz0 = pp[0].n[i0++];
               if (n1++ == pp[1].x[i1])
                   siz1 = pp[1].n[i1++];
40               p0++;
               p1++;
45           }
       }

       /* pct homology:
        * if penalizing endgaps, base is the shorter seq
        * else, knock off overhangs and take shorter core
        */
       if (endgaps)
           lx = (len0 < len1)? len0 : len1;
       else
55          lx = (lx < ly)? lx : ly;

       pct = 100.*((double)nm)/(double)lx;
       fprintf(fx, "\n");
       fprintf(fx, "< %d match%s in an overlap of %d: %.2f percent similarity\n",
           nm, (nm == 1)? "" : "es", lx, pct);

```

Table 3I

```

fprintf(fx, "< gaps in first sequence: %d", gapx);
5  if (gapx) {
    (void) sprintf(outh, " (%d %s%s)",
        gapx, (dna)? "base":"residue", (ngapx == 1)? "":"s");
    fprintf(fx, "%s", outh);

    fprintf(fx, ", gaps in second sequence: %d", gapy);
10  if (gapy) {
    (void) sprintf(outh, " (%d %s%s)",
        gapy, (dna)? "base":"residue", (ngapy == 1)? "":"s");
    fprintf(fx, "%s", outh);
    }
15  if (dna)
    fprintf(fx,
        "\n< score: %d (match = %d, mismatch = %d, gap penalty = %d + %d per base)\n",
        smax, DMAT, DMIS, DINS0, DINS1);
    else
20  fprintf(fx,
        "\n< score: %d (Dayhoff PAM 250 matrix, gap penalty = %d + %d per residue)\n",
        smax, PINS0, PINS1);

    if (endgaps)
25  fprintf(fx,
        "< endgaps penalized. left endgap: %d %s%s, right endgap: %d %s%s\n",
        firstgap, (dna)? "base" : "residue", (firstgap == 1)? "":"s",
        lastgap, (dna)? "base" : "residue", (lastgap == 1)? "":"s");
    else
30  fprintf(fx, "< endgaps not penalized\n");
}

static nm; /* matches in core -- for checking */
static lmax; /* lengths of stripped file names */
static ij[2]; /* jmp index for a path */
35 static nc[2]; /* number at start of current line */
static ni[2]; /* current elem number -- for gapping */
static siz[2];
static char *ps[2]; /* ptr to current element */
static char *po[2]; /* ptr to next output char slot */
40 static char out[2][P_LINE]; /* output line */
static char star[P_LINE]; /* set by stars() */

/*
45 * print alignment of described in struct path pp[]
*/
static
pr_align()
{
50  int nn; /* char count */
  int more;
  register i;

  for (i = 0, lmax = 0; i < 2; i++) {
55  nn = strlen(name[i]);
    if (nn > lmax)
        lmax = nn;

    nc[i] = 1;
    ni[i] = 1;
    siz[i] = ij[i] = 0;
60  ps[i] = seq[i];
    po[i] = out[i];

```

...getmat

pr_align

Table 3J**...pr_align**

```

5      for (nn = nm = 0, more = 1; more;) {
          for (i = more = 0; i < 2; i++) {
              /*
              * do we have more of this sequence?
              */
              if (!*ps[i])
10                 continue;

                more++;

                if (pp[i].spc) { /* leading space */
                    *po[i]++ = ' ';
                    pp[i].spc--;
                }
                else if (siz[i]) { /* in a gap */
                    *po[i]++ = '-';
                    siz[i]--;
20                }
                else { /* we're putting a seq element
                    */
                    *po[i] = *ps[i];
                    if (islower(*ps[i]))
                        *ps[i] = toupper(*ps[i]);
                    po[i]++;
                    ps[i]++;

                    /*
                    * are we at next gap for this seq?
                    */
                    if (ni[i] == pp[i].x[ij[i]]) {
                        /*
                        * we need to merge all gaps
                        * at this location
                        */
                        siz[i] = pp[i].n[ij[i]] + 1;
                        while (ni[i] == pp[i].x[ij[i]])
                            siz[i] += pp[i].n[ij[i]] + 1;
40                    }
                    ni[i]++;
                }
            }
            if (++nn == olen || !more && nn) {
25                dumpblock();
                for (i = 0; i < 2; i++)
                    po[i] = out[i];
                nn = 0;
            }
        }
    }

    /*
    * dump a block of lines, including numbers, stars: pr_align()
    */
    static
    dumpblock()
    {
        register i;
60        for (i = 0; i < 2; i++)
            *po[i]-- = '\0';
    }

```

dumpblock

Table 3K

...dumpblock

```

5      (void) puts("\n", fx);
      for (i = 0; i < 2; i++) {
          if (*out[i] && (*out[i] != ' ' || *(po[i]) != ' ')) {
              if (i == 0)
                  nums(i);
              if (i == 0 && *out[1])
                  stars();
              putline(i);
              if (i == 0 && *out[1])
                  fprintf(fx, star);
              if (i == 1)
                  nums(i);
          }
      }
}

20 /*
   * put out a number line: dumpblock()
   */
static
nums(ix)                                nums
25 {
    int    ix;        /* index in out[] holding seq line */
    {
        char    nline[P_LINE];
        register i, j;
        register char *pn, *px, *py;

30        for (pn = nline, i = 0; i < lmax+P_SPC; i++, pn++)
            *pn = ' ';
        for (i = nc[ix], py = out[ix]; *py; py++, pn++) {
            if (*py == ' ' || *py == '-')
                *pn = ' ';
            else {
35                if (i%10 == 0 || (i == 1 && nc[ix] != 1)) {
                    j = (i < 0)? -i : i;
                    for (px = pn; j /= 10, px--)
                        *px = j%10 + '0';
                    if (i < 0)
                        *px = '-';
                }
                else
40                    *pn = ' ';
                i++;
            }
        }
        *pn = '\0';
        nc[ix] = i;
        for (pn = nline; *pn; pn++)
            (void) puts(*pn, fx);
        (void) puts("\n", fx);
    }
}

55 /*
   * put out a line (name, [num], seq, [num]): dumpblock()
   */
static
putline(ix)                                putline
60 {
    int    ix;

```

Table 3L

...putline

```

5      int          i;
      register char *px;

      for (px = namex[ix], i = 0; *px && *px != ':'; px++, i++)
          (void) puts(*px, fx);
10     for (; i < lmax+P_SPC; i++)
          (void) puts(' ', fx);

      /* these count from 1:
       * ni[] is current element (from 1)
       * nc[] is number at start of current line
       */
15     for (px = out[ix]; *px; px++)
          (void) puts(*px&0x7F, fx);
      (void) puts('\n', fx);
20 }

/*
 * put a line of stars (seqs always in out[0], out[1]): dumpblock()
 */
25 static
stars()
{
    int          i;
    register char *p0, *p1, cx, *px;

    if (!*out[0] || (*out[0] == ' ' && *(p0[0]) == ' ') ||
        !*out[1] || (*out[1] == ' ' && *(p0[1]) == ' '))
30         return;
    px = star;
    for (i = lmax+P_SPC; i; i--)
        *px++ = ' ';

    for (p0 = out[0], p1 = out[1]; *p0 && *p1; p0++, p1++) {
40         if (isalpha(*p0) && isalpha(*p1)) {
            if (xbm[*p0-'A']&xbm[*p1-'A']) {
                cx = '*';
                nm++;
            }
            else if (ldna && _day[*p0-'A'][*p1-'A'] > 0)
50                 cx = '.';
            else
                cx = ' ';
        }
        else
            cx = ' ';
        *px++ = cx;
    }
    *px++ = '\n';
55     *px = '\0';
}

```

stars

Table 3M

```

/*
 * strip path or prefix from pn, return len: pr_align()
 */
5 static
stripname(pn)
char *pn; /* file name (may be path) */
{
10 register char *px, *py;

py = 0;
for (px = pn; *px; px++)
    if (*px == '/')
        py = px + 1;
15 if (py)
    (void) strcpy(pn, py);
    return(strlen(pn));
}
20

```

stripname

Table 3N

```

/*
 * cleanup() -- cleanup any tmp file
 * getseq() -- read in seq, set dna, len, maxlen
25 * g_calloc() -- calloc() with error checking
 * readjmps() -- get the good jmps, from tmp file if necessary
 * writejumps() -- write a filled array of jmps to a tmp file: nw()
 */
#include "nw.h"
#include <sys/file.h>
30
char *jname = "/tmp/homgXXXXXX"; /* tmp file for jmps */
FILE *fj;
35 int cleanup(); /* cleanup tmp file */
long lseek();

/*
 * remove any tmp file if we blow
40 */
cleanup(i)
int i;
{
45 if (fj)
    (void) unlink(jname);
    exit(i);
}

/*
50 * read, return ptr to seq, set dna, len, maxlen
 * skip lines starting with ';', '<', or '>'
 * seq in upper or lower case
 */
char *
55 getseq(file, len)
char *file; /* file name */
int *len; /* seq len */
{
60 char line[1024], *pseq;
register char *px, *py;
int natgc, then;
FILE *fp;

if ((fp = fopen(file, "r")) == 0) {

```

cleanup

getseq

```

    fprintf(stderr, "%s: can't read %s\n", prog, file);
    exit(1);
}
den = natgc = 0;
5 while (fgets(line, 1024, fp)) {
    if (*line == ';' || *line == '<' || *line == '>')
        continue;
    for (px = line; *px != '\n'; px++)
10     if (isupper(*px) || islower(*px))
        den++;
}
if ((pseq = malloc((unsigned)(den+6))) == 0) {
    fprintf(stderr, "%s: malloc() failed to get %d bytes for %s\n", prog, den+6, file);
15     exit(1);
}
pseq[0] = pseq[1] = pseq[2] = pseq[3] = '\0';

```

Table 30

...getseq

```

20 py = pseq + 4;
   *len = den;
   rewind(fp);

   while (fgets(line, 1024, fp)) {
25       if (*line == ';' || *line == '<' || *line == '>')
           continue;
       for (px = line; *px != '\n'; px++) {
           if (isupper(*px))
30               *py++ = *px;
           else if (islower(*px))
               *py++ = toupper(*px);
           if (index("ATGCU", *(py-1))
               natgc++;
       }
       *py++ = '\0';
       *py = '\0';
       (void) fclose(fp);
       dna = natgc > (den/3);
40       return(pseq+4);
   }

   char *
g_alloc(msg, nx, sz)
50 char *msg; /* program, calling routine */
   int nx, sz; /* number and size of elements */
   {
       char *px, *calloc();
55     if ((px = calloc((unsigned)nx, (unsigned)sz)) == 0) {
         if (*msg) {
             fprintf(stderr, "%s: g_alloc() failed %s (n=%d, sz=%d)\n", prog, msg, nx, sz);
             exit(1);
         }
       }
       return(px);
   }

   /*
60  * get final jmps from dx[] or tmp file. set pp[], reset dmax: main()
   */
   readjumps()
   {
       int fd = -1;

```

readjumps

```

int          siz, i0, i1;
register     i, j, xx;

if (f) {
    (void) fclose(f);
    if ((fd = open(jname, O_RDONLY, 0)) < 0) {
        fprintf(stderr, "%s: can't open() %s\n", prog, jname);
        cleanup(1);
    }
}
for (i = i0 = i1 = 0, dmax0 = dmax, xx = len0; ; i++) {
    while (1) {
        for (j = dx[dmax].ijmp; j >= 0 && dx[dmax].jp.x[j] >= xx; j--)

```

Table 3P

...readjmps

```

    if (j < 0 && dx[dmax].offset && f) {
        (void) lseek(fd, dx[dmax].offset, 0);
        (void) read(fd, (char *)&dx[dmax].jp, sizeof(struct jmp));
        (void) read(fd, (char *)&dx[dmax].offset, sizeof(dx[dmax].offset));
        dx[dmax].ijmp = MAXJMP-1;
    }
    else
        break;
}
if (i >= JMPS) {
    fprintf(stderr, "%s: too many gaps in alignment\n", prog);
    cleanup(1);
}
if (j >= 0) {
    siz = dx[dmax].jp.n[j];
    xx = dx[dmax].jp.x[j];
    dmax += siz;
    if (siz < 0) { /* gap in second seq */
        pp[1].n[i1] = -siz;
        xx += siz;

        /* id = xx - yy + len1 - 1
        */
        pp[1].x[i1] = xx - dmax + len1 - 1;
        gapy ++;
        ngapy -= siz;
/* ignore MAXGAP when doing endgaps */
        siz = (-siz < MAXGAP || endgaps)? -siz : MAXGAP;
        i1 ++;
    }
    else if (siz > 0) { /* gap in first seq */
        pp[0].n[i0] = siz;
        pp[0].x[i0] = xx;
        gapx ++;
        ngapx += siz;
/* ignore MAXGAP when doing endgaps */
        siz = (siz < MAXGAP || endgaps)? siz : MAXGAP;
        i0 ++;
    }
}
else
    break;
}

/* reverse the order of jmps
*/
for (j = 0, i0--; j < i0; j++, i0--) {
    i = pp[0].n[j]; pp[0].n[j] = pp[0].n[i0]; pp[0].n[i0] = i;

```

```

        i = pp[0].x[j]; pp[0].x[j] = pp[0].x[i0]; pp[0].x[i0] = i;
    }
    for (j = 0, i1--; j < i1; j++, i1--) {
        i = pp[1].n[j]; pp[1].n[j] = pp[1].n[i1]; pp[1].n[i1] = i;
        i = pp[1].x[j]; pp[1].x[j] = pp[1].x[i1]; pp[1].x[i1] = i;
    }
    if (fd >= 0)
        (void) close(fd);
    if (fj) {
        (void) unlink(jname);
        fj = 0;
        offset = 0;
    }

```

Table 3Q

```

15  /*
    * write a filled jmp struct offset of the prev one (if any): nw()
    */

```

```

writejmps(ix)

```

writejmps

```

20  {
    int ix;
    char *mktmp();

    if (!fj) {
        if (mktmp(jname) < 0) {
            fprintf(stderr, "%s: can't mktmp() %s\n", prog, jname);
            cleanup(1);
        }
        if ((fj = fopen(jname, "w")) == 0) {
            fprintf(stderr, "%s: can't write %s\n", prog, jname);
            exit(1);
        }
    }
    (void) fwrite((char *)&dx[ix].jp, sizeof(struct jmp), 1, fj);
    (void) fwrite((char *)&dx[ix].offset, sizeof(dx[ix].offset), 1, fj);
35  }

```

What is claimed is:

1. An isolated DNA molecule selected from the group consisting of: (1) a DNA molecule encoding an hIL-1Ra1 polypeptide comprising the amino acid sequence of amino acid residues from about 37 to about 203 of Figure 2 (SEQ ID NO:5); (2) a DNA molecule encoding an hIL-1Ra1 polypeptide comprising the amino acid sequence of amino acid residues from about 15 to about 193 of Figure 3 (SEQ ID NO:7); (3) a DNA molecule encoding an hIL-1Ra2 polypeptide comprising the amino acid sequence of amino acid residues from about 1 to about 134 of Figure 5 (SEQ ID NO:10); (4) a DNA molecule encoding an hIL-1Ra3 polypeptide comprising the amino acid sequence of amino acid residues from about 95 to about 134 of Figure 7 (SEQ ID NO:13); (5) a DNA molecule encoding a mL-1Ra3 polypeptide comprising the amino acid sequence of amino acid residues from about 95 to about 134 of Figure 9 (SEQ ID NO:16); (6) a DNA molecule encoding an hIL-1Ra1L polypeptide comprising the amino acid sequence of amino acid residues from about 26 to about 207 of Figure 15 (SEQ ID NO:19); (7) a DNA molecule encoding an hIL-1Ra1S polypeptide comprising the amino acid sequence of amino acid residues from about 26 to about 167 of Figure 16 (SEQ ID NO:21); (8) a DNA molecule encoding an hIL-1Ra1V polypeptide comprising the amino acid sequence of amino acid residues from about 46 to about 218 of Figure 19 (SEQ ID NO:25); and (9) the complement of any of the DNA molecules of (1)-(8).

2. The isolated DNA molecule of claim 1 selected from the group consisting of: (1) a DNA molecule encoding an hIL-1Ra1 polypeptide comprising the amino acid sequence of amino acid residues from about 37 to about 203 of Figure 2 (SEQ ID NO:5); (2) a DNA molecule encoding an hIL-1Ra1 polypeptide comprising the amino acid sequence of amino acid residues from about 15 to about 193 of Figure 3 (SEQ ID NO:7); (3) a DNA molecule encoding an hIL-1Ra2 polypeptide comprising the amino acid sequence of amino acid residues from about 1 to about 134 of Figure 5 (SEQ ID NO:10); (4) a DNA molecule encoding an hIL-1Ra3 polypeptide comprising the amino acid sequence of amino acid residues from about 95 to about 134 of Figure 7 (SEQ ID NO:13); (5) a DNA molecule encoding a mL-1Ra3 polypeptide comprising the amino acid sequence of amino acid residues from about 95 to about 134 of Figure 9 (SEQ ID NO:16); and (6) the complement of any of the DNA molecules of (1)-(5).

3. The isolated DNA molecule of Claim 1 selected from the group consisting of: (1) a DNA molecule encoding an hIL-1Ra1L polypeptide comprising the amino acid sequence of amino acid residues from about 26 to about 207 of Figure 15 (SEQ ID NO:19); (2) a DNA molecule encoding an hIL-1Ra1S polypeptide comprising the amino acid sequence of amino acid residues from about 26 to about 167 of Figure 16 (SEQ ID NO:21); (3) a DNA molecule encoding an hIL-1Ra1V polypeptide comprising the amino acid sequence of amino acid residues from about 46 to about 218 of Figure 19 (SEQ ID NO:25); and (4) the complement of any of the DNA molecules of (1)-(3).

4. The isolated DNA molecule of Claim 3 selected from the group consisting of: (1) a DNA molecule encoding an hIL-1Ra1L polypeptide comprising the amino acid sequence of amino acid residues from about 1 to about 207 of Figure 15 (SEQ ID NO:19); (2) a DNA molecule encoding an hIL-1Ra1S polypeptide comprising the amino acid sequence of amino acid residues from about 1 to about 167 of Figure 16 (SEQ ID NO:21); (3) a DNA molecule encoding an hIL-

1Ra1V polypeptide comprising the amino acid sequence of amino acid residues from about 1 to about 218 of Figure 19 (SEQ ID NO:25); and (4) the complement of any of the DNA molecules of (1)-(2).

5 5. The isolated DNA molecule of Claim 2 selected from the group consisting of: (1) a DNA molecule encoding an hIL-1Ra3 polypeptide comprising the amino acid sequence of amino acid residues from about 34 to about 155 of Figure 7 (SEQ ID NO:13); (2) a DNA molecule encoding a mL-1Ra3 polypeptide comprising the amino acid sequence of amino acid residues from about 34 to about 155 of Figure 9 (SEQ ID NO:16); and (3) the complement of any of the DNA molecules of (1)-(2).

10 6. The isolated DNA molecule of Claim 5 selected from the group consisting of: (1) a DNA molecule encoding an hIL-1Ra3 polypeptide comprising the amino acid sequence of amino acid residues from about 2 to about 155 of Figure 7 (SEQ ID NO:13); (2) a DNA molecule encoding a mL-1Ra3 polypeptide comprising the amino acid sequence of amino acid residues from about 2 to about 155 of Figure 9 (SEQ ID NO:16); and (3) the complement of any of the DNA molecules of (1)-(2).

15 7. The isolated DNA molecule of Claim 1 selected from the group consisting of: (1) a DNA molecule which encodes an hIL-1Ra1 polypeptide, and which comprises the nucleic acid sequence of nucleotide positions from about 118 to about 618 in the sense strand of Figure 2 (SEQ ID NO:4); (2) a DNA molecule which encodes an hIL-1Ra1 polypeptide, and which
20 comprises the nucleic acid sequence of nucleotide positions from about 145 to about 681 in the sense strand of Figure 3 (SEQ ID NO:6); (3) a DNA molecule which encodes an hIL-1Ra2 polypeptide, and which comprises the nucleic acid sequence of nucleotide positions from about 96 to about 497 in the sense strand of Figure 5 (SEQ ID NO:9); (4) a DNA molecule which encodes an hIL-1Ra3 polypeptide, and which comprises the nucleic acid sequence of nucleotide
25 positions from about 283 to about 402 in the sense strand of Figure 7 (SEQ ID NO:12); (5) a DNA molecule which encodes a mL-1Ra3 polypeptide, and which comprises the nucleic acid sequence of nucleotide positions from about 427 to about 546 in the sense strand of Figure 9 (SEQ ID NO:15); (6) a DNA molecule which encodes an hIL-1Ra1L polypeptide, and which comprises the nucleic acid sequence of nucleotide positions from about 79 to about 624 in the sense strand of
30 Figure 15 (SEQ ID NO:18); (7) a DNA molecule which encodes an hIL-1Ra1S polypeptide, and which comprises the nucleic acid sequence of nucleotide positions from about 79 to about 504 in the sense strand of Figure 16 (SEQ ID NO:20); (8) a DNA molecule which encodes an hIL-1Ra1V polypeptide, and which comprises the nucleic acid sequence of nucleotide positions from about 208 to about 726 in the sense strand of Figure 19 (SEQ ID NO:24); and (9) the complement
35 of any of the DNA molecules of (1)-(8).

8. The isolated DNA molecule of Claim 7 selected from the group consisting of: (1) a DNA molecule which encodes an hIL-1Ra1 polypeptide, and which comprises the nucleic acid sequence of nucleotide positions from about 118 to about 618 in the sense strand of Figure 2 (SEQ ID NO:4); (2) a DNA molecule which encodes an hIL-1Ra1 polypeptide, and which
40 comprises the nucleic acid sequence of nucleotide positions from about 145 to about 681 in the sense strand of Figure 3 (SEQ ID NO:6); (3) a DNA molecule which encodes an hIL-1Ra2

polypeptide, and which comprises the nucleic acid sequence of nucleotide positions from about 96 to about 497 in the sense strand of Figure 5 (SEQ ID NO:9); (4) a DNA molecule which encodes an hIL-1Ra3 polypeptide, and which comprises the nucleic acid sequence of nucleotide positions from about 283 to about 402 in the sense strand of Figure 7 (SEQ ID NO:12); (5) a DNA molecule which encodes a mL-1Ra3 polypeptide, and which comprises the nucleic acid sequence of nucleotide positions from about 427 to about 546 in the sense strand of Figure 9 (SEQ ID NO:15); and (6) the complement of any of the DNA molecules of (1)-(5).

9. The isolated DNA molecule of Claim 7 selected from the group consisting of: (1) a DNA molecule which encodes an hIL-1Ra1L polypeptide, and which comprises the nucleic acid sequence of nucleotide positions from about 79 to about 624 in the sense strand of Figure 15 (SEQ ID NO:18); (2) a DNA molecule which encodes an hIL-1Ra1S polypeptide, and which comprises the nucleic acid sequence of nucleotide positions from about 79 to about 504 in the sense strand of Figure 16 (SEQ ID NO:20); (3) a DNA molecule which encodes an hIL-1Ra1V polypeptide, and which comprises the nucleic acid sequence of nucleotide positions from about 208 to about 726 in the sense strand of Figure 19 (SEQ ID NO:24); and (4) the complement of any of the DNA molecules of (1)-(3).

10. The isolated DNA molecule of Claim 9 selected from the group consisting of: (1) a DNA molecule which encodes an hIL-1Ra1L polypeptide, and which comprises the nucleic acid sequence of nucleotide positions from about 4 to about 624 in the sense strand of Figure 15 (SEQ ID NO:18); (2) a DNA molecule which encodes an hIL-1Ra1S polypeptide, and which comprises the nucleic acid sequence of nucleotide positions from about 4 to about 504 in the sense strand of Figure 16 (SEQ ID NO:20); (3) a DNA molecule which encodes an hIL-1Ra1V polypeptide, and which comprises the nucleic acid sequence of nucleotide positions from about 73 to about 726 in the sense strand of Figure 19 (SEQ ID NO:24); and (4) the complement of any of the DNA molecules of (1)-(3).

11. The isolated nucleic acid molecule of Claim 8 selected from the group consisting of: (1) a DNA molecule comprising the nucleic acid sequence of nucleotide positions from about 103 to about 681 in the sense strand of Figure 3 (SEQ ID NO:6); (2) a DNA molecule comprising the nucleic acid sequence of nucleotide positions from about 100 to about 465 in the sense strand of Figure 7 (SEQ ID NO:12); (3) a DNA molecule comprising the nucleic acid sequence of nucleotide positions from about 244 to about 609 in the sense strand of Figure 9 (SEQ ID NO:15); and (4) the complement of any of the DNA molecules of (1)-(3).

12. The isolated nucleic acid molecule of Claim 8 comprising (a) the complete DNA sequence in the sense strand of Figure 2 (SEQ ID NO:4), Figure 3 (SEQ ID NO:6), Figure 5 (SEQ ID NO:9), Figure 7 (SEQ ID NO:12), or Figure 9 (SEQ ID NO:15), or (b) the complement of (a).

13. An isolated nucleic acid molecule encoding an IL-11p polypeptide, comprising DNA hybridizing to the complement of a nucleic acid sequence selected from the group consisting of: (1) the nucleic acid sequence consisting of nucleotide positions from about 238 to about 465 in the sense strand of Figure 7 (SEQ ID NO:12); (2) the nucleic acid sequence consisting of nucleotide positions from about 427 to about 609 in the sense strand of Figure 9 (SEQ ID

NO:15); and (3) the nucleic acid sequence consisting of nucleotide positions from about 114 to about 135 in the sense strand of Figure 15 (SEQ ID NO:18).

14. An isolated nucleic acid molecule comprising (a) a DNA molecule encoding a polypeptide selected from the group consisting of: (1) a polypeptide comprising the entire amino acid sequence encoded by the longest open reading frame in the cDNA insert in the vector deposited as ATCC Deposit No. 203588; (2) a polypeptide comprising the entire amino acid sequence, or the entire amino acid sequence excluding the 36 N-terminal amino acid residues of such sequence, encoded by the longest open reading frame in the cDNA insert in the vector deposited as ATCC Deposit No. 203587; (3) a polypeptide comprising the entire amino acid sequence encoded by the longest open reading frame in the cDNA insert in the vector deposited as ATCC Deposit No. 203586; (4) a polypeptide comprising the entire amino acid sequence, or the entire amino acid sequence excluding the N-terminal amino acid residue of such sequence, encoded by the longest open reading frame in the cDNA insert in the vector deposited as ATCC Deposit No. 203589; (5) a polypeptide comprising the entire amino acid sequence, or the entire amino acid sequence excluding the N-terminal amino acid residue of such sequence, encoded by the cDNA insert in the vector deposited as ATCC Deposit No. 203590; (6) a polypeptide comprising the entire amino acid sequence, or the entire amino acid sequence excluding the 34 N-terminal amino acid residues of such sequence, encoded by the longest open reading frame in the cDNA insert in the vector deposited as ATCC Deposit No. 203846; (7) a polypeptide comprising the entire amino acid sequence encoded by the longest open reading frame in the cDNA insert in the vector deposited as ATCC Deposit No. 203855; and (8) a polypeptide comprising the entire amino acid sequence, or the entire amino acid sequence excluding the 45 N-terminal amino acid residues of such sequence, encoded by the longest open reading frame in the cDNA insert in the vector deposited as ATCC Deposit No. 203973; or (b) the complement of the DNA molecule of (a).

15. An isolated nucleic acid molecule comprising (a) DNA encoding the IL-11p polypeptide of Claim 21, or (b) the complement of the DNA of (a).

16. A vector comprising the nucleic acid of Claim 1.

17. The vector of Claim 16 operably linked to control sequences recognized by a host cell transfected with the vector.

18. A host cell comprising the vector of Claim 16.

19. A process for producing an IL-11p polypeptide comprising the steps of: (1) culturing a host cell comprising the DNA molecule of claim 15 under conditions suitable for expression of the IL-11p polypeptide encoded by the DNA molecule; and (2) recovering said IL-11p polypeptide from the cell culture.

20. An isolated IL-11p polypeptide encoded by the nucleic acid molecule of Claim 1.

21. An isolated IL-11p polypeptide selected from the group consisting of: (1) an hIL-1Ra1V polypeptide consisting of an amino acid sequence having at least an 80% sequence identity to the sequence of amino acid residues from about 46 to about 218 of Figure 19 (SEQ ID NO:25); (2) an hIL-1Ra3 polypeptide consisting of an amino acid sequence having at least an 80% sequence identity to the sequence of amino acid residues from about 95 to about 134 of

Figure 7 (SEQ ID NO:13); and (3) a mL-1Ra3 polypeptide consisting of an amino acid sequence having at least an 80% sequence identity to the sequence of amino acid residues from about 95 to about 134 of Figure 9 (SEQ ID NO:16).

22. An isolated IL-11p polypeptide selected from the group consisting of: (1) an hIL-1Ra1 polypeptide comprising amino acid residues from about 37 to about 203 of Figure 2 (SEQ ID NO:5); (2) an hIL-1Ra1 polypeptide comprising amino acid residues from about 15 to about 193 of Figure 3 (SEQ ID NO:7); (3) an hIL-1Ra2 polypeptide comprising amino acid residues from about 1 to about 134 of Figure 5 (SEQ ID NO:10); (4) an hIL-1Ra3 polypeptide comprising amino acid residues from about 95 to about 134 of Figure 7 (SEQ ID NO:13); (5) a mL-1Ra3 polypeptide comprising amino acid residues from about 95 to about 134 of Figure 9 (SEQ ID NO:16); (6) an hIL-1Ra1L polypeptide comprising amino acid residues from about 26 to about 207 of Figure 15 (SEQ ID NO:19); (7) an hIL-1Ra1S polypeptide comprising amino acid residues from about 26 to about 167 of Figure 16 (SEQ ID NO:21); and (8) an hIL-1Ra1V polypeptide comprising amino acid residues from about 46 to about 218 of Figure 19 (SEQ ID NO:25).

23. The isolated IL-11p polypeptide of Claim 22 selected from the group consisting of: (1) an hIL-1Ra1L polypeptide comprising amino acid residues from about 1 to about 207 of Figure 15 (SEQ ID NO:19); (2) an hIL-1Ra1S polypeptide comprising amino acid residues from about 1 to about 167 of Figure 16 (SEQ ID NO:21); and (3) an hIL-1Ra1V polypeptide comprising amino acid residues from about 1 to about 218 of Figure 19 (SEQ ID NO:25).

24. The isolated polypeptide of Claim 22 selected from the group consisting of: (1) an hIL-1Ra3 polypeptide comprising amino acid residues from about 34 to about 155 of Figure 7 (SEQ ID NO:13); and (2) a mL-1Ra3 polypeptide comprising amino acid residues from about 34 to about 155 of Figure 9 (SEQ ID NO:16).

25. The polypeptide encoded by the DNA molecule of (a) in claim 14.

26. The IL-11p polypeptide of Claim 22 that comprises a native amino acid sequence of the IL-11p fused at its C-terminus or N-terminus to a heterologous amino acid sequence.

27. The IL-11p polypeptide of Claim 26, wherein said heterologous amino acid sequence is an epitope tag sequence.

28. The IL-11p polypeptide of Claim 26, wherein said heterologous amino acid sequence is a Fc region of an immunoglobulin.

29. An antibody which specifically binds to the IL-11p polypeptide of claim 22.

30. The antibody of Claim 29, wherein said antibody is a monoclonal antibody.

206120*99569960

1 GGCACGAGG AAGCTTCCA GGTATCTG AGCACTTG AAGTCTGAG AGCTACTGCC CTACAGAAAG TACTAGTCC CCTAAGAGTG GCCTGCGAC
CCGTGCTCC TTGGAGAGT CCAATAGAC TGGTGAGAC TTTCAGACTC TCGATGAGG GATGTCTTTC ATATGACAG GGAATTCGAC GCGACCGTG
*start insert

101 TGATGTTACT CTGCTGTG GAGTACAAC TCCATTAGA AACACTGC CAGCACTTA AGCACTCA CACCTCAGA TTGGCTTGA GAAGATTTG
ACTACATAGA CGACGACAC CTCATGTGA AGGATATCT TTGTGTGAG GTCTGGAT TCTGTGAGT GTGGAGTCT CACGGAAT CTCTTCAAC
1 M L L L L L E Y N F P I E N N C Q H L K T I H T F R *possible splice donor

201 GGTGCAAGG TCATGCGA GACACCACT TAAGAGATA GTGAATAGT CTGCATGTGA GAGCTGAGA TCCATATGCA GCTGTGATA GAGGGAAGC
CCGAGTTCTT AGTACTGCT CTGTGGTGA ATTCTCTH CACTGATCA GAGCTACT CTGCACTCT AGGATACAT CCGACTAT CTCTCCCTTG
301 AGAAACAAA GGAAGACA GCTTAAGAA GCGTTAAGA GCAACCCAC CATCTGAC AGTCACTGC CCAGCTGGG GGCCTCTGT CTTATCAAA
TCTTTGGTT CTCTCTCT CGGAATCT CGGTGGTG GTAGAGCTG TCAGTGACG GGTCCGACC CCGGGACAA GAATAGTTT

401 CAGTGCCTG AGCTCTTTC AGAGTCCAA AGGTGAGAA CTAAACCG AAGAAATCA GCATTCATGA CCAGATCAC AAGTACTGG TCTTGACTC
V K N L N P K K F S I H D Q H K V L V L D S
1 GTTACGAC TCAGAGAGG TCTCAGGT TCACTTCTT GAATTTGGC TTCTTTAAGT GGTAACTACT GTTCTAGTG TTCTAGTACC AGCACTGAG
*85066..AH1282..Asc.f, 5'Tag: TTTCCTTTGGCGCCG
*85066..AH1284..Nsi.f, 5'Tag: TTTCCTTTGATCATGATGAGCATGACAA
*possible splice acceptor
*ORF

501 TGGGAATCTC ATAGAGTTC CAGATAAAA CTCATACGC CCAGAGATCT TCTTGATTT AGCCTATCC TTGACTCAG CCTGTCGGA GAGAGGAAGT
AGCCTTAGAG TATCTCAG GTCTATTTT GATGATGCG GGTCTTAGA AGAAGCTAA TCGAGTAGG AACTGAGT GAGAGGCT CTTCCTTCA
24 G N L I A V P D K N Y I R P E I F A L A S S L S A S A E K G S

601 CCGATCTCC TTGGGCTTC TAAGGGGAG TTGTGTCT ACTGTGACA GATTAAGA CAAATCATC CATCCCTTCA GCTGAGAG GAGAATGA
GGCTTAGAG ACCCTTCC ATTTCCCTC AATACAGA TGACATTT CTATTTCT GTTTCATAG GTAGGAAGT CAGCTTTC CTCTTGAT
57 P I L L G V S K G E F C L Y C D K D K G Q S H P S L Q L K K E K L M

Fig. 1A

206120' 99569860

701 TGAAGCTGGC TGCCCAAG GATCAGC GCGGCCCTT CATCTTTAT AGGCTCAG TGAGCTCTCTG GATCAGCTG GATCGCGCG CTCACCCCG
 ACTCGACG AGGGTTTC CTGAGCTG CGCGGGGA GTAGAMATA TCCGAGTCC ACCGAGGAC CTGTAGAGC CTCAGCGCC GATGCGGCC
 91 K L A A Q K E S A R K F I F Y R A Q V G S W N M L E S A A H P G
 801 ATGGTTCATC TGCACCTCTT GCAATTGTAA TGAGCTGTT GGGTGACG ATTAATTGA GACAGAAA CACATTGAAT TTTCATTTC ACCAGTTTC
 TACCAGTAG AGTGAGGA CGTACACTT ACTGAGCAA CCCACGTGCTT TATTTAGCT CTGACCTTT GTGTAACTTA AAGTAAAGT TGTCAAACG
 124 W F I C T S C N C N E P V G V T D K F E N R K H I E F S F Q P V C
 901 AAGCTGAA TGAGCCCCAG TGAGGTACG GATTAGAAA CTGCCCCAT GACGCTTC CTCGTAAT TTGACTAAT GTATAAAG ACCAACCTG
 TTTCGACTT ACTCGGCTC ACTCAGTCG CTATCTTT GACGGGTAA CTGCGGAG GAGGATTA ACTGATTA CATTATTG TGTGTCGAC
 157 K A E M S P S E V S D O
 ~85066.AH1283.r
 ~85066.AH1285.Not.r. 5'Tag: TTTCCTTTGCGCGCTTA

1001 CTCCT
 GATGA

FIG 1B

206120:99569860

1 TAAATCAGCA TGTCTGCACT TCTGATCTTA GCTCTTGTG GAGCTGAGT TGTGTACTAG AAGACGATG AGCAAGACT TCCGCGCGCG TAATCAGCTC
 1 M S A L L I L A L V G A A V A D Y K D D D D K L A A A N S A L *insert st
 *orf
 101 TTTCGAGAG TCCAAAGGTG AAGAACTTAA ACCCGAGAA ATTCAACATT CATGACGAG ATCACAAAGT ACTGTCCTG GACTCTGGGA ATCTATAGC
 AATAGTGGT ACAGACTGA AGACTAGAT CGAGACAGC CTGACACTA AGACTGATG TTCTGTCTAC TGTCTTTGA AGCGCGGCGG TTAGTCTGAG
 32 C R G P K V K N L N P K K F S I H D Q D H K V L V L D S G N L I A
 201 AGTTCAGAT AAAACTACA TACCCACAGA GATCTCTTGT GATAGGCT CATCTTCG CTACGCTCT GCGAGAAAG GAATCCGAT TCTCTGCGG
 TCAAGCTCA TTITGATGT ATCGGGTCT CTAGAGAGAA CGTATCGA GTAGGACTC AGTCTGAGA CGCTCTTC CTTCAAGCTA AGAGACCCC
 65 V P D K N Y I R P E I F A L A S S L S A S A E K G S P I L L G
 301 GTCTCTAAG GGGATTTTG TCTCTACTGT GACAGAGATA AAGACAAAG TCATCCATCC CTTCAGCTGA AGAGAGAGAA ACTGATGAAG CTGCTGCGC
 CAGAGATTTC CCTCAAAAC AGAGATGACA CTGTCTCTAT TTCTGTGTTT AGTAGTGATC GAATCGACT TCTTCTCTTT TGACTACTTC GAGCGACGG
 98 V S K G E F C L Y C D K D K G Q S H P S L Q L K K E K L M K L A A Q
 401 AAAAGGATC AGCAGCGG CCTTCATCT TTTATAGGTC TCAGTGGGC TCTTGAACA TGTGTGATC GCGGCTCAC CCGGATGCT TCAATCTGAC
 TTCTGCTTAG TCTGCGGCG GGGAGATAGA AATATCCG AGTCCACCG AGGACTTGT AGACTCTAG CGGCGAGTG GGGCTTACCA AGTAGAGTG
 132 K E S A R R P F I F Y R A V G S W N M L E S A A H P G W F I C T
 501 CTCTGCAAT TGTATGAC CTGTGGGT GACAGATAA TTGAGAAC TTTGAGACA GGAACACAT TGAAITTTA TTTCACACG TTTCACAAAG TGAATGAGC
 GAGACGTTA ACATTAATCG GACAACCCA CTGTCTATT AAATCTTGT CTTTGTGTA ACTTAAAGT AAAGTTGTC AAAGTTTCG ACTTACTCG
 165 S C N C N E P V G V T D K F E N R K H I E F S F Q P V C K A E M S
 601 CCAGTGAGG TCAGCAATTA GGTACCACT GCACTTAGA GGAATCCGGG
 GGTACTTCC AGTCTAAT CCATAGCTA CTTGAGATCT CCTAGGCGC
 198 P S E V S D O *inserts ends

FIG. 2

206720-99569660

1 GGCAGAGC AGCCTTCA GGTATCTG AGCAGCTG AAGTCTGAG AGCTACTGCC CTACAGAAAG TTACTAGTGC CCTAAGCTG GCCTGGCAC
 CCTGCTCG ITCGGAAGT CCATGAGC TCGGTGAGC TTTCAGACTC TTGATGAGG GATGTCITTC ATGATCAGC GGAATTCGAC CGGACCGTG

 101 TGAATGACT GTGCTGTG GAGTCACT TCCTATGA AAGCACTGC CAGCAGCTTA AGACCACTCA CACCTTCAGA GTGAGAACT TAAACCGGA
 ACTACATGA CGACGACAC CTCATCTTGA AGGATATCT TTGTGTGAG GTCTGGAT TTGTTGATG GTGAGTGT GTGAGTGT CACTTCTGA ATTGGGCTT
 1 M L L L L E Y N F P I E N N C Q H L K T T H T F R V K N L N P K
 "Met

 201 GAAATTCAGC ATTATGACC AGGATCAAA AGTACTGGT CUGGACTCTG GGAATCTCAT AGCATCTTCA GATAAAACT ACATAGCGCC AGAGATCTTC
 34 K F S I H D Q D H K V L V L D S G N L I A V P D K N Y I R P E I F
 301 TTTCATTAG CTCATCCTT GAGCTACCC CTCGCGAGA AAGGAGTCC GATCTCTCTG GGGTCTCTA AAGGGAGTT TTGTCTTAC TTGTCAAGG
 AAACGTAATC GAGTAGGAA CTCGAGTCGG AGACCTCTT TTCTTCAGC CTGAGAGAC CCCAGAGAT TTCCTCTGAA ACAGAGATG AGCAGTTC
 67 F A L A S S L S A S A E K G S P I L L G V S K G E F C L Y C D K D
 401 ATAAAGACA AATCATCA TCCCTTCAGC TGAAGAAGA GAACTGATG AAGTGGTG CCCAAGA AATCAGACCC CGGCTCTCA TCTTTTATG
 TATTCTCTGT TTCTAGTGT AGGGAATCG ACTTCTCTCT CTTTGACTAC TTGCACCCAG GGTTTTCTCT TAGTGTGCG GCGGGAAGT AGAAATATC
 101 K G Q S H P S L Q L K K E K L M K L A A Q K E S A R R P F I F Y R
 501 GGTCTAGGTG GATCTCTGA ACATCTGA GTGGCGCT CACCCGGAT GGTTCATG CACCTCTCTC AATTGTAAG AGCTGTGG GTGACAGAT
 CCGATCCAC CCGAGACT TTACGACT CACCGCGA GTGGGCTTA CCAAGTAGAC GTGAGAGAGC TTAACATTAC TCGGACAACC CCACCTGTA
 134 A Q V G S W N M L E S A A H P G W F I C T S C N C N E P V G V T D
 601 AAATTTGAGA ACAGGAACA CATTGAATTT TCATTGAC CAGTTGCAA ACTGTGAATG AGCCCAAGT AGTCAAGCA TTAGAAACT GCACCATTA
 167 K F E N R K H I E F S F Q P V C K A E M S P S E V S D O
 701 AGCCTCTCTT CGCTAATTTG AACTAATGT ATAAAGAAC CAACCTGCT CACT
 TCGGGAAGA GCAATTAAC TTGATTAACA TATTTTGT GTTTGGACA GTGA

FIG. 3

2061207 99569800

1 CCAGGCCAA GNTCCCAAC CATGAATTTT GTTCACAA GTCCAAAGT GAGAGCTTA AACCGAAG AATTCAGAT TCATGACGAG GATCACAAAG
 GGTCCGGGTT GGNAGGGGTG GTACTTAAAC CAAGTGTTGT CAGCTTCCA CTTCCTGAT TTGGGTTCT TTAAGTCTA AGTACTGTC CTAGTGTTTC

 101 TACTGGCTG GACTCTGGG ATCTCATAGC AGTTCAGAT AAAAATCTACA TAGGCCNCA GATCTCTTTT GCAATGACT CATCTTTGAG CTCAGCCTCT
 ATACCCGAG CTGAGACCT TAGAGTATCG TCAAGTCTA TTTTGTATGT ATGCGGCTCT CTAGAAGAA CGTAATCGGA STAGAGATC GAGTCGGAGA

 201 CCGGAGAAAG GAGTCGGAT TCTCTGGGG CTCTCTAAG GGGATTTTG TCTCTACTGT GACAGAGATA AAGGACAAAG TCATCCATCC CTCAGCTGA
 CGCTCTTTC CTTCAGGCTA AAGAGACCC CAGAGATTC CCGTGAAC AGAGATGACA CTGTCTCTAT TTCTGTGTTT AGTAGGTAGG GAGTGCGACT
 1 I L L G V S K G E F C L Y C D K D K G Q S H P S L Q L K
 ^orf
 ^84664.p1

 301 AGAAGAGAA ACTGATGAG CTGGCTGCC AAAGGAATC AGCAGCCGG CCCTTCATCT TTTATGGGC TCAGTGGGC TCCTGGACA TGCTGAGTC
 TCTCTCTTT TGACTCTTC GAGCGAGGG TTTTCTTAG TGTGCGGCC GGGAGTNGA AAATCCCG AGTCCACCG AGACCTGTG ACAGCTCAG
 29 K E K L M K L A A Q K E S A R R P F I F Y R A Q V G S W N M L E S

 401 GGGGCTCAC CCGGATGGT TCATCTGCAC CTCTGCAAT TGTATGAGC CTGTGGGGT GACAGATAA TTGAGACA GGAACACAT TGAATTTCA
 CCGCGAGTG GGGCTACCA AGTAGACGTG GAGGACOTTA ACATTAATCG GACACCCCA CTGTCTATTT AAATCTGTG CCTTCTGTA ACTTAAAGT
 62 A A H P G W F I C T S C N C N E P V G V T D K F E N R K H I E F S

 501 TTTCACAG TTTCACAG TGAATGAGC CCGAGTGGG TCAGCATTA GGAATGCC CCATGAGCG CTTCTCTGC TAAATTGAAC TAAATTGATA
 AAAGTTGTC AAAGTTTCG ACTTTACTCG GGTCACTCC AGTGTCTAT CTTTGGCGG GSTRACTCG GGAAGAGCG ATTAACCTTG ATTAACATAT
 95 P Q P V C K A E M S P S E V S D O
 ^84664.r1

 601 AAAACCCCA ACTGCTCAC TAAABAAA
 TTTGGGGTT TGGAGGTG ATTTTTTT

FIG. 4

206120-99569860

1 GTGACACAC GGTCTCCAGC CTGCTGAGC CAGACTGAG TCCCTGGAG TGTGATGAA GACCTTCTT TCCAGGTGC TGAGACACC ACACATAGAG
 1 CACTGGTG CGAGCTTC GACACCTTC GTCTATGTC AGGAGACCTG ACATCAHTT CTGGGAAGA AGSETCCAGS ACTCTGTGTS TGTATGATC
 M R
 'MET
 92929.AH421.Asc.f. 5'Tag: AAGGGAAAGGCGCC'

101 AGGCACTCCA GAGACCTG ATGGTGAGG AAGGCGCTC TATCAATCA TCTCTGTGG TGTATCACA TGCAGTATC CAGAGCTCT TGAAGAGGC
 TCGTGAAGT CCTCTGGAC TACCACTCC TTCCGGCAG ATAGTAGT AGTCAACG ACATAGTGT AGTTCATAG GTTCCGAGA ACTGTCTCG
 3 G T P G D A D G G R A V Y Q S I T V A V I T C K Y P E A L E Q G

201 AGAGGGATC CCATTATTI GGAATCCAG ATCCAGAA TGTGTTGTA TTGAGAGG GTTGAAGC AGCCACAT GCACATAAA GAGCAGAGA
 TCTCCCTAG GGTAAATAA CCTTAGTC TTAGTCTTT AGCAACAT AACACTCTC CAACCTCTG TCGGTGTAA CGTGATTTT CTGTCTCTT
 36 R G D P I Y L G I Q N P E M C L Y C E K V G E Q P T L Q L K E Q K I

301 TGTATGACT GTATGCGA CCGACCCG TGAACCTT CCTTTCTAC CTGTCCAGA CTGTAGGAC CTCACCTT GAGTCTGG CCTTCCGGA
 AGTACTAGA CATACGCTT GGTCTGGC ACTTGGAA GGAAGATG GCAGGTCTT GACATCTG GAGTGGGA CTCAGACCC GGAAGCCT
 70 H D L Y G Q P E P V K P P L F Y R A K T G R T S T L E S V A F P D

401 CTGTCTCAT GCTCTCCA AGAGACCA GGCATCTT CTGACTCAG AACCTGGAA GTATACAC ACTGCTTGG AMTAAAT AATGACTGA
 GACCAATGA CGAGAGAT TCTCTCTGT CGGTGATA GACTGATC TTGACCTT CATTCTGT TGACGAACT TATATTATA TTACTACT
 103 W F I A S K R D Q P I I L T S E L G K S Y N T A F E L N I N D O
 *92929.AH1

501 ACTAGGCTA GAGTGGAG CTGTGCTT GCTATAGT TTCTGTTC CANTGTT TGTCTGAT TTCTATG TCAATTCAC GCTGTCTG
 TGAATGAT CTCACCTC GACCAAGA CAGATTCA AGACCAAG GTTACAAA AGCAGTGA AAGATACAG AGTAAAGTG CGACCAAG
 136

601 AGAGAGAG AGGTGCTG TTATCATC ATTATATAT GAGAGAG CATTACTTC ATGAGACTG AAGACAGA TGTGCTCA GAGCAGAG
 TCTGTCCTG TTCCAGAC ATATAGTG TAAATATA CTCTCTTC GTATGAG TATGCTGAC TTCTGTCT ACACGAGT CTTCTCTCTC

701 AGCTGGTGT PATAGGCTG TCTCTCAG CTGTGCTGT AGGACACA AGGACTGCT ATGCTACT TTAGCTCA AGACCAAC ACTGAGTTT
 TCAAGCAC ATATTCAG AGAGATC GACCAACA CTGCGTGT TCTGAGCC TACTACTGA ATTTGAGT TCTGTGTTG TGAATGAAA

FIG. 5A

801 CTTCTAGGGG TGGGTATGAA GATCTTCAG AGCTATGCG GATTACCCAC CTAGCACAGA SCTGATCTCT GTTCTGTGTT TGCCTTATTC
 GAGGATCCCG ACCCATCTT CTACGAGTC TCGAGTACGC SCATGGGGT CTACGGTACT GATCGTGCT GAACTAGAGA CAAAGACAAA ACCGAATATAG
 901 CCTCTTGGGA TGTATGCTAC CAGTCTTAT ATGTGCGAA TATACCTCAT TGTGTGTAT AGACCTTCT TAGCATTAAG ACCTTGTAAC CAAATATAT
 GAGAGACCTT ACTATGATAG GTGAGAAATA TACAGGGTT ATATGGAGTA ACACACATTA TCTTGGGAAG ATCGTAATTC TGGAACTATT GTTTTATTTA
 1001 TCTTGGGGTG GGTATGAGA TGGTTCGAG CTCATGCGG TTACCCAGA TGGCATGACT AGCAGAGCG TGATCTCTGT TTTCTGTTTT CTTTATTCOC
 AGAGCCCCAC CACTACTCT ACGAGTCTC GAGTACGCG AATGGGTGCT ACCGACTAGA TGGTGCTCG ACTAGAGACA AAGACAAAC GAAATAGGG
 1101 TCTTGGGATG ATATCTCCA GTCTTATAT GTTCCAAATA TACTCTATG TGTGTATAG AACCTCTTA SCATTAAGAC CTGTAAACA AAATATATTC
 AGAGCCCTAC TTATGTAGT CAGAAATTA CACGGTAT ATGGAGTAC ACAGATATTC TTGGAGAT COTATCTG GAACNTTGT TTTTATTAAG
 1201 TTGTGTTAAG TTAATCTTT TTGTGCTTAA TTGTATATG TATCTTAA GTTATATTA CTGTGTGAT TATATATA ATTAAGCTAA AACGATATA
 AACATATTC AATTAGTAA AACAGGAT AACATATAC ATATGATTT CATTATTTT GAAACATATA AATATATAT TATTTGATTT TTGACTATAT
 1301 AAATTAAGAA AGATTAAGT G
 TTTATTTCTT TCTCATTTGA C

FIG. 5B

206120-9559980

1 AAGCTGCTGG AGCCACGATT CAGTCCCTCG GACTGTAGAT AAAGACCTTT TCTTGCCAGG TGCTGAGACA ACCACTAT GAGAGCACT CCAGGAGACG
TTGAGAGACC TCGGTGCTAA GTCAAGGGAC CTGACAATCTA TTTCGTGGAA AGAACGCTCC ACGACTCTGT TGGTGTGATA CTCTCGGTGA GGTCTCTTGC

101 CTGATGCTGG AGGAGGGGCC GTCTATCAAT CAACTACTGT TGCTGTTATC ACATGCAAGT ATCCAGAGGC TCTTGAGCAA GGCAGAGGGG ATCCCAATTA
GACTACCAAC TCCTTCCCGG CAGATAGTGA GTTAGTGACA ACGACAATAG TGTACTTCA TGGTCTCCG AGAATCGTT CCGTCTCCC TGGGTAAAT

201 TTGCGGAATC CAGAATCCAG AAATGTGTTT GTATGTGAG AAGTTTGA
AAACCTTTAG GTCTTAGTTC TTACACAA CATACACTC TTCCAACCT

FIG. 6

206720 9955950

1 ATGCTCTGA GTGGGCGT GTGCTTCGA ATGAGGACT CGGCAITGAA GGTGCTTTAT CTGCATATA ACCAGTTCT AGCTGGAGG GTGCTGAG
 TACCAGACT CACCCGCGCA CACGAGGCT TACTCTCTGA GCGTAACCTT CCACGAATA TGTGGAAGA TCGACTCCC GAGCTAGCTC
 1 M V L S G A L C F R M K D S A L K V L Y L H N N Q L L A G G L H A G
 101 GGAAGTGCAT TAAAGTGA GAGATCAGC TGTTCGCCAA TGTGTGCTG GATGCCAGCC TGTCCCGCT CATCTGGGT GTCCAGGCTG GAAGCCAGTG
 35 K V I K G E E I S V V P N R W L D A S L S P V I L G V Q G G S Q C
 201 CTTGTCTATGT GGGTGGGCG TCTTACTA TCTTACTA GATCATGGA GCTCTATCTT GTGCCAAGG ATCCAGAG GTTCACTTC
 GAGCAGTACA CCCCACCCG TCTCTGGCTG AGATTGTGAT CTGGTCACT TGTGTACTT CGGATAGAA CACGTTCC TTAGTCTTC GAGTGGAG
 68 L S C G V G Q E P T L T L E P V N I M E L Y L G A R E S K S F T F
 301 TACCGGCGG ACATGGGCT CACCTCCAGC TTGAGTCGG CTGCTTACC GGGCTGGTC CTGTGACGG TGCCTGAGC CGATCAGCT GTGAGACTCA
 ATGGGCGCC TGTACCCCGA GTGAGGTGCG AAGCTCAGCC GACGATGGG CCGCACCIAG GACAGGTGCC AGGACTTGG GCTAGTCGGA CAGTCTGAGT
 101 Y R R D M G L T S S F E S A A Y P G W F L C T V P E A D Q P V R L T
 401 CCGAGCTTC CGAGATGGT GGTGGAATG CCCCATCAC AGACTTCTAC TTCCAGCAGT GTGACTAG
 GGTGCAAGG GCTCTTACA CCGACTTAC GGGGTAGTG TCTGAGATG AAGGTGTCA CACTGATC
 135 Q L P E N G G W N A P I T D F Y F Q Q C D O

FIG. 7

206120-9956960

1 GCTCCCGCA GAGAAAGGA ACATCTGAG GGGAGCTAC ACCTGTGGA GTCTGAATG GTCTGAATG GGGCGCTGTG CTTCGGAATG AAGGACTGG
 GAGGGCGGT CCTCTTCTT TGTAGATC CCTGAGATG TGGGACACT CGAGTTCTAC CAGGACTAC CCGCGACAC GAAGGTTAC TTCTGAGCC
 1 A P A R R K E H S E G S L H P V E L K M V L S G A L C F R M K D S A
 101 CATCGAGGT GCTTATCTG CATATATAC AGCTTCTAG TGGAGGCTG CATGCAAGGA AGTCATTA AAGTGAAG ATCAGCGTGG TCCCAATCG
 GFLCTTCCA GGAATAGAC GTATATGG TCGAGATCG ACCTCCGAC GTAGTCCCT TCCGTAAT TCACCTTCTC TATCGCACCC AGGGTTAGC
 35 L K V L Y L M N N Q L L A G G L H A G K V I K G E E I S V V P N R
 201 GTGGTGAAT GCAGGCTGT CCCCCTCAT CTGGGTGTC GAGGTGGA GCGAGTCCCT GTGATGSG GTGGGCGAG AGNCACTCT AGAT
 CACCGACCTA GGTGGACA GGGGCGATA GACCCAGG GTCCACTT CAGTACGGA CAGTACACC CACCCGTC TCAGTAGA TTGTA
 68 W L D A S L S P V I L G V Q G G S Q C L S C G V G Q E X T L I

FIG. 8

206120-59569860

1 ATAGGAATTT TGGCCCTCGA GGCACAGAT TGGGACGAG GGGAGCTGC TTCTACTTA GTCTCAAT TTTCAGCCT TGCTTTGCC TAAATTTCC
 TTATCCCTTAA ACCGGAGCT CCGTTCTTA ACCCTGCTC CCTCGGACG MAGTGAAT CCAGATTTA AAAGTCGA ACAGAAAGG ATTTTAAGG

101 TGCTGTATT TTCAAAATAG GGTCTACATA CTGTGGAGT CATGATGTT CTGATGGGG CATATGCTT CGATGCGAG GATTCAGCCT TGAGGTACT
 ACAGACAATA AAGTTTATC CCAGATGTAT GACACTCGA GTACTACCA GACTACCCC GTATAGGA GGTCTACTT CTAAGTCGA ACTTTCATGA
 M V L S G A L C F R M K D S A L K V L
 ^orf

201 GTATCTGAC AATAACGAC TGCTGGCTGG AGGACTGAC GCAGAGAAG TCATTAAAG TGAAGAGATC AGTGTGTCC CAATCGGGC ACTGGATGCC
 CATAGACTGG TTAATGTCG AGCAAGGACC TCCTGAGGTG CGTCTTTCC AGTAAATTTCC ACTCTCTTAG TCACACAGG GTTTAGCCCG GTTACTCAGG
 20 Y L H N N Q L L A G G L H A E K V I K G E I S V V P N R A L D A

301 AGTCTGTCCC CTGTCTCTCT GGGCTTCTA GGAAGAGCC AGTGGCTATC TTGTGGACA GAGAAAGGSC CAATTCGAA TGAATGACCA GTGACATCA
 TCAGACAGG GACGATGGA CCCCAAGT CTCTCTTGG TCAGGATAG AACACCTGT CTCTTTCCG GTTAGACTT TGAATCTGGT CACTGTGAT
 53 S L S P V I L G V Q G S Q C L S C G T E K G P I L K L E P V N I M

401 TGGAGCTCTA CCTCGGGCC AAGAAATCAA AGAGTTTAC CTCTACCG CGGATATGG GTCTTACTTC CAGCTTGAA TCCCTGTCT ACCAGGCTG
 ACCTCGAGT GGAGCCCCG TTCTCTTAGT TCTCGAGTG GAAGATGCC GGCCTATACC CAGAAATGGAG GTGAGACTT AGGGACGGA TGGGTCCGAC
 87 E L Y L G A K E S K S F T F Y R R D M G L T S F E S A A Y P G W

501 GTTCTCTTGC ACTTACCGC AAGCTGACA GCTCTGAGG CTCACTCAGA TCCCTGAGGA CCCGCTGGG GATGCTCCA TCACAGACTT CTACTTTTAC
 CAGAGAGAC TGGATGGCC TTCACTGCTT CGACATTC GAGTGAATCT AGGACTCTT GGGGGGACC CTACAGGGT AGTCTGTGA GATGAAATC
 120 F L C T S P E A D Q P V R L T Q I F E D P A W D A P I T D F Y F Q

601 CAGTGTGACT AGGGCTGCT GTTCCCAAA ACTCCNTRAG CAGAGCGA GTAGCGTG CGGCTCTCG ATGAGGNTA GAGAGACGA GAGAGCTCCAC
 GTCAACATGA TCCGAGGCA CCAGGGGTTT TGAATATTC GTTCGGTCT CATCCGTAC CCGCGAGGAC TATCTCTAT CTCTGTGT CCTCGAGGTG
 153 Q C D O

701 AGTAGTGGC TPACTGCTCT CTTCTCTAC TGAGCTCCG CTTCGACCT AAGGACACA GACACTCTT TCTCTGCTT CCGAGTGTG GTAAATTTTC
 TCATCCACG ATGAGAGGA GAGAGGATG ACCTGAGGSC GAAGACTGGA TTCCGTGTGT CTGTGAGAGA AGAGAGCTA GGGTCAAGC CATTTAGAG

FIG. 9A

206120' 99568880

801 TGGTATTTCG AGCTCAGCTT GTAGATCTCT TCGATTGGA TGGTACTACC TCTGTGTGGS AACCAATAG AGACCAAGTA GGACCAACAA AGACCAACAT
 ACCATTAACC TCGATTACCA CACTTAAGAA AGCTTAACCT ACCGATGGS AGACCAACCC TTGGSTATTG CTTGGTGCAT CCGTGTGTTT TCTGTGTGTA

901 AAAAGATTCT TGGGTGAGAA AGAGGTGGGA ACTGTTCTTA CAGCTTAGA TCTGACAG TACTCTCAGA GTCTTGCCAT TCGTTATGTT CTGGAGAAAG
 TTTTCTNAGA ACCCACTTCT TCTCAGCTT TGCAGATAT GTACATCT AGCTGTGTC ATGGATCTT CAGCAAGTA AGGAATACAA GACCTCTTTC

1001 TGGAGGGGGG GTCCACAGA CTCTCTGGS CTGCTGGGC CTTTCCCTC AACCTTCTG ACATCTCGS CCTCTCTAT TCTTGGCTTC ATTCTCTGAC
 ACTTCCCCC CAGTGTCTT GAAAGAGACC GACCAACCG GGAAGGGAG TTGGAAGAC TGTAGCTC GGAAGGATG AGAACGGAG TAAAGACCG

1101 CCTGAACCGA GAGGTGATA TCGAGTATAC TGACAGAGA TGACCAGGA CACTGTCTG GTTGAACC AGAGGGACA ATTAATAACC CTGATCTTGG
 GGACTTGGCT CTCCCATAT ATCTCTATG ACTGTCTCT ACTGTCTCT GTACAGAGC CAACTTGG TCTCCCTCT TATTTTGG GACTAGACC

1201 TCTCTACTCA CATATAAGA AGCTTGTAAG CATTAAGTGG GAAGATTTG CTACTAATA ACATCTTGS TAATTCATC TTAATTAAAA TATCTCTTC
 AGAGATGAGT GTATTTTCT TCGACACTT GTATTCACC CTCTCTAAC GATGATTTAT TGTATGAAAC ATTAATGATG AATTAATTTT AATGGAAG

1301 TATATTATAT ATTTTAAAA AAAAAAAA AAAAAAAA AAAAAAAA AAAAAATGC GGCCTAAGC TTATTCCTT TAGGA
 AATATATATA TAAATTTT TTTTTTTT TTTTTTTT TTTTTTTT TTTTGTAGC CCGCTTGC AATAGATTA ATCT

*insert ends

FIG. 9B

206120: 99569660

1 GGAGCGTCTT TCTACTTAG GTCTCAATT TTCCAGCTT GTCTTGCTT AATATTCCT GCTGTTTAT TCATAATAG GTCTACATAC TGTGAGCTC
 CCTCGGACG AAGATGATC AGATGATC TCCAGCTT TCCAGCTT AATATTCCT GCTGTTTAT TCATAATAG GTCTACATAC TGTGAGCTC
 101 ATGATGCTTC TGAATGAGG ACTATGCTC GGAATGAGG ATTGAGCTT GAGTACTG TATCTGACA ATAACCACT GCTGCTCGA GGAATGACG
 TACTACCAAG ACTACCCCG TGTAGAGAG GCTTACTTC TAAGTGGAA CTTCATGAC ATGAGCTGT TATTGTGGA GACCGAATC CCTGACGTG
 1 M V L S G A L C F R M K D S A L K V L Y L H N N Q L L A G G L H A
 'orf
 201 CAGAGAAGT CATTAAGT GAGGATCA GTGTTGTCC AATCGGCA CTGATGCGA GTCTGTCCC TGTATCTTG GGTTCACG GAGAAGCA
 GTCTGTCCA GTATTTCCA CTCTCTAGT CACAACAGG TTGAGCCGT GACTACCGT CAGACAGG ACAGTAGAC CGCAAGTTC CTCTCTCGT
 35 E K V I K G E E I S V V P N R A L D A S L S P V I L G V Q G G S Q
 301 GTGCTATCT TGTGAGCAG AGAAGGCC AATCTGAA CTTGAGCAG TGAACATCAT GAGCTCTAC CTCGAGGCA AG
 CAGGATAG ACACCTGTC TGTTCGCG TTGAGCTTT GACTGTGTC ATTGTGATTA CTTCGAGATG GAGCCCGT TC
 68 C L S C G T E K G P I L K L E P V N I M E L Y L G A K

FIG. 10

hIL1Ra 1 MEICRGRLRSHLITLLLFHSETICR[S]GRKSSKMGAFRIW[D]VNQKTFYL
hIL1Rabeta 1MRGTPGDA[D]GGRAVYQSMCK[P]ITG
tango77MGSEDWEKDEPCCLEDPA[G]SPLEP
hIL1Ra1S 1MGSEDWEKDEPCCLEDPA[V]SPLEP
hIL1Ra1L 1MGSEDWEKDEPCCLEDPA[V]SPLEP
hIL1Ra1V 1MSFVGENSGVKMGSEDWEKDEPCCLEDPA[G]SPLEP
hIL1Ra2 1MRGTPGDA[D]GGG.....
hILRa3 1MVLSGALCFRMK[D]S[L]KVLYL

hIL1Ra 51 RNN.....QLV[AGY]LG
hIL1Rabeta 26 TIN.....DLNQGVVTLGGONLVA[P]RSDSV
tango77 26 [G]P.....SLPTMNFV
hIL1Ra1S 26 [G]P.....SLPTMNFV
hIL1Ra1L 26 [G]PSLPAMNFVHTSPKVKNLNPKKFSIHDQDHKVLVDSGNLIA[V]PDKNYI
hIL1Ra1V 37 [G]PSLPAMNFVHTSPKVKNLNPKKFSIHDQDHKVLVDSGNLIA[V]PDKNYI
hILRa2 13RAVYQS..
hILRa3 22 HNN.....QLLAGGLHA


hIL1Ra 63 PNVNLEEKIDV[P]F.....IEPHALF[GIN]G[KM]C[L]S[C]V[K]SGDETRIQ[LE]A
hIL1Rabeta 52 TPVTVAVITCKY[P]E[AL]EQGR[G]P[I]Y[LG]IONP[E]MCLYCE[KV]GE[G].....P[IT]L
tango77 34 HTK[IFF]ALASSLSSASAE[K]GSPILLGVSKGEFCLYCDKDKGOSHPSLQL
hIL1Ra1S 34 HTK[IFF]ALASSLSSASAE[K]GSPILLGVSKGEFCLYCDKDKGOSHPSLQL
hIL1Ra1L 76 RPE[IFF]ALASSLSSASAE[K]GSPILLGVSKGEFCLYCDKDKGOSHPSLQL
hIL1Ra1V 87 RPE[IFF]ALASSLSSASAE[K]GSPILLGVSKGEFCLYCDKDKGOSHPSLQL
hILRa2 15ITVAVITCKY[P]E[AL]EQGR[G]P[I]Y[LG]IONP[E]MCLYCE[KV]GE[G].....P[IT]L
hILRa3 34 GKVL[K]GEEI[S]VVP[PN]RWLDASL[S]P[V]I[LG]V[GG]S[GC]L[S]C[GV]G.....DEPTLT[L]EP

hIL1Ra 108 VNITOLSENRRKOD..K[R]FAFIR..SDSG[P]TITSF[ES]AAC[P]GWF[L]CTAMEA[D]
hIL1Rabeta 100 [K]E[Q]K[L]M[b]L[Y]G[G]P[E]..P[V]K[P]L[F]YRA[K]T[G]R[IT]S[TL]ES[V]A[F]P[D]WFI[AS]S[K]..R[D]
tango77 85 KKEKLMKLAACKESARRPFI[F]YRAQVGSWNMLESAAHPGWFICTSCNCNE
hIL1Ra1S 85 KKEKLMKLAACKESARRPFI[F]YRAQVGSWNMLESAAHPGWFICTSCNCNE
hIL1Ra1L 125 KKEKLMKLAACKESARRPFI[F]YRAQVGSWNMLESAAHPGWFICTSCNCNE
hIL1Ra1V 136 KKEKLMKLAACKESARRPFI[F]YRAQVGSWNMLESAAHPGWFICTSCNCNE
hILRa2 85 [K]E[Q]K[L]M[b]L[Y]G[G]P[E]..P[V]K[P]L[F]YRA[K]T[G]R[IT]S[TL]ES[V]A[F]P[D]WFI[AS]S[K]..R[D]
hILRa3 83 VNIMELYL[G]AKES..KSFTFYR..RDM[G]LTS[SF]ESAAY[P]GWF[ICT]VPEAD[D]

hIL1Ra 155 [P]V[S]LT[N]MPDEG...VMVTK[P]Y[F]Q[ED]E.....
hIL1Rabeta 148 [P]I[L]TSELGKS...YNT[A]FELNIND.....
tango77 135 P[V]GVTDKFENR...KHIEFSFOPVCKAEMSPSEVSD
hIL1Ra1S 135 P[V]GVTDKFENR...KHIEFSFOPVCKAEMSPSEVSD
hIL1Ra1L 175 P[V]GVTDKFENR...KHIEFSFOPVCKAEMSPSEVSD
hIL1Ra1V 186 P[V]GVTDKFENR...KHIEFSFOPVCKAEMSPSEVSD
hILRa2 113 [P]I[L]TSELGKS...YNT[A]FELNIND.....
hILRa3 130 [P]V[R]L[QL]P[EN]GGWNAPI[D]G[Y]F[Q]OCD.....

Fig. 12

A



B





FIG. 11






FIG. 13A




FIG. 14

206120 99559860

1 AAATGGGCT CTGAGCATG GGAAGAAT GAGCCCGCT GCTCTTAGA AGACCCGCT GTAGCCAGG CCCAGCCTC CCGGCATGA
 TTTTACCGA, GACTCTGAC CTTTCTTCTA CTGCGGTCA GAGCAATCT TCTGGGCGA CATTCGGGG ACCTTGTCC GGGTTCGAG GGGCGTACT
 1 M G S E D W E K D E P Q C C L E D P A V S P L E P G P S L P A M N
 *insert starts
 *orf

101 ATTTTGTCA CACAGTCCA AAGTGAGA ACTTAACCC GAGAAATTC AGATTTCAG ACCAGATCA CAAATCTAG GTCTGGACT CTGGGAATCT
 TAAACAAAGT GTTCACTTCT TGAATTGGG CTTCTTTAAG TGTAAATGAC TGTGCTTAGT GTTTCATGAC CAGGACCTGA GACCTTAGA
 34 F V H T S P K V K N L N P K K F S I H D Q D H K V L V L D S G N L
 201 CATGACGTT CCAGATAAA ACTACATAG CCGAGATC TTCTTTGCAT TAGCCTCATC CTTGAGCTCA GCCTCTGGG AGAAGGAAG TCGGATTCG
 GTATTCGTCAA GGTCTATTIT TGTATGTC GGTCTCTAG AAGAAGCTA ATCGGAGTAG GACTCGAGT CCGAGAGCC TCTTCTTC TCAGCTAGAG
 67 I A V P D K N Y I R P E I F F A L A S S L S S A S A E K G S P I L
 301 CTGGGGTCT CTAAGGGGA GTTTTCTCT TACTGTGACA AGATAAAG ACAAGATCAT CCATCCCTTC AGCTAAGAA GGAGAAATG ATGAGATGG
 GACCCCAAGA GATTTCCCTT CAAACAGAG ATGACACTGT TCCATTTCC TGTTCAGTA GGTAGGGAAG TCGACTTCTT CCTCTTGGAC TACTTCGAC
 100 L G V S K G E F C L Y C D K D K G Q S H P S L Q L K K E K L M K L A
 401 CTGCGCAAAA GGAATCACA CCGCGCCTT TCATCTTTTA TAGGGCTCAG GTGGCTCTT GGAACATGCT GGAATCGGG GTCAACCCG GATGGTTCAT
 GACGGGTITT CTTTACTGCT GCGCCGGGA AGTAGAAAT ATCCCGAGTC CACCCAGCA CTTGTACGA CCTCAGCCG CAGGTGGGG CTACCAAGA
 134 A Q K E S A R R P F I F Y R A Q V G S W N M L E S A A H P G W F I
 501 CTGACCTCC TCAATTGTA ATGAGCTGT TGGGTGACA GATTAATTG AGACAGGA ACACATTGA TTTTCATTTT AACAGTTTG CAAAGCTGA
 GACTGTGAG AGTTAATCAT TACTTCGACA ACCCACTGT CTAATTAAAC TCTGTCTCTT TGTGTACTT AAAGTAAAG TTGCTAAC GTTTCGACTT
 167 C T S C N C N E P V G V T D K F E N R K H I E F S F Q P V C K A E
 601 ATGAGCCCA GTGAGGTGAC GATTA
 TACTCGGGGT GACTCCATC GATTA
 200 M S P S E V S D

FIG. 15

206120-99569860

1 AAATGGGCT CTGAGGACTG GGAAAAGAT GAAACCCAGT GCTGCTTAGA AGACCCGGCT GTAAGCCGCC TGGAACCCAGG CCGAGGCTTC CCGCCATGGA
 TTATTACCCGA GACTCCCTGAC CCTTTTCTA CTGGGGTCA CGAGGATCT TCTGGGCCGA CATTCGGGGG ACCTTGTCTC GGGTTCGGAG GGGCGTACT
 1 M G S E D W E K D E P Q C C L E D P A V S P L E P G P S L P A M N
 *insert starts
 .orf
 101 ATTTTGTCA CACAAGATC TTCTTTGCAT TAGCTCATC CTTGAGTCA GCTCTGGG AGAAGGAG TCCGATCTTC CTGGGGTCT CTAAAGGGA
 TAAACAAGT GTGTTTCTAG AAGAAAGTA ATCGAGTAG GAATCGAGT CGAGAGCCG TCTTTCTTTC AGCTTAAGAG GACCCCGAGA GATTTCCCT
 34 F V H T K I F F A L A S S L S S A S A E K G S P I L L G V S K G E
 201 GTTTTCTCT TACTGTGACA AGGTAAGG ACAAGTAT CCATCCCTTC AGCTGAAGA GAGAAGACTG ATGAAGCTG CTGCCCAAA GGATCAGCA
 CAAACAAGT ATGCACTGT TCTATTTC TGTTCAGTA GGTAGGAG TCGACTTCTT CCTTTTGAC TACTTGCAC GACGGTTT CCTTAGTCT
 67 F C L Y C D K D K G Q S H P S L Q L K K E K L M K L A A Q K E S A
 301 CGCCGCCCT TCATCTTTTA TAGGGCTCAG GTGGCTCTT GGAATGCT GAGTCTGGG GCTCACCCG GATGTTCT ATGCACTCC TSCAATTGA
 CGGCGCGGA AGTAGAAT ATCCGAGTC CACCGAGGA CCTGTGAGA CCTCAAGC CGAGTGGGC CTACCAAGTA GACTGGAGG AGTTAACAT
 100 R R P F I F Y R A Q V G S W N M L E S A A H P G W F I C T S C N C N
 401 ATGAGCTGT TGGGTGACA GATAATTG AGACAGGA ACACATTGA TTTCATTC AACAGTTG AACAGTGA ATGAGCCCA GTGAGTCA
 TACTCGACA ACCCACTGT CTATTAACT TCTGTCTT TGTGACTT AAAGTAAAG TTGTCAAC GTTCTGACTT TACTCGGGT CACTCAAGT
 134 E P V G V T D K F E N R K H I E F S F Q P V C K A E M S P S E V S
 501 CGATTA
 GCTAAT
 167 D

FIG. 16

1 AACCGAAGA AATTCACGAT TCATGACCAG GATCACAAG TACTGTCTCT GGACTCTGGG AATCTCATG CAGTTCCAGA TAAACTAC ATACGCCGAG
 TTGGCTTCT TTAAGTCGTA AGTACTGTC CTAGTGTTTC ATGACCAAGA CTTGAGACCC TTGAGTATC GTCAAGGTCT ATTTTGATG TATGCGGTC
 1 N P K K F S I H D Q D H K V L V L D S G N L I A V P D K N Y I R P E
 101 AGATCTCTT TGCATTAGC TCATCCTTGA GCTCAGCTC TCGGAGAA GGAAGTCCGA TTCTCTGGG GGTCTCTTAA GGGAGTTTT GTCTTACTG
 TCTAGAGAA AGTAATTCG AGTACTGTC CGAGTGGAG AGGCTCTTT CTCTCAGGT AGAGGACCC CAGAGATTT CCCTCAAA CAGAGATGAC
 35 I F F A L A S S L S S A S A E K G S P I L L G V S K G E F C L Y C
 201 TGCACAGGT AAGGACAMA GTCATCATC CCTTCAGCTG AAGAAGSAGA AACTGATGAA GCTGCTGTC CAAAGGAAT CAGCAGCCG GCCTTCATC
 ACTGTCTCTA TTCTCTGTTT CAGTAGGTAG GGAAGTCGAC TTCTTCTCT TTGACTACTT CGAGCGACGG GTTTTCTTA GTCTGCGGC CGGAGTAGT
 68 D K D K Q S H P S L Q L K K E K L M K L A A Q K E S A R R P F I
 301 TTTTATAGG CTCAGGTGG CTCCTGGAAC ATGCTGGAGT CCGCGCTCA CCGCGAGTG TTCACTGCA CCTCTGCAA TTGTATGAG CTGTGTGGG
 AAAATATCC GAGTCCACCC GAGGACCTTG TAGGACTCA GCGCGGAGT GCGGCTTACC AAGTAGACT GGAGAGCTT AACATTACT GCACACCC
 101 F Y R A Q V G S W N M L E S A A H P G W F I C T S C N C N E P V G V
 401 TGACAGATAA ATTTGAGAC AGGAACACA TTGAATTTT ATTTCACCA GTTTGCAAG CTGAATGAG CCCAGTGAG GTCAAGGATT AGMAACTGC
 ACTGTCTATT TAAACTCTTG TCTTTTGTT ACTTAAAG TAAAGTTGTT CAAAGTTTCT GACTTACTT CCGGTCACTC CAGTCGCTAA TCTTTTGAAG
 135 T D K P E N R K H I E F S F Q P V C K A E N S P S E V S D O
 501 CCCATTGAAC GCCTTCTCG CTAATTGAA CTAATTGTAT AAAACACCA TAAGTCTCA C
 GGGTAACTTG CGGAGGAGC GATTAACTT GATTACATA TTTTGTGTT TTAGCAGAT G

FIG. 17

```

hILiRa1      1  MSFVGENSEGVKMGSEDEKDEPQCCLDPAGSPLEPGPSLPTMNFVHTSP  M
hILiRa1L    1  MSFVGENSEGVKMGSEDEKDEPQCCLDPAGSPLEPGPSLPTMNFVHTSP  M
hILiRa1S    1  MSFVGENSEGVKMGSEDEKDEPQCCLDPAGSPLEPGPSLPTMNFVHTSP  M

hILiRa1      1  VKNLNPKKFSIHODDHKKVLVLDSGN
hILiRa1L    2  GSEDEKDEKDEPQCCLDPAGSPLEPGPSLPTMNFVHTSPKVKLNLPKKFSI
hILiRa1V    51  VKNLNPKKFSIHODDHKKVLVLDSGN
hILiRa1S    2  GSEDEKDEKDEPQCCLDPAGSPLEPGPSLPTMNFVHTSPKVKLNLPKKFSI

hILiRa1      26  LIAVPDKNYIRPEIFFALASSLSSASAEKGPSILLG
hILiRa1L    56  HODDHKKVLVLDSGNLIAVPDKNYIRPEIFFALASSLSSASAEKGPSILLG
hILiRa1V    77  LIAVPDKNYIRPEIFFALASSLSSASAEKGPSILLG
hILiRa1S    28  SLPAMNFVHTKLIFFALASSLSSASAEKGPSILLG

hILiRa1      62  VSKGEFCLYCDKDKGQSHPSLQKKLKKMLAAQKESARRPIFYRAQVG
hILiRa1L    102  VSKGEFCLYCDKDKGQSHPSLQKKLKKMLAAQKESARRPIFYRAQVG
hILiRa1V    113  VSKGEFCLYCDKDKGQSHPSLQKKLKKMLAAQKESARRPIFYRAQVG
hILiRa1S    62  VSKGEFCLYCDKDKGQSHPSLQKKLKKMLAAQKESARRPIFYRAQVG

hILiRa1      112  SNWMLESAAHPGWFICTSCNCNEPVGVTDKFENRKHIIESFQPVCKAE
hILiRa1L    152  SNWMLESAAHPGWFICTSCNCNEPVGVTDKFENRKHIIESFQPVCKAE
hILiRa1V    163  SNWMLESAAHPGWFICTSCNCNEPVGVTDKFENRKHIIESFQPVCKAE
hILiRa1S    112  SNWMLESAAHPGWFICTSCNCNEPVGVTDKFENRKHIIESFQPVCKAE

hILiRa1      162  PSEVSD
hILiRa1L    202  PSEVSD
hILiRa1V    213  PSEVSD
hILiRa1S    162  PSEVSD

```

[illegible]

Fig. 18

206120 99569860

1 GGCCTCGAG GCCAAGAAT CGGCACGAG CTTCATCCA TTTTGTGTTG AGTAATAAAC TCAAGTTGA AAATGCTTT TGTGGGGAG AACTCAGAG
 CGCGAGCTC CGGTCTTAA GCGCTGCTC GAAGTAAGT AAGAGAAC TCATTATTTG AGTTGCAJCT TTACAGBAA ACACCCCTTC TGTAGCTTC
 M S F V G E N S G V
 1
 101 TGAAATSG CTCGAGGAC TGGAAAGA AFGAACCCCA GTCTGCTTGA GAAGACCGG CTGGAAGCC CCGCAACCA GGCACCAAT TCCCACCAT
 ACTTTACC GAGACTGCTG ACCCTTTTC TACTTTTCT CACGAGGAT CTTCGCGC GACCTTCGG GACCTTGT CGGCTTCGG AGGGTGGA
 11 K M G S E D W E K D E P Q C C L E D P A G S P L E P G P S L P T H
 201 GAATTTGTTT CACACAGTC CAAGGTGAA GACTTAAC CCGAGGAAT TCACATCA TCACAGGAT CACAAATAC TGGTCTTGA CTCTGGGAAT
 CTTAAACAA GTGTGTCAG GTTTCACCT CTGGAATTTG GGTCTTTTAA AGTCGTAAGT ACTGTCCTA GTGTTTCAAG ACCAGACCT GAGACCTTA
 44 N F V H T S P K V K N L N P K K F S I H D Q D H K V L V L D S G N
 301 CTCATGACG TTCCAGATA AAACATACATA GGCACAGAGA TCTTCTTTGC ATTAGCTCA TCCTTGAGCT CAGCTCTGC GAGAAAGA AGTCGATTC
 GAGTATCTTC AAGTCTATT TTGTGATAT GCGGTCTCT AGAAGAAGC TATCGGAGT AGGAATCTGA GTGCGAGAG CTCTTTCT TCAGCTAAG
 77 L I A V P D K N Y I R P E I F F A L A S S L S S A S A E K G S P I L
 401 TCTGGGGT CTCTAAGGG GAGTTTGTG TCTACTGTGA CAAGATATA GGCATAAA GGCATAATC ATCCATCCT TCAGCTGAG AGGAGAAC TGATGAAGCT
 AGGACCCCA GAGATTTCC CTCAACAG AGATGACACT GTCTTATTT TGTGTTTCCAG TAGGTAGGA AGTCGACTTC TTCTCTTTG ACTACTGA
 111 L G V S K G E F C L Y C D K D K G Q S H P S L Q L K K E K L M K L
 501 GGTGCTCCAA AGGAATCAG CAGCGCGGC CTTCATCTTT TATAGGGCTC AGGTGGGCTC CTGGAACATG CTGAGTCGC CGGTCACCC CGATGTTTC
 CGACGGGTT TTCTTAGTC GTGCGCGCG GAGTAGAAA ATATCCCGAG TCACCCCGAG GACCTTGTAC GACCTCAGC GCGAGTGGG GCTTACCAAG
 144 A A Q K E S A R R P F I F Y R A Q V G S W N M L E S A A H P G W F
 601 ATCTCACT CTGCAATG TAAAGGCT CTGCGGTGA CAGTAAAT TGAAGACAG AAGACATG AATTTCAIT TCAACCAATT TCAAAAGCTG
 TGAAGTGA GAGCTTAC ATTACTGGA CAACCCACT GTCTATTTAA TTTGTGTAAC TTAAGTAA AGTTGTCAA AGTTTCAG
 177 I C T S C N C N E P V G V T D K F E N R K H I E F S F Q P V C K A E
 701 AAATGAGCC CAGTGGCT AGCATAGG AACTGCCCC ATTGAAGCC TTCTCGTA ATTGACTA ATTGTATAA AAGACCAAC CTGCTACTA
 TTACTCGG GTACTCCAG TCCTATTC TTTGACGGG TAACTTCGG AAGAGGAT TAACTGAT TAACTTT TGTGTTG GACGAGTAT
 211 M S P S E V S D O
 801 AAAAAAAA AAAAAAAGT TTGCGCGCGC AGCTTAT
 TTTTTTTTT TTTTTTGA AAGCGCGG TCGAATA

Fig. 19

COMBINED DECLARATION FOR PATENT APPLICATION AND POWER OF ATTORNEY

As a below named inventor, I hereby declare that:

My residence, post office and citizenship are as stated below next to my name,

I believe I am the original and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled:

IL-1 RELATED POLYPEPTIDES

the specification of which (check only one item below):

☐ is attached hereto.

☐ was filed as United States application Serial No. _____ on _____ and was amended on _____ (if applicable.).

☒ was filed as PCT international application Number PCT/US99/30720 on 22 December 1999 and was amended under PCT Article 19 on () (if applicable) (now assigned as U.S.S.N 09/869,566, filed June 29, 2001).

I hereby state that I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose information which is material to the examination of this application in accordance with Title 37, Code of Federal Regulations, §1.56(a).

I hereby claim foreign priority benefits under Title 35, United States Code, §119 of any foreign application(s) for patent or inventor's certificate or of any PCT international application(s) designating at least one country other than the United States of America listed below and have also identified below any foreign application(s) for patent or inventor's certificate or any PCT international application(s) designating at least one country other than the United States of America filed by me on the same subject matter having a filing date before that of the application(s) of which priority is claimed.

PRIOR FOREIGN/PCT APPLICATION/U.S. PROVISIONAL APPLICATIONS WHICH CLAIMS PRIORITY UNDER 35 U.S.C. 119:

COUNTRY	APPLICATION NUMBER	DATE OF FILING (day, month, year)	PRIORITY CLAIMED UNDER 35 USC 119
US	60/113,430	23 December 1998	
US	60/116,843	22 January 1999	
US	60/129,122	13 April 1999	

I hereby claim the benefit under Title 35, United States Code, §120 of any United States application(s) or PCT international application(s) designating the United States of America that is/are listed below and, insofar as the subject matter of such of the claims of this application is not disclosed in that/those prior application(s) in the manner provided by the first paragraph of Title 35, United States Code §112, I acknowledge the duty to disclose material information a defined in Title 37, Code of Federal Regulations, §1.56(a) which occurred between the filing date of the prior application(s) and the national or PCT international filing date of this application:

COMBINED DECLARATION FOR PATENT APPLICATION AND POWER OF ATTORNEY (Continued)
PRIOR U.S. APPLICATIONS OR PCT INTERNATIONAL APPLICATIONS DESIGNATING
THE U.S. FOR BENEFIT UNDER 35 U.S.C. 120:

U.S. APPLICATIONS		STATUS (check one)		
U.S. Application Number	U.S. Filing Date	Patented	Pending	Abandoned

PCT APPLICATIONS DESIGNATING THE U.S.

PCT Application No.	PCT Filing Date	U.S. Serial Numbers			
PCT/US99/30720	22 December 1999				

POWER OF ATTORNEY: As a named inventor, I hereby appoint the following attorney(s) and/or agent(s) to prosecute this application and transact all business in the Patent and Trademark Office connected therewith.



09157

PATENT TRADEMARK OFFICE

Send correspondence to

Genentech, Inc.
 Attn: Craig G. Svoboda
 1 DNA Way
 South San Francisco, CA 94080-4990
 Telephone: (650) 225-1489

I hereby declare that all statements made herein of my own knowledge and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment or both, under Section 1001 of Title 18 of the United States Code and that willful false statements may jeopardize the validity of the application or any patent issue thereon.

The undersigned hereby authorizes the U.S. attorney or agent named herein at accept and follow instructions from his foreign patent agent as to any action to be taken in the Patent and Trademark Office regarding this application without direct communication between the U.S. attorney or agent and the undersigned. In the event of a change in the persons from whom instructions may be taken, the U.S. attorney or agent named herein will be so notified by the undersigned.

Full name of sole or first inventor

Audrey Goddard

Inventor's Name

Inventor's signature

Date

110 Congo Street, San Francisco, California 94131

Residence

Canada

Citizenship

1 DNA Way, South San Francisco, CA 94080-4990

Post Office Address

Full name of second joint inventor, if any

Guohua James Pan

Inventor's Name

Inventor's signature

Date

14 Tewsley Place, Etobicoke, Ontario, Canada, M9P 1N7

Residence

Canada

Citizenship

1 DNA Way, South San Francisco, CA 94080-4990

Post Office Address

00000566-074902

COMBINED DECLARATION FOR PATENT APPLICATION AND POWER OF ATTORNEY

As a below named inventor, I hereby declare that:

My residence, post office and citizenship are as stated below next to my name,

I believe I am the original and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled:

IL-1 RELATED POLYPEPTIDES

the specification of which (check only one item below):

☐ is attached hereto.

☐ was filed as United States application Serial No. _____ on _____ and was amended on _____ (if applicable.).

☒ was filed as PCT international application Number PCT/US99/30720 on 22 December 1999 and was amended under PCT Article 19 on () (if applicable) (now assigned as U.S.S.N 09/869,566, filed June 29, 2001).

I hereby state that I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose information which is material to the examination of this application in accordance with Title 37, Code of Federal Regulations, §1.56(a).

I hereby claim foreign priority benefits under Title 35, United States Code, 119 of any foreign application(s) for patent or inventor's certificate or of any PCT international application(s) designating at least one country other than the United States of America listed below and have also identified below any foreign application(s) for patent or inventor's certificate or any PCT international application(s) designating at least one country other than the United States of America filed by me on the same subject matter having a filing date before that of the application(s) of which priority is claimed.

PRIOR FOREIGN/PCT APPLICATION/U.S. PROVISIONAL APPLICATIONS WHICH CLAIMS PRIORITY UNDER 35 U.S.C. 119:

COUNTRY	APPLICATION NUMBER	DATE OF FILING (day, month, year)	PRIORITY CLAIMED UNDER 35 USC 119
US	60/113,430	23 December 1998	
US	60/116,843	22 January 1999	
US	60/129,122	13 April 1999	

I hereby claim the benefit under Title 35, United States Code, §120 of any United States application(s) or PCT international application(s) designating the United States of America that is/are listed below and, insofar as the subject matter of such of the claims of this application is not disclosed in that/those prior application(s) in the manner provided by the first paragraph of Title 35, United States Code §112, I acknowledge the duty to disclose material information as defined in Title 37, Code of Federal Regulations, §1.56(a) which occurred between the filing date of the prior application(s) and the national or PCT international filing date of this application:

COMBINED DECLARATION FOR PATENT APPLICATION AND POWER OF ATTORNEY (Continued)
PRIOR U.S. APPLICATIONS OR PCT INTERNATIONAL APPLICATIONS DESIGNATING
THE U.S. FOR BENEFIT UNDER 35 U.S.C. 120:

U.S. APPLICATIONS		STATUS (check one)		
U.S. Application Number	U.S. Filing Date	Patented	Pending	Abandoned

PCT APPLICATIONS DESIGNATING THE U.S.

PCT Application No.	PCT Filing Date	U.S. Serial Numbers			
PCT/US99/30720	22 December 1999				

POWER OF ATTORNEY: As a named inventor, I hereby appoint the following attorney(s) and/or agent(s) to prosecute this application and transact all business in the Patent and Trademark Office connected therewith.



Send correspondence to

Genentech, Inc.
 Attn: Craig G. Svoboda
 1 DNA Way
 South San Francisco, CA 94080-4990
 Telephone: (650) 225-1489

I hereby declare that all statements made herein of my own knowledge and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment or both, under Section 1001 of Title 18 of the United States Code and that willful false statements may jeopardize the validity of the application or any patent issue thereon.

The undersigned hereby authorizes the U.S. attorney or agent named herein to accept and follow instructions from his foreign patent agent as to any action to be taken in the Patent and Trademark Office regarding this application without direct communication between the U.S. attorney or agent and the undersigned. In the event of a change in the persons from whom instructions may be taken, the U.S. attorney or agent named herein will be so notified by the undersigned.

Full name of sole or first inventor

Audrey Goddard
 Inventor's Name

A. Goddard
 Inventor's signature

2/5/02
 Date

110 Congo Street, San Francisco, California 94131
 Residence

Canada
 Citizenship

1 DNA Way, South San Francisco, CA 94080-4990
 Post Office Address

Full name of second joint inventor, if any

Guohua James Pan

Inventor's Name

Inventor's signature

Date

14 Tewsley Place, Etobicoke, Ontario, Canada, M9P 1N7

Residence

Canada

Citizenship

1 DNA Way, South San Francisco, CA 94080-4990

Post Office Address

09369566-021902

#5

Sequence Listing

<110> Goddard, Audrey D.
Fan, Guohua James

<120> IL-1 Related Polypeptides

<130> P2534-3US

<140> US 09/869,566

<141> 2001-06-29

<150> US 60/113,430

<151> 1998-12-23

<150> US 60/116,843

<151> 1999-01-22

<150> US 60/129,122

<151> 1999-04-13

<150> PCT/US99/30720

<151> 1999-12-22

<160> 32

<210> 1

<211> 1006

<212> DNA

<213> Homo sapiens

<400> 1

ggcacgaggc aagccttcca ggttatcgtg acgcaccttg aaagtctgag 50
agctactgcc ctacagaaag ttactagtgc cctaaagctg gcgctggcac 100
tgatgttact gctgctgttg gagtacaact tcctataga aaacaactgc 150
cagcacctta agaccactca caccttcaga gtggccttga gaaagatttg 200
gggtcaagga tcatgagcga gaacaccact taagaggata gtgaactagt 250
ctgcatgtga gacgctgaga tcctatgtca ggctgtgata ggaggggaaac 300
agaaacccaaa ggaaagaaca gctttaagaa gcgcttaaga gccaccacc 350
cattcttgac agtcaactgc ccagcctggg ggcccoctgtt ctttatcaaa 400
caagtgcctg agctctttgc agagggtcaa aggtgaagaa cttaaacccg 450
aagaaattca gcattcatga ccaggatcac aaagtactgg tcctggactc 500
tggaatctc atagcagttc cagataaaaa ctacatacgc ccagagatct 550
tctttgcatt agcctcatcc ttgagctcag cctctgcgga gaaaggaagt 600
ccgattctcc tgggggtctc taaaggggag ttttgtctct actgtgacaa 650
ggataaagga caaagtcac ccatccctca gctgaagaag gagaactga 700
tgaagctggc tgcccaaaag gaatcagcac gccggccctt catcttttat 750

agggtcagg tgggtctctg gaacatgctg gagtcggcgg ctacccccgg 800
 atggttcacg tgcacctctt gcaattgtaa tgagcctgtt ggggtgacag 850
 ataaatttga gaacaggaaa cacattgaat tticatttca accagtttgc 900
 aaagctgaaa tgagccccag tgaggtcagc gattaggaaa ctgccccatt 950
 gaagccttc ctgcctaatt tgaactaatt gtataaaaac accaaacctg 1000
 ctact 1006

<210> 2
 <211> 26
 <212> PRT
 <213> Homo sapiens

<400> 2
 Met Leu Leu Leu Leu Glu Tyr Asn Phe Pro Ile Glu Asn Asn
 1 5 10 15
 Cys Gln His Leu Lys Thr Thr His Thr Phe Arg
 20 25

<210> 3
 <211> 167
 <212> PRT
 <213> Homo sapiens

<400> 3
 Val Lys Asn Leu Asn Pro Lys Lys Phe Ser Ile His Asp Gln Asp
 1 5 10 15
 His Lys Val Leu Val Leu Asp Ser Gly Asn Leu Ile Ala Val Pro
 20 25 30
 Asp Lys Asn Tyr Ile Arg Pro Glu Ile Phe Phe Ala Leu Ala Ser
 35 40 45
 Ser Leu Ser Ser Ala Ser Ala Glu Lys Gly Ser Pro Ile Leu Leu
 50 55 60
 Gly Val Ser Lys Gly Glu Phe Cys Leu Tyr Cys Asp Lys Asp Lys
 65 70 75
 Gly Gln Ser His Pro Ser Leu Gln Leu Lys Lys Glu Lys Leu Met
 80 85 90
 Lys Leu Ala Ala Gln Lys Glu Ser Ala Arg Arg Pro Phe Ile Phe
 95 100 105
 Tyr Arg Ala Gln Val Gly Ser Trp Asn Met Leu Glu Ser Ala Ala
 110 115 120
 His Pro Gly Trp Phe Ile Cys Thr Ser Cys Asn Cys Asn Glu Pro
 125 130 135
 Val Gly Val Thr Asp Lys Phe Glu Asn Arg Lys His Ile Glu Phe
 140 145 150

Ser Phe Gln Pro Val Cys Lys Ala Glu Met Ser Pro Ser Glu Val
155 160 165

Ser Asp

<210> 4
<211> 650
<212> DNA
<213> Artificial Sequence

<220>
<223> recombinant DNA

<400> 4
taattcacca tgtctgcact tctgatccta gctcttggtg gagctgcagt 50
tgctgactac aaagacgatg acgacaagct tgcggccgcg aattcagctc 100
tttgcagagg tccaaagggt aagaacttaa acccgaagaa attcagcatt 150
catgaccagg atcacaaagt actggctcctg gactctggga atctcatagc 200
agttccagat aaaaactaca tacgcccaga gatcttcttt gcattagcct 250
catccttgag ctacgctctt gcggagaaag gaagtccgat tctcctgggg 300
gtctctaaag gggagttttg tctctactgt gacaaggata aaggacaaag 350
tcatccatcc ctacagctga agaaggagaa actgatgaag ctggctgcc 400
aaaaggaatc agcacgccgg ccttcatct tttatagggc tcagggtggg 450
tcttggaaca tgctggagtc ggcggctcac ccggatggt tcatctgcac 500
ctctgcaat tgtaatgagc ctggtggggg gacagataaa ttgagaaca 550
ggaaacacat tgaattttca ttcaaccag ttgcaaagc tgaatgagc 600
cccagtggg tcagcgatta gggaccagt cgactctaga ggatcccg 650

<210> 5
<211> 203
<212> PRT
<213> Artificial Sequence

<220>
<223> recombinant protein

<400> 5
Met Ser Ala Leu Leu Ile Leu Ala Leu Val Gly Ala Ala Val Ala
1 5 10 15
Asp Tyr Lys Asp Asp Asp Asp Lys Leu Ala Ala Ala Asn Ser Ala
20 25 30
Leu Cys Arg Gly Pro Lys Val Lys Asn Leu Asn Pro Lys Lys Phe
35 40 45
Ser Ile His Asp Gln Asp His Lys Val Leu Val Leu Asp Ser Gly
50 55 60

Asn	Leu	Ile	Ala	Val	Pro	Asp	Lys	Asn	Tyr	Ile	Arg	Pro	Glu	Ile	
				65					70					75	
Phe	Phe	Ala	Leu	Ala	Ser	Ser	Leu	Ser	Ser	Ala	Ser	Ala	Glu	Lys	
				80					85					90	
Gly	Ser	Pro	Ile	Leu	Leu	Gly	Val	Ser	Lys	Gly	Glu	Phe	Cys	Leu	
				95					100					105	
Tyr	Cys	Asp	Lys	Asp	Lys	Gly	Gln	Ser	His	Pro	Ser	Leu	Gln	Leu	
				110					115					120	
Lys	Lys	Glu	Lys	Leu	Met	Lys	Leu	Ala	Ala	Gln	Lys	Glu	Ser	Ala	
				125					130					135	
Arg	Arg	Pro	Phe	Ile	Phe	Tyr	Arg	Ala	Gln	Val	Gly	Ser	Trp	Asn	
				140					145					150	
Met	Leu	Glu	Ser	Ala	Ala	His	Pro	Gly	Trp	Phe	Ile	Cys	Thr	Ser	
				155					160					165	
Cys	Asn	Cys	Asn	Glu	Pro	Val	Gly	Val	Thr	Asp	Lys	Phe	Glu	Asn	
				170					175					180	
Arg	Lys	His	Ile	Glu	Phe	Ser	Phe	Gln	Pro	Val	Cys	Lys	Ala	Glu	
				185					190					195	
Met	Ser	Pro	Ser	Glu	Val	Ser	Asp								
				200											

<210> 6
 <211> 754
 <212> DNA
 <213> Homo sapiens

<400> 6
 ggcacgaggc aagccttcca ggttatcgtg acgcaccttg aaagtctgag 50
 agctactgcc ctacagaaag ttactagtgc cctaaagctg gcgctggcac 100
 tgatgttact gctgctgttg gagtacaact tccctataga aaacaactgc 150
 cagcacctta agaccactca caccttcaga gtgaagaact taaacccgaa 200
 gaaattcagc attcatgacc aggatcacaa agtactggtc ctggactctg 250
 ggaatctcat agcagttcca gataaaaact acatacgccc agagatcttc 300
 tttgcattag cctcatcctt gagctcagcc tctgcggaga aaggaagtcc 350
 gattctcctg ggggtctcta aaggggagtt ttgtctctac tgtgacaagg 400
 ataaaggaca aagtcattcca tcccttcagc tgaagaagga gaaactgatg 450
 aagctggctg cccaaaagga atcagcacgc cggcccttca tcttttatag 500
 ggctcaggtg ggctcctgga acatgctgga gtcggcggct caccocggat 550
 ggttcactcg cacctcctgc aattgtaatg agcctgttgg ggtgacagat 600

aaatttgaga acaggaaca cattgaattt tcatttcaac cagtttgcaa 650
 agctgaaatg agccccagtg aggtcagcga ttaggaaact gccccattga 700
 acgccttctt cgctaatttg aactaattgt ataaaaacac caaacctgct 750
 cact 754

<210> 7
 <211> 193
 <212> PRT
 <213> Homo sapiens

<400> 7
 Met Leu Leu Leu Leu Leu Glu Tyr Asn Phe Pro Ile Glu Asn Asn
 1 5 10 15
 Cys Gln His Leu Lys Thr Thr His Thr Phe Arg Val Lys Asn Leu
 20 25 30
 Asn Pro Lys Lys Phe Ser Ile His Asp Gln Asp His Lys Val Leu
 35 40 45
 Val Leu Asp Ser Gly Asn Leu Ile Ala Val Pro Asp Lys Asn Tyr
 50 55 60
 Ile Arg Pro Glu Ile Phe Phe Ala Leu Ala Ser Ser Leu Ser Ser
 65 70 75
 Ala Ser Ala Glu Lys Gly Ser Pro Ile Leu Leu Gly Val Ser Lys
 80 85 90
 Gly Glu Phe Cys Leu Tyr Cys Asp Lys Asp Lys Gly Gln Ser His
 95 100 105
 Pro Ser Leu Gln Leu Lys Lys Glu Lys Leu Met Lys Leu Ala Ala
 110 115 120
 Gln Lys Glu Ser Ala Arg Arg Pro Phe Ile Phe Tyr Arg Ala Gln
 125 130 135
 Val Gly Ser Trp Asn Met Leu Glu Ser Ala Ala His Pro Gly Trp
 140 145 150
 Phe Ile Cys Thr Ser Cys Asn Cys Asn Glu Pro Val Gly Val Thr
 155 160 165
 Asp Lys Phe Glu Asn Arg Lys His Ile Glu Phe Ser Phe Gln Pro
 170 175 180
 Val Cys Lys Ala Glu Met Ser Pro Ser Glu Val Ser Asp
 185 190

<210> 8
 <211> 629
 <212> DNA
 <213> Homo sapiens

<220>
 <221> unsure
 <222> 13

09669566.021902

<223> unknown base

<400> 8

ccaggcccaa gntccccac catgaatttt gttcacacaa gtcgaaaggt 50
 gaagagctta aacccgaaga aattcagcat tcattgaccag gatcacaaag 100
 tactggcctg gactctggga atctcatagc agttccagat aaaaactaca 150
 tacgcccaga gatcttcttt gcattagcct catccttgag ctgagcctct 200
 gcgagaaag gaagtcgat tctcctggg gtctctaaag gggagttttg 250
 tctctactgt gacaaggata aaggacaaag tcattcatcc cttagctga 300
 agaaggagaa actgatgaag ctggctgcc aaaaggaatc agcacgcgg 350
 ccttcatct tttatagggc tcagtgggc tcttggaaca tgctggagtc 400
 ggcggtcac ccggatggt tcattgcac ctctgcaat tgtaatgagc 450
 ctgttggtg gacagataaa ttgagaaca ggaacacat tgaattttca 500
 tttcaaccag tttgcaaagc tgaaatgagc ccagtgagg tcagcgatta 550
 ggaactgcc ccattgaacg ccttctcgc taatttgaac taattgtata 600
 aaaaacccaa acctgctcac taacaaaaa 629

<210> 9

<211> 1321

<212> DNA

<213> Homo sapiens

<400> 9

gtcgacccac gcgtccgaag ctgtggagc cagattcag tcccctggac 50
 tgtagataaa gacctttct tgccaggtgc tgagacaacc acactatgag 100
 aggcactcca ggagacgctg atgttgagg aagggccgtc tatcaatcaa 150
 tcaactgttc tgttatcaca tgcaagtatc cagaggctct tgagcaaggc 200
 agaggggato ccatttattt gggaatccag aatccagaaa tgtgtttgta 250
 ttgtgagaag gttggagaac agcccacatt gcagctaaaa gaggagaaga 300
 tcattgatct gtatggccaa cccagcccg tgaaccctt cctttttac 350
 cgtgccaaga ctggtaggac ctccaccctt gactctgtg ccttcccggg 400
 ctgttcatc gctcctcca agagagacca gccatcatt ctgacttcag 450
 aacttgaggaa gtcatacaac actgccttg aattaaatat aaatgactga 500
 actcagccta gagggtgcag ctgtgtctt gtcttaaatg ttctggttcc 550
 caatgtgtt tcgtctacat ttcttagtg toattttcac gctgtgctg 600
 agacaggagc aggtgtctg ttatcatctc attttataat gaagaagaag 650

caattacttc atagcaactg aagaacagga tgtggcctca gaagcaggag 700
 agctgggtgg tataaggctg tcctctcaag ctgggtctgt gtaggccaca 750
 aggcattctg atgagtact ttaagaacta aagaccaaac actgagcttt 800
 cttctagggg tgggtatgaa gatgcttcag agctcatgag cgttaccac 850
 gatggcatga ctagcacaga gctgatctct gttctgttt tgetttattc 900
 cctcttggga tgatatcatc cagtctttat atgttgccaa tatacctcat 950
 tgtgtgtaat agaaccttct tagcattaag acctgttaa caaaaaaat 1000
 tcttgggggt ggtatgaaga tgcttcagag ctcatgcgag ttaccacaga 1050
 tggcatgact agcacagagc tgatctctgt tctgtttttg ctttattccc 1100
 tcttgggatg atatcatcca gtctttatat gttgccaata tacctcattg 1150
 tgtgtaatag aacctcttta gcattaagac ctgttaaaca aaaataattc 1200
 ttgtgttaag ttaaatcatt ttgtctctaa ttgtaatgtg taatcttaaa 1250
 gttaaataaa ctttgtgtat ttatataata ataaagctaa aactgatata 1300
 aaetaaagaa agagtaaact g 1321

<210> 10
 <211> 134
 <212> PRT
 <213> Homo sapiens

<400> 10
 Met Arg Gly Thr Pro Gly Asp Ala Asp Gly Gly Gly Arg Ala Val
 1 5 10 15
 Tyr Gln Ser Ile Thr Val Ala Val Ile Thr Cys Lys Tyr Pro Glu
 20 25 30
 Ala Leu Glu Gln Gly Arg Gly Asp Pro Ile Tyr Leu Gly Ile Gln
 35 40 45
 Asn Pro Glu Met Cys Leu Tyr Cys Glu Lys Val Gly Glu Gln Pro
 50 55 60
 Thr Leu Gln Leu Lys Glu Gln Lys Ile Met Asp Leu Tyr Gly Gln
 65 70 75
 Pro Glu Pro Val Lys Pro Phe Leu Phe Tyr Arg Ala Lys Thr Gly
 80 85 90
 Arg Thr Ser Thr Leu Glu Ser Val Ala Phe Pro Asp Trp Phe Ile
 95 100 105
 Ala Ser Ser Lys Arg Asp Gln Pro Ile Ile Leu Thr Ser Glu Leu
 110 115 120
 Gly Lys Ser Tyr Asn Thr Ala Phe Glu Leu Asn Ile Asn Asp
 125 130

<210> 11
 <211> 249
 <212> DNA
 <213> Homo sapiens

<400> 11
 aagctgctgg agccacgatt cagtcccctg gactgtagat aaagaccctt 50
 tcttgccagg tgctgagaca accacactat gagaggcact ccaggagacg 100
 ctgatggtgg aggaagggcc gtctatcaat caatcaactgt tgctgtttatc 150
 acatgcaagt atccagaggc tcttgagcaa ggcagagggg atcccattta 200
 ttgggaatc cagaatccag aaatgtgttt gtattgtgag aaggttgga 249

<210> 12
 <211> 468
 <212> DNA
 <213> Homo sapiens

<400> 12
 atggtcctga gtggggcgct gtgcttccga atgaaggact cggcattgaa 50
 ggtgctttat ctgcataata accagcttct agctggaggg ctgcatgcag 100
 ggaaggtcat taaaggtgaa gagatcagcg tggctcccaa tcggtggctg 150
 gatgccagcc tgtccccctg catcctgggt gtccagggtg gaagccagtg 200
 cctgtcatgt ggggtggggc aggagccgac tctaacacta gagccagtga 250
 acatcatgga gctctatctt ggtgccaagg aatccaagag cttcaccttc 300
 taccggcggg acatggggct cacctccagc ttcgagtcgg ctgcctaccc 350
 gggctgggtc ctgtgcacgg tgccotgaagc cgatcagcct gtcagactca 400
 cccagcttcc cgaqaatggt ggctggaatg cccccatcac agactttcac 450
 ttccagcagt gtgactag 468

<210> 13
 <211> 155
 <212> PRT
 <213> Homo sapiens

<400> 13
 Met Val Leu Ser Gly Ala Leu Cys Phe Arg Met Lys Asp Ser Ala
 1 5 10 15
 Leu Lys Val Leu Tyr Leu His Asn Asn Gln Leu Leu Ala Gly Gly
 20 25 30
 Leu His Ala Gly Lys Val Ile Lys Gly Glu Glu Ile Ser Val Val
 35 40 45
 Pro Asn Arg Trp Leu Asp Ala Ser Leu Ser Pro Val Ile Leu Gly
 50 55 60
 Val Gln Gly Gly Ser Gln Cys Leu Ser Cys Gly Val Gly Gln Glu

09869566 121002

65	70	75
Pro Thr Leu Thr Leu Glu Pro Val Asn Ile Met Glu Leu Tyr Leu		
80	85	90
Gly Ala Lys Glu Ser Lys Ser Phe Thr Phe Tyr Arg Arg Asp Met		
95	100	105
Gly Leu Thr Ser Ser Phe Glu Ser Ala Ala Tyr Pro Gly Trp Phe		
110	115	120
Leu Cys Thr Val Pro Glu Ala Asp Gln Pro Val Arg Leu Thr Gln		
125	130	135
Leu Pro Glu Asn Gly Gly Trp Asn Ala Pro Ile Thr Asp Phe Tyr		
140	145	150
Phe Gln Gln Cys Asp		
155		

<210> 14
 <211> 295
 <212> DNA
 <213> Homo sapiens

<220>
 <221> unsure
 <222> 283
 <223> unknown base

<400> 14
 gctcccgccgga ggagaaagga acattctgag gggagtctac accctgtgga 50
 gctcaagatg gtcctgagtg gggcgctgtg cttcogaatg aaggactcgg 100
 cattgaaggt gctttatctg cataataacc agcttctagc tggagggctg 150
 catgcaggga aggtcattaa aggtgaagag atcagcgtgg tcccgaatcg 200
 gtggctggat gccagcctgt ccccgctcat cctgggtgtc cagggtggaa 250
 gccagtgcct gtcatgtggg gtggggcagg agnccgactct aacat 295

<210> 15
 <211> 1385
 <212> DNA
 <213> Mus musculus

<400> 15
 atagggaatt tggccctcga ggccaagaat tcggcacgag gggagcctgc 50
 ttctactta ggtctcaaat ttccagcct tgtctttgcc taaaatttcc 100
 tgctgtttat ttcaaaatag ggtctacata ctgtggagct catgatggtt 150
 ctgagtgggg cactatgctt ccgaatgaag gattcagcct tgaagggtact 200
 gtatctgcac aataaccagc tgctggctgg aggactgcac gcagagaagg 250
 tcattaaagg tgaggagatc agtgttgtcc caaatcgggc actggatgcc 300

agtctgtccc ctgtcatcct gggcgttcaa ggaggaagcc agtgccatc 350
 ttgtgggaca gagaagggc caattctgaa acttgagcca gtgaacatca 400
 tggagctcta cctcggggcc aaggaatcaa agagcttcac cttctaccgg 450
 cgggatatgg gtcttacctc cagcttcgaa tccgctgcct acccaggctg 500
 gttctctgc acctaccgg aagctgacca gccgtgcagg ctcaactcaga 550
 tccctgagga ccccgctgg gatgctccca tcacagactt ctactttcag 600
 cagtgtgact agggctgcgt ggtcccaaa actccataag cagaggcaga 650
 gtaggcagtg gcggctcctg atagaggata gagagacaga ggagctccac 700
 agtaggtggc ttactcctct ccttcctac tggactcccg cttctgacct 750
 aaggcacaca gacactctct tctcctgcat ccagtgctg gtaaatcttc 800
 tggattttgg agctcaatgt gtagattott tcagattgga tggactacc 850
 tctggtgtgg aaccaatag aaaccacgta ggaccaaaa agagcaacat 900
 aaaagattct tgggtgaaga agaggtggga actgttcata catagtaaga 950
 tctgacacag tacctcagaa gtccctgcat tccttatgtt ctggagaaa 1000
 tggagggggg gtcaccaaga ctttctctgg ctggctgggg ctttccctc 1050
 aacctttctg acatctgcag cctctctcat tcttgccttc attctctggc 1100
 cctgaaccga gaggggtgata tcaggatagc tgacagaaga tgaccaggca 1150
 cactgtcctg gtttgaacc agaggggaca ataaaaaacc ctgattctgg 1200
 tctctactca cataaaaaga agcttgtgaa cattaagtg gaagagattg 1250
 ctactaaata acataccttg taatttcac ttaattaaaa tatacttctc 1300
 tatattatat attttaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 1350
 aaaaacatgc ggcgcgaagc ttattccatt tagga 1385

<210> 16

<211> 155

<212> PRT

<213> Mus musculus

<400> 16

Met	Val	Leu	Ser	Gly	Ala	Leu	Cys	Phe	Arg	Met	Lys	Asp	Ser	Ala
	1				5				10				15	

Leu	Lys	Val	Leu	Tyr	Leu	His	Asn	Asn	Gln	Leu	Leu	Ala	Gly	Gly
			20						25				30	

Leu	His	Ala	Glu	Lys	Val	Ile	Lys	Gly	Glu	Glu	Ile	Ser	Val	Val
			35						40				45	

Pro	Asn	Arg	Ala	Leu	Asp	Ala	Ser	Leu	Ser	Pro	Val	Ile	Leu	Gly
				50					55				60	

Val Gln Gly Gly Ser Gln Cys Leu Ser Cys Gly Thr Glu Lys Gly
65 70 75

Pro Ile Leu Lys Leu Glu Pro Val Asn Ile Met Glu Leu Tyr Leu
80 85 90

Gly Ala Lys Glu Ser Lys Ser Phe Thr Phe Tyr Arg Arg Asp Met
95 100 105

Gly Leu Thr Ser Ser Phe Glu Ser Ala Ala Tyr Pro Gly Trp Phe
110 115 120

Leu Cys Thr Ser Pro Glu Ala Asp Gln Pro Val Arg Leu Thr Gln
125 130 135

Ile Pro Glu Asp Pro Ala Trp Asp Ala Pro Ile Thr Asp Phe Tyr
140 145 150

Phe Gln Gln Cys Asp
155

<210> 17
<211> 382
<212> DNA
<213> Mus musculus

<400> 17
ggagcctgct ttctacttag gtctcaaatt ttccagcctt gtctttgctt 50
aaaatttctt gctgtttatt tcaaaataggt gtctacatac tgtggagctc 100
atgatgggtc tgagtggggc actatgcttc cgaatgaagg attcagcctt 150
gaaggtagct tatctgcaca ataaccagct gctggctgga ggactgcacg 200
cagagaaggt cattaaaggt gaggagatca gtgttgctcc aaatcgggca 250
ctgqatqcca gtctgtcccc tgtcatcctg ggctttcaaa gaggaagcca 300
tgtcctatct tgtgggacag agaaagggcc aattctgaaa cttgagccag 350
tgaacatcat ggagctctac ctcggggcca ag 382

<210> 18
<211> 626
<212> DNA
<213> Homo sapiens

<400> 18
aaaatgggct ctgaggactg ggaaaaagat gaacccagct gctgcttaga 50
agacccggct gtaagcccc ttgaaccagg cccaagcctc cccgccatga 100
attttgttca cacaagtcca aaggtgaaga acttaaaccc gaagaaattc 150
agcattcatg accaggatca caaagtactg gtcctggact ctgggaatct 200
catagcagtt ccagataaaa actacatacg cccagagatc ttctttgcat 250
tagcctcacc cttgagctca gctctgcgg agaaaggaag tccgattctc 300

ctgggggtct ctaaagggga gttttgtctc tactgtgaca aggataaagg 350
 acaaagtcat ccattcccttc agctgaagaa ggagaaactg atgaagctgg 400
 ctgccccaaa ggaatcagca cgccggccct tcattcttta tagggctcag 450
 gtgggtctct ggaacatgct ggagtcggcg gctcaccocg gatggttcat 500
 ctgcacctcc tgcaattgta atgagcctgt tggggtgaca gataaatttg 550
 agaacaggaa acacattgaa ttttcatttc aaccagtttg caaagctgaa 600
 atgagcccca gtgaggtcag cgatta 626

<210> 19
 <211> 207
 <212> PRT
 <213> Homo sapiens

<400> 19
 Met Gly Ser Glu Asp Trp Glu Lys Asp Glu Pro Gln Cys Cys Leu
 1 5 10 15
 Glu Asp Pro Ala Val Ser Pro Leu Glu Pro Gly Pro Ser Leu Pro
 20 25 30
 Ala Met Asn Phe Val His Thr Ser Pro Lys Val Lys Asn Leu Asn
 35 40 45
 Pro Lys Lys Phe Ser Ile His Asp Gln Asp His Lys Val Leu Val
 50 55 60
 Leu Asp Ser Gly Asn Leu Ile Ala Val Pro Asp Lys Asn Tyr Ile
 65 70 75
 Arg Pro Glu Ile Phe Phe Ala Leu Ala Ser Ser Leu Ser Ser Ala
 80 85 90
 Ser Ala Glu Lys Gly Ser Pro Ile Leu Leu Gly Val Ser Lys Gly
 95 100 105
 Glu Phe Cys Leu Tyr Cys Asp Lys Asp Lys Gly Gln Ser His Pro
 110 115 120
 Ser Leu Gln Leu Lys Lys Glu Lys Leu Met Lys Leu Ala Ala Gln
 125 130 135
 Lys Glu Ser Ala Arg Arg Pro Phe Ile Phe Tyr Arg Ala Gln Val
 140 145 150
 Gly Ser Trp Asn Met Leu Glu Ser Ala Ala His Pro Gly Trp Phe
 155 160 165
 Ile Cys Thr Ser Cys Asn Cys Asn Glu Pro Val Gly Val Thr Asp
 170 175 180
 Lys Phe Glu Asn Arg Lys His Ile Glu Phe Ser Phe Gln Pro Val
 185 190 195
 Cys Lys Ala Glu Met Ser Pro Ser Glu Val Ser Asp

<210> 20
 <211> 506
 <212> DNA
 <213> Homo sapiens

<400> 20
 aaaatgggct ctgaggactg ggaaaaagat gaaccccgct gctgcttaga 50
 agaccgggct gtaagccccc tggaaccagg cccaagcctc cccgccatga 100
 attttgttca cacaaagatc ttctttgcat tagcctcatc cttgagctca 150
 gcctctcgcg agaaaggaag tccgattctc ctgggggtct ctaaaggggga 200
 gttttgtctc tactgtgaca aggataaagg acaaagtcac ccatcccttc 250
 agctgaagaa ggagaaactg atgaagctgg ctgcccaaaa ggaatcagca 300
 cgccggccct tcatctttta tagggctcag gtggggtcct ggaacatgct 350
 ggagtcggcg gctcaccocg gatggttcat ctgcacctcc tgcaattgta 400
 atgagcctgt tggggtgaca gataaatttg agaacaggaa acacattgaa 450
 ttttcatttc aaccagtttg caaagctgaa atgagcccca gtgaggtcag 500
 cgatta 506

<210> 21
 <211> 167
 <212> PRT
 <213> Homo sapiens

<400> 21
 Met Gly Ser Glu Asp Trp Glu Lys Asp Glu Pro Gln Cys Cys Leu
 1 5 10 15
 Glu Asp Pro Ala Val Ser Pro Leu Glu Pro Gly Pro Ser Leu Pro
 20 25 30
 Ala Met Asn Phe Val His Thr Lys Ile Phe Phe Ala Leu Ala Ser
 35 40 45
 Ser Leu Ser Ser Ala Ser Ala Glu Lys Gly Ser Pro Ile Leu Leu
 50 55 60
 Gly Val Ser Lys Gly Glu Phe Cys Leu Tyr Cys Asp Lys Asp Lys
 65 70 75
 Gly Gln Ser His Pro Ser Leu Gln Leu Lys Lys Glu Lys Leu Met
 80 85 90
 Lys Leu Ala Ala Gln Lys Glu Ser Ala Arg Arg Pro Phe Ile Phe
 95 100 105
 Tyr Arg Ala Gln Val Gly Ser Trp Asn Met Leu Glu Ser Ala Ala
 110 115 120
 His Pro Gly Trp Phe Ile Cys Thr Ser Cys Asn Cys Asn Glu Pro

	125		130		135
Val Gly Val Thr Asp Lys Phe Glu Asn Arg Lys His Ile Glu Phe					
	140		145		150
Ser Phe Gln Pro Val Cys Lys Ala Glu Met Ser Pro Ser Glu Val					
	155		160		165

Ser Asp

<210> 22
 <211> 561
 <212> DNA
 <213> Homo sapiens

<400> 22
 aacccgaaga aattcagcat tcatgaccag gatcacaaag tactggtcct 50
 ggactctggg aatctcatag cagtccaga taaaaactac atacgccag 100
 agatcttctt tgcattagcc tcatcctga gctcagcctc tggcgagaaa 150
 ggaagtccga ttctcctggg ggtctctaaa ggggagttt+ gtctctactg 200
 tgacaaggat aaaggacaaa gtcattccatc ccttcagctg aagaaggaga 250
 aactgatgaa gctggctgcc caaaaggaat cagcacgccg gcccttcac 300
 ttttataggg ctgagtgagg ctccctggaac atgctggagt cgcgcgctca 350
 ccccgatagg ttcattctga cctcctgcaa ttgtaattg cctgttgggg 400
 tgacagataa atttgagaac aggaaacaca ttgaatttct atttcaacca 450
 gtttgcaaa ctgaaatgag cccagtgag gtcagcgatt aggaaactgc 500
 cccattgaac gccctcctcg ctaatttgaa ctaattgtat aaaaacacca 550
 aacctgctca c 561

<210> 23
 <211> 561
 <212> DNA
 <213> Homo sapiens

<400> 23
 ttgggcttct ttaagtcgta agtactggtc ctagtgttct atgaccagga 50
 cctgagacct ttagagtatc gtcaaggctc atttttgatg tatcggggtc 100
 tctagaagaa acgtaatcgg agtaggaact cgagtcggag acgcctcttt 150
 ccttcaggct aagaggacct ccagagattt cccctcaaaa cagagatgac 200
 actgttctta ttctcgttt cagtaggtag ggaagtcgac ttcttctct 250
 ttgactactt cgaccgacgg gttttcctta gtcgtgcggc ogggaagtag 300
 aaaatatccc gagtccacct gaggaccttg tacgacctca gccgccgagt 350

ggggcctacc aagtagacgt ggaggacgtt aacattactc ggacaacccc 400
 actgtctatt taaactcttg tcctttgtgt aacttaaaag taaagtgggt 450
 caaacgttgc gactttactc ggggtcactc cagtcgctaa tcctttgacg 500
 gggttaacttg cggaaggagc gattaaactt gattaacata tttttgtgggt 550
 ttggacgagt g 561

<210> 24
 <211> 839
 <212> DNA
 <213> Homo sapiens

<400> 24
 ggccctcgag gccagaatt cggcacgagg ctctattcca ttttctgttg 50
 agtaataaac tcaacgttga aaatgtcctt tgtgggggag aactcaggag 100
 tgaaaatggg ctctgaggac tgggaaaaag atgaacccca gtgctgctta 150
 gaagaccogg ctggaagccc cctggaacca ggcccaagcc tccccaccat 200
 gaattttgtt cacacaagtc caaagggtgaa gaacttaaac ccgaagaagt 250
 tcagcattca tgaccaggat cacaaagtac tggctcctgga ctctgggaat 300
 ctcatagcag ttccagataa aaactacata cgcccagaga tcttctttgc 350
 attagcctca tccttgagct cagcctctgc ggagaaagga agtccgattc 400
 tcctgggggt ctctaagggt gagttttgtc tctactgtga caaggataaa 450
 ggacaaagtc atccatccct tcagctgaag aaggagaaac tgatgaagct 500
 ggtgcgccaa aaggaatcag caccgccggc cttcatcttt tatagggttc 550
 agtggggctc ctggaacatg ctggagtcgq cggctcaccg cggatggttc 600
 atctgcacct cctgcaattg taatgagcct gttgggggtga cgataaatt 650
 tgagaacagg aaacacattg aattttcatt tcaaccagtt tgcaaagctg 700
 aaatgagccc cagtgggttc agcgattagg aaactgcccc attgaacgcc 750
 ttctcgcta attgaaacta attgtataaa aacaccaaac ctgctcacta 800
 aaaaaaaaaa aaaaaaacgt ttgcggccgc aagcttatt 839

<210> 25
 <211> 218
 <212> PRT
 <213> Homo sapiens

<400> 25
 Met Ser Phe Val Gly Glu Asn Ser Gly Val Lys Met Gly Ser Glu
 1 5 10 15
 Asp Trp Glu Lys Asp Glu Pro Gln Cys Cys Leu Glu Asp Pro Ala
 20 25 30

Gly Ser Pro Leu Glu Pro Gly Pro Ser Leu Pro Thr Met Asn Phe
 35 40 45
 Val His Thr Ser Pro Lys Val Lys Asn Leu Asn Pro Lys Lys Phe
 50 55 60
 Ser Ile His Asp Gln Asp His Lys Val Leu Val Leu Asp Ser Gly
 65 70 75
 Asn Leu Ile Ala Val Pro Asp Lys Asn Tyr Ile Arg Pro Glu Ile
 80 85 90
 Phe Phe Ala Leu Ala Ser Ser Leu Ser Ser Ala Ser Ala Glu Lys
 95 100 105
 Gly Ser Pro Ile Leu Leu Gly Val Ser Lys Gly Glu Phe Cys Leu
 110 115 120
 Tyr Cys Asp Lys Asp Lys Gly Gln Ser His Pro Ser Leu Gln Leu
 125 130 135
 Lys Lys Glu Lys Leu Met Lys Leu Ala Ala Gln Lys Glu Ser Ala
 140 145 150
 Arg Arg Pro Phe Ile Phe Tyr Arg Ala Gln Val Gly Ser Trp Asn
 155 160 165
 Met Leu Glu Ser Ala Ala His Pro Gly Trp Phe Ile Cys Thr Ser
 170 175 180
 Cys Asn Cys Asn Glu Pro Val Gly Val Thr Asp Lys Phe Glu Asn
 185 190 195
 Arg Lys His Ile Glu Phe Ser Phe Gln Pro Val Cys Lys Ala Glu
 200 205 210
 Met Ser Pro Ser Glu Val Ser Asp
 215

<210> 26

<211> 177

<212> PRT

<213> Homo sapiens

<400> 26

Met Glu Ile Cys Arg Gly Leu Arg Ser His Leu Ile Thr Leu Leu
 1 5 10 15
 Leu Phe Leu Phe His Ser Glu Thr Ile Cys Arg Pro Ser Gly Arg
 20 25 30
 Lys Ser Ser Lys Met Gln Ala Phe Arg Ile Trp Asp Val Asn Gln
 35 40 45
 Lys Thr Phe Tyr Leu Arg Asn Asn Gln Leu Val Ala Gly Tyr Leu
 50 55 60
 Gln Gly Pro Asn Val Asn Leu Glu Glu Lys Ile Asp Val Val Pro
 65 70 75

Ile Glu Pro His Ala Leu Phe Leu Gly Ile His Gly Gly Lys Met
80 85 90
Cys Leu Ser Cys Val Lys Ser Gly Asp Glu Thr Arg Leu Gln Leu
95 100 105
Glu Ala Val Asn Ile Thr Asp Leu Ser Glu Asn Arg Lys Gln Asp
110 115 120
Lys Arg Phe Ala Phe Ile Arg Ser Asp Ser Gly Pro Thr Thr Ser
125 130 135
Phe Glu Ser Ala Ala Cys Pro Gly Trp Phe Leu Cys Thr Ala Met
140 145 150
Glu Ala Asp Gln Pro Val Ser Leu Thr Asn Met Pro Asp Glu Gly
155 160 165
Val Met Val Thr Lys Phe Tyr Phe Gln Glu Asp Glu
170 175

<210> 27
<211> 169
<212> PRT
<213> Homo sapiens

<400> 27
Met Arg Gly Thr Pro Gly Asp Ala Asp Gly Gly Gly Arg Ala Val
1 5 10 15
Tyr Gln Ser Met Cys Lys Pro Ile Thr Gly Thr Ile Asn Asp Leu
20 25 30
Asn Gln Gln Val Trp Thr Leu Gln Gly Gln Asn Leu Val Ala Val
35 40 45
Pro Arg Ser Asp Ser Val Thr Pro Val Thr Val Ala Val Ile Thr
50 55 60
Cys Lys Tyr Pro Glu Ala Leu Glu Gln Gly Arg Gly Asp Pro Ile
65 70 75
Tyr Leu Gly Ile Gln Asn Pro Glu Met Cys Leu Tyr Cys Glu Lys
80 85 90
Val Gly Glu Gln Pro Thr Leu Gln Leu Lys Glu Gln Lys Ile Met
95 100 105
Asp Leu Tyr Gly Gln Pro Glu Pro Val Lys Pro Phe Leu Phe Tyr
110 115 120
Arg Ala Lys Thr Gly Arg Thr Ser Thr Leu Glu Ser Val Ala Phe
125 130 135
Pro Asp Trp Phe Ile Ala Ser Ser Lys Arg Asp Gln Pro Ile Ile
140 145 150
Leu Thr Ser Glu Leu Gly Lys Ser Tyr Asn Thr Ala Phe Glu Leu
155 160 165
Asn Ile Asn Asp

<210> 28
 <211> 167
 <212> PRT
 <213> Homo sapiens

<400> 28
 Met Gly Ser Glu Asp Trp Glu Lys Asp Glu Pro Gln Cys Cys Leu
 1 5 10 15
 Glu Asp Pro Ala Gly Ser Pro Leu Glu Pro Gly Pro Ser Leu Pro
 20 25 30
 Thr Met Asn Phe Val His Thr Lys Ile Phe Phe Ala Leu Ala Ser
 35 40 45
 Ser Leu Ser Ser Ala Ser Ala Glu Lys Gly Ser Pro Ile Leu Leu
 50 55 60
 Gly Val Ser Lys Gly Glu Phe Cys Leu Tyr Cys Asp Lys Asp Lys
 65 70 75
 Gly Gln Ser His Pro Ser Leu Gln Leu Lys Lys Glu Lys Leu Met
 80 85 90
 Lys Leu Ala Ala Gln Lys Glu Ser Ala Arg Arg Pro Phe Ile Phe
 95 100 105
 Tyr Arg Ala Gln Val Gly Ser Trp Asn Met Leu Glu Ser Ala Ala
 110 115 120
 His Pro Gly Trp Phe Ile Cys Thr Ser Cys Asn Cys Asn Glu Pro
 125 130 135
 Val Gly Val Thr Asp Lys Phe Glu Asn Arg Lys His Ile Glu Phe
 140 145 150
 Ser Phe Gln Pro Val Cys Lys Ala Glu Met Ser Pro Ser Glu Val
 155 160 165
 Ser Asp

<210> 29
 <211> 31
 <212> DNA
 <213> Homo sapiens

<400> 29
 ggcggaatcca aaatgggctc tgaggactgg g 31

<210> 30
 <211> 30
 <212> DNA
 <213> Homo sapiens

<400> 30
 gcggaattct aatcgctgac ctcaactgggg 30

<210> 31

<211> 9
<212> PRT
<213> Homo sapiens

<400> 31
Val His Thr Ser Pro Lys Val Lys Asn
1 5

<210> 32
<211> 10
<212> PRT
<213> Homo sapiens

<400> 32
Val Leu Ser Gly Ala Leu Cys Phe Arg Met
1 5 10

09869566.021902